

BOSTON
MEDICAL LIBRARY



IN THE
Francis A. Countway
Library of Medicine
BOSTON

Gift of

New England Journal of Medicine

Symp.
E
161 I
89
621—

Basic Aspects

DEDICATION

We dedicate this book to the Organizing Committee: E. Johannison, P. Melzer, H.S. Jacobs, P.C. Sizonenko, F.H. Schroder and F. Comite and to all our colleagues whose work with GnRH analogues is increasing the quality of life for individuals suffering the wide range of medical conditions benefited by the use of these agents.

B. H. Vickery
B. Lunenfeld

VOLUME I

Basic Aspects

GNRH ANALOGUES
IN CANCER AND
HUMAN
REPRODUCTION

Edited by
B. H. Vickery and B. Lunenfeld



KLUWER ACADEMIC PUBLISHERS
DORDRECHT / BOSTON / LONDON

Distributors

for the United States and Canada: Kluwer Academic Publishers, PO Box 358,
Accord Station, Hingham, MA 02018-0358, USA

for all other countries: Kluwer Academic Publishers Group, Distribution Center,
PO Box 322, 3300 AH Dordrecht, The Netherlands

British Library Cataloguing in Publication Data

GnRH analogues in cancer and human reproduction.

Vol.1. Basic aspects

1. Women. Reproductive system. Cancer. Drug therapy

I. Vickery, B. H. (Brian H.), 1941- II. Lunenfeld, Bruno

616.99 465061

ISBN 0-7923-8927-1

ISBN 0-7923-8932-1 set

Copyright

© 1989 by Kluwer Academic Publishers

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior permission from the publishers, Kluwer Academic Publishers BV, PO Box 17, 3300 AA Dordrecht, The Netherlands.

Published in the United Kingdom by Kluwer Academic Publishers, PO Box 55,
Lancaster, UK.

Kluwer Academic Publishers BV incorporates the publishing programmes of D. Reidel,
Martinus Nijhoff, Dr W. Junk and MTP Press.

Printed in Great Britain by Butler and Tanner Ltd., Frome and London.

572
1
89
xy 1

CONTENTS

Contents of other volumes	vii
Preface	xiii
List of Contributors	xv
Introduction <i>B.H. Vickery and B. Lunenfeld</i>	1
1 Analogues of LHRH: the present and the future <i>A.V. Schally, S. Bajusz, T.W. Redding, A. Zalatnai and A.M. Comaru-Schally</i>	5
2 Chemistry of GnRH analogues <i>A.S. Dutta</i>	33
3 Active reduced size analogues <i>F. Haviv, C.A. Palabrica, E.N. Bush, G. Diaz, S.K. Burt, J. Greer and E.S. Johnson</i>	45
4 Intracellular aspects <i>L. Kiesel and B. Runnebaum</i>	53
5 Extrapituitary actions <i>J.G.M. Klijn and J.A. Foekens</i>	71
6 First generation GnRH antagonist: ovulation inhibition by weekly treatment <i>D. Kenigsberg and G.D. Hodgen</i>	85
7 Second generation GnRH antagonist: characterization of pituitary response in monkeys <i>C.F. Chillik, J. Itskovitz, D.W. Hahn, J.L. McGuire, D.R. Danforth and G.D. Hodgen</i>	95
8 The reproductive, endocrine and pharmacological evaluation of a new LHRH antagonist <i>D.W. Hahn, A. Phillips, J.L. McGuire, D.M. Ritchie, R.J. Capetola, C. Bowers and K. Folkers</i>	105

9	Pharmacokinetics of GnRH analogues	
	<i>J. Sandow, G. Jerabek-Sandow and B. Krauss</i>	113
10	Molecular mechanisms of GnRH action on mammary tumors and uterine leiomyomata	
	<i>J. Levy, T. Segal, A. Wiznitzer, V. Insler and Y. Sharoni</i>	127
11	Kinetics of sustained release delivery systems	
	<i>R. Deghengi</i>	137
12	Formulations of GnRH analogues for therapeutic use	
	<i>B.J.A. Furr and F.G. Hutchinson</i>	143
	Index	157

CONTENTS OF OTHER VOLUMES

Volume II. GnRH Analogues in Reproduction and Gynecology

Preface

- 1 Is there a rationale for GnRH analogue therapy in endometriosis?
I.A. Brosens, F. Cornillie and P. Puttemans
- 2 GnRH agonists in the management of endometriosis: the results of two randomized trials
W.P. Dmowski, E. Radwanska, Z. Binor, I. Tummon and P. Pepping
- 3 Comparison of buserelin to danazol therapy in endometriosis
A. Lemay
- 4 LHRH agonist treatment of endometriosis
A. Conti
- 5 A Phase III clinical trial of buserelin versus danazol in the treatment of endometriosis
W.H.M. Matta and R.W. Shaw
- 6 Management of pelvic endometriosis with a GnRH analogue
D.F.H. Li and P.C. Ho
- 7 The use of GnRH analogues to control menstrual bleeding
R.W. Shaw
- 8 Rationale, indications and management of GnRH analogues in ovulation induction protocols
B. Lunenfeld, V. Insler, E. Lunenfeld, G. Potashnik and L. Levy
- 9 Physiologic effects of a gonadotropin releasing hormone antagonist in normal women
J.E. Hall, T.D. Brodie, T.M. Badger, J. Rivier, W. Vale, P.M. Conn and W.F. Crowley, Jr.

- 10 Pharmacologic hypogonadotropism can be used to advantage for ovulation induction protocols
G. Bettendorf, W. Braendle, C.H. Lindner, V. Lichtenburg, M. Luckhardt and T. Schlotfeld
- 11 Induction of ovulation by combined GnRH-analogue/hMG/hCG treatment
M. Breckwoldt, F. Geithövel, J. Neulen and H. Schillinger
- 12 Combined buserelin and exogenous gonadotropins in ovulation induction in infertile women with normal menstrual rhythm
R. Fleming, M. Carter, M.P.R. Hamilton, M.E. Jamieson, M.J. Haxton, W.P. Black and J.R.T. Coutts
- 13 Combined buserelin and exogenous gonadotrophin therapy for the treatment of infertility in women with polycystic ovarian disease
J.R.T. Coutts, S. Finnie, C. Conaghan, M.J. Haxton, W.P. Black and R. Fleming
- 14 Comparison between short and long protocols using a combination of LHRH agonist and gonadotropins
L. Mettler
- 15 Induction of ovulation with the combination of LHRH analog in a short protocol and exogenous gonadotropins
P. Barriere, P. Lopes and B. Charbonnel
- 16 The use of GnRH analogues for synchronization of oocyte donation
P. Devroey, M. Camus, J. De Schacht, I. Kahn, J. Smits, C. Staesson, L. Van Waesberghe, A. Wisanto and A.C. Van Steirteghem
- 17 Programmed oocyte retrieval for IVF: clinical and biological effects of different protocols of pituitary suppression and follicular stimulation
S. Maschiach, Z. Ben-Rafael, A. Elenbogen, S. Lipitz, J. Blankenstein, D. Levran, A. Davidson, E. Rudak and J. Dor
- 18 Improved pregnancy rate in IVF/ET by combined long-acting GnRH analog and gonadotropins
R. Ron-El, E. Caspi, H. Nachum, A. Golan, A. Herman, Y. Soffer and Z. Weinraub

Index

Volume III. Benign and Malignant Tumors

Preface

- 1 The use of LHRH agonists in the treatment of uterine fibroids
D.L. Healey
- 2 Traitement des fibromes uterins par vaporization nasales de busereline
R. Erny, E. Milliet and L. Boubli
- 3 Advances in the treatment of leiomyomata uteri with leuprolide
A.J. Friedman
- 4 Relevance of an LHRH agonist to the treatment of uterine fibromyomas
J. Cohen and D. Elia
- 5 Comparison of treatment of uterine leiomyomata with 3 different GnRH agonist analogs
Z. Blumenfeld, M. Dirnfeld, D. Beck, H. Abramovici and J. M. Brandes
- 6 Sequential buserelin – medroxyprogesterone acetate treatment of uterine leiomyomata
G. Benagiano, A. Morini, A. Abbondante, V. Aleandri, F. Piccinno and D. Sala
- 7 Combined endoscopical and endocrinological treatment of uterine fibroids
J. Huber
- 8 Novel approaches to the treatment of advanced prostate cancer
A. Manni, R. Santen, A. Boucher, A. Lipton, H. Harvey, M. Simmonds, D. White-Hershey, M. Bartholemew, R. Caplan, R. Gordon, T. Rohner, J. Drago, J. Wettlaufer and L. Glode
- 9 Maximal androgen blockage in prostate cancer: today's therapy of choice and 5-year clinical experience
F. Labrie, A. Dupont, L. Cusan, G. Manhes, Y. Lacourciere, G. Monfette and J. Emond
- 10 Medical versus surgical orchiectomy in advanced prostatic cancer
H. Parmar, R. H. Phillips, L. Edwards and S. L. Lightman

- 11 Orchiectomy versus total androgen blockade. A Phase III EORTC 30853
L. Denis, F. Keuppens, D. Newling, P.H. Smith, F. Calais da Silva, R. Sylvester, M. De Pauw, P. Onegana and members of the EORTC GN Group, Belgium

- 12 Towards the best treatment for advanced prostatic cancer
J. Waxman, J. Sandow, P. Abel, N. Farah, J. Fleming, J. Cox, E.P.N. O'Donoghue, K. Sikora and G. Williams

- 13 Clinical studies of leuprolide depot formulation in metastatic prostatic cancer
L. Swanson, J. Seely, R. Browneller and D. Max

- 14 Treatment of advanced prostrate carcinomas by an LHRH analogue with or without an antiandrogen
I. Papadopoulos, F. Schaumkell, H. Bartermann and H. Wand

- 15 The use of GnRH analogues in ovarian cancer
G. Emons, G.S. Pahwa, R. Strum, R. Knuppen and F. Oberheuser

- 16 GnRH analogues in treatment of ovarian cancer
W. Jäger, W. Wildt and N. Lang

- 17 Partial tumor remission in recurrent low malignant stromal endometriosis following GnRH agonist treatment
Th. Bremen, S. Waibel, G. Bender, P. Heuschen and G. Leyendecker

- 18 Endocrine, pharmacokinetic and clinical effect of LHRH analogue treatment in patients with malignant and benign breast disease
J.G.M. Klijn, A.N. van Geel, F.H. de Jong and J. Sandow

- 19 LHRH analogs as adjuncts to somatostatin analogs in the treatment of pancreatic cancer
A.M. Comaru-Schally and A.V. Schally

- 20 An FSH producing tumor: response to GnRH analogue
M. Berezin, H. Halkin and B. Lunenfeld

Index

Volume IV. Precocious Puberty, Contraception, Safety Issues

Preface

- 1 Reversible inhibition of sexual maturation in male rats: a safety evaluation for the use of LHRH analogues in precocious puberty
J. Sandow, J. Engelbart, W. von Rechenberg, M. Hahn and S. Kille
- 2 Mechanism of action of GnRH analogues in sexual maturation and function
M.L. Aubert, R. Kreuter and P.C. Sizonenko
- 3 Should all girls with central precocious puberty be treated by LHRH analogue?
R. Brauner and R. Rappaport
- 4 Long-term experience with a superactive GnRH analogue in the treatment of precocious puberty
R. Kauli, A.V. Schally and Z. Laron
- 5 Effects of GnRH analogues on growth and bone maturation: a role for growth factors and adrenal androgens
S.L.S. Drop, W. Oostdijk, R.J.H. Odink, W.G. Sippell, C.J. Partsch and R. Hummelink
- 6 Growth patterns and skeletal maturation during sex steroid suppression and reactivation in central precocious puberty
P.A. Boepple, M.J. Mansfield, J.D. Crawford, J.F. Crigler, Jr., K. Link, R.M. Blizzard and W.F. Crowley, Jr.
- 7 Precocious puberty: pathophysiology and treatment
S.L. Kaplan, D. Stephure, J. Knezvic and M.M. Grumbach
- 8 Effects of GnRH agonists on pituitary gonadotropins: dissociated secretion of subunits in children with precocious puberty treated with [D-Trp⁶]LHRH microcapsules
M. Roger, N. Lahlou, J.L. Chaussain, C. Fouprie, R. Francois, J.E. Toublanc, R. Scholler and A.V. Schally
- 9 Testicular changes in patients treated with buserelin for precocious puberty
W.J. Tze, B. Gilks, H. Kitson, R.M. Couch, G.P. Taylor and P. Merat

- 10 Influence of buserelin on testicular tissue after long-term treatment for precocious puberty and prostatic cancer
F. Hadziselimovic
- 11 GnRH analogues in the development of a male contraceptive: importance of testosterone substitution and reversibility of pituitary and testicular suppression
G.F. Weinbauer and E. Nieschlag
- 12 Clinical pharmacology of LHRH antagonists
S.N. Pavlou, J. Rivier, W. Vale, T. Kamilaris
- 13 Clinical studies with the Na¹-Glu GnRH antagonist
Ph. Bouchard, S. Hajri, N. Lahlou, M. Roger, S. Brailly, S. Dubourdieu, R. Frydman, B. Charbonnel, S. Delivet, G. Schaison, Y.Y. Tsong and I.M. Spitz
- 14 The clinical relevance of bone mass measurements
I. Fogelmann and A. Rodin
- 15 Bone calcium dynamics in women with declining estrogen levels
D. Kenigsberg and M.E. Hull
- 16 Effects of GnRH agonist induced ovarian suppression on calcium homeostasis and bone metabolism in premenopausal women
S. Waibel, H.W. Minne, Th. Bremen, S.H. Sharla, R. Schiffl, C. Wüster, R. Zeigler and G. Leyendecker
- 17 GnRH analogs and safety
F. Comite
- 18 Management of post-marketing surveillance following short and long-term use of GnRH analogues
C.A. Schaffenburg

Index

PREFACE

These four volumes comprising "GnRH Analogues in Cancer and Human Reproduction" are a distillation of the presentations of the invited speakers at a landmark International Symposium bearing the same name, organized by one of us (B.L.) and held in Geneva, Switzerland in February 1988. The Symposium was truly interdisciplinary spanning gonadal hormone dependent disease including various forms of cancer and ranging to control of fertility, both pro- and conception. The international flavor can be caught from the 480 participants and 259 contributors drawn from 14 countries. The Symposium, and therefore this book, would not have been possible without the backing of The International Committee for Research in Reproduction and the sponsorship of the International Society of Gynecologic Endocrinology, The Swiss Society of Fertility and Sterility, The University of Geneva School of Medicine, The Swiss Society of Endocrinology and The US Foundation for Studies in Reproduction Inc., and help from the World Health Organization.

B. H. Vickery
B. Lunenfeld
June 1989

LIST OF CONTRIBUTORS

TO THE SERIES

A. Abbondante

First Institute of Obstetrics and
Gynecology
University "La Sapienza"
Rome, Italy

P. Abel

Department of Urology
Hammersmith Hospital
DuCane Road
London W12 0HS, UK

H. Abramovici

Departments of Obstetrics and
Gynecology
Rambam Medical Center and Carmel
Hospital, Technion
Israel Institute of Technology
Haifa 31096, Israel

V. Aleandri

First Institute of Obstetrics and
Gynecology
University "La Sapienza"
Rome, Italy

Michel L. Aubert

Department of Pediatrics and Genetics
Division of Biology of Growth and
Reproduction
University of Geneva Medical School
1211 Geneva 4, Switzerland

Tom M. Badger

Reproductive Endocrine Unit
Vincent Memorial Research Laboratories
Boston, MA 02114, USA

Sandor Bajusz

Veterans Administration Medical Center
1601 Perdido Street
New Orleans, LA 71046, USA

P. Barriere

IVF Department
CHU Nantes
44035-Nantes Cedex 01, France

H. Bartermann

Urologische Universitätsklinik Kiel
Arnold-Heller Strasse 7
D-2300 Kiel 1, FRG

M. Bartholomew

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

D. Beck

Departments of Obstetrics and
Gynecology
Ramban Medical Center and Carmel
Hospital, Technion
Israel Institute of Technology
Haifa 31096, Israel

G. Benagiano

First Institute of Obstetrics and
Gynecology
University "La Sapienza"
Rome, Italy

G. Bender

Department of Obstetrics and Gynecology
Städtische Kliniken
Grafenstrasse 9
D-6100 Darmstadt, FRG

Z. Ben-Rafael

Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

M. Berezin

Institute of Endocrinology
The Chaim Sheba Medical Center
Tel-Hashomer 52621, Israel

G. Bettendorf

UKE Frauenklinik
Division of Endocrinology
Martinistrasse 52
2000 Hamburg 20, FRG

Zvi Binor

Section of Reproductive
Endocrinology/Infertility
Department of Obstetrics and Gynecology
Rush Medical College
Chicago, IL 60612, USA

W.P. Black

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

J. Blankstein

Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

Robert M. Blizzard

Department of Pediatrics
University of Virginia Medical School
PO Box 386
Charlottesville, VA 22901, USA

Zeev Blumenfeld

Departments of Obstetrics and
Gynecology
Ramban Medical Center and Carmel
Hospital
Technion, Israel Institute of Technology
Haifa 31096, Israel

Paul A. Boepple

Reproductive Endocrine Unit
Massachusetts General Hospital
Rear Blossom Street
Boston, MA 02114, USA

L. Boubli

Gynecologie-Obstetrique
Hopital Michel Levy, Annexe Conception
84a Rue de Lodi
13281 Marseille Cedex 6, France

A. Boucher

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

Cyril Bowers

Endocrine Unit
Tulane University School of Medicine
New Orleans, LA 70112, USA

W. Braendel

UKE Frauenklinik
Division of Endocrinology
Martinistrasse 52
2000 Hamburg 20, FRG

J.M. Brandes

Departments of Obstetrics and
Gynecology
Ramban Medical Center and Carmel
Hospital, Technion
Israel Institute of Technology
Haifa 31096, Israel

R. Brauner

Unit of Pediatric Endocrinology and
Diabetes
Hospital des Enfants Malades
75015 Paris, France

M. Breckwoldt

Department of Obstetrics and Gynecology
Division of Clinical Endocrinology
University of Freiburg
D-7800 Freiburg-im-Breisgau, FRG

Th. Bremen

Department of Obstetrics and Gynecology
Staetische Kliniken
Grafenstrasse 9
D-6100 Darmstadt, FRG

Todd D. Brodie

Reproductive Endocrine Unit
Vincent Memorial Research Laboratories
Boston, MA 02114, USA

I.A. Brosens

Laboratory for Gynaecological
Physiopathology
UZ Gasthuisberg
KU Leuven, Belgium

Robert Browneller

Abbott Laboratories
Abbott Park, IL 60064, USA

S.K. Burt

Pharmaceutical Discovery Division
Abbott Laboratories
Abbott Park, IL 60064, USA

Eugene N. Bush

Pharmaceutical Discovery Division
Abbott Laboratories
Abbott Park, IL 60064, USA

M. Camus

Center for Reproductive Medicine
Medical Campus, Vrije University Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

R.J. Capetola

Ortho Pharmaceutical Corporation
Route 202
Raritan, NJ 08869, USA

R. Caplan

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvanian State University
PO Box 850, Hershey, PA 17033, USA

M. Carter

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

E. Caspi

Department of Obstetrics and Gynecology
Assaf Harofe Medical Center
Zerefin, Israel

B. Charbonnel

IVF Department
CHU Nantes
44035-Nantes Cedex 01, France

J.L. Chaussain

Foundation de Recherche en
Homonologie
PB110
94268 Fresnes Cedex, France

Claudia Chillik

Department of Obstetrics and Gynecology
Eastern Virginia Medical School
Norfolk, VA-23507, USA

Jean Cohen

Clinique Marignan
3 rue Marignan
75008 Paris, France

Ana Maria Comaru-Schally

Veterans Administration Medical Center
1601 Perdido Street
New Orleans, LA 70146, USA

Florence Comite

Department of Medicine and Gynecology
Yale University School of Medicine
New Haven, CT 06510-8063, USA

C. Conaghan

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

P. Michael Conn

Department of Pharmacology
University of Iowa College of Medicine
Iowa City, IO 52242, USA

Angelo Conti

Ch. de Mornex 6
1003 Lausanne, Switzerland

F. Cornillie

Laboratory for Gynaecological
Pathophysiology
UZ Gasthuisberg
KU Leuven, Belgium

R.M. Couch

Department of Pediatrics and Pathology
University of British Columbia
BC's Children's Hospital
Vancouver, BC V6H 3V4, Canada

J.R.T. Coutts

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

J. Cox

Department of Urology
Central Middlesex Hospital
London, UK

John D. Crawford
Children's Service
Massachusetts General Hospital
Fruit Street
Boston, MA 02114, USA

John F. Crigler, Jr.
Department of Medicine
Division of Endocrinology, Children's
Hospital
300 Longwood Avenue
Boston, MA 02115, USA

William F. Crowley Jr.
Departments of Medicine and Gynecology
Massachusetts General Hospital
Boston, MA 02114, USA

Lionel Cusan
Department of Molecular Endocrinology
Laval University Medical Center
Quebec G1V 4G2, Canada

Douglas R. Danforth
Department of Obstetrics and Gynecology
Eastern University Medical School
Norfolk, VA 23507, USA

F. Calais da Silva
A.Z. Middelheim
Antwerp 2020, Belgium

Adi Davidson
Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

R. Deghenghi
Debiopharm SA
1003 Lausanne, Switzerland

F.H. de Jong
Department of Medicine II and Clinical
Endocrinology
Erasmus University
Rotterdam, The Netherlands

L. Denis
A.Z. Middelheim
Antwerp 2020, Belgium

M. De Pauw
A.Z. Middelheim
Antwerp 2020, Belgium

J. De Schacht
Center for Reproductive Medicine
Medical Campus, Vrije Universiteit Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

P. Devroey
Center for Reproductive Medicine
Medical Campus, Vrije Universiteit Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

Gilbert Diaz
Pharmaceutical Discovery Division
Abbott Laboratories
Abbott Park, IL 60064, USA

M. Dirnfeld
Departments of Obstetrics and
Gynecology
Ramban Medical Center and Carmel
Hospital, Technion
Israel Institute of Technology
Haifa 31096, Israel

W. Paul Dmowski
Section of Reproductive
Endocrinology/Infertility
Department of Obstetrics and Gynecology
Rush Medical College
Chicago, IL 60612, USA

Joshua Dor
Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

J. Drago
Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

S.L.S. Drop
Sophia Children's Hospital
160 Gordelweg
3038 GE Rotterdam, The Netherlands

André Dupont

Department of Molecular Endocrinology
Laval University Medical Center
Quebec G1V 4G2, Canada

Anand S. Dutta

Pharmaceutical Division
Imperial Chemical Industries plc
Mereside, Alderley Park
Macclesfield, Cheshire SK10 4TG, UK

L. Edwards

Westminster Hospital
London SW1 2AP, UK

Adreian Elenbogen

Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

D. Elia

Clinique Marignan
3 rue Marignan
75008 Paris, France

Jean Emond

Laval University Medical Center
Quebec G1V 4G2, Canada

G. Emons

Klinik für Frauenheilkunde und
Geburtshilfe
Institute für Biochemische Endokrinologie
Medizinische Universität zu Lübeck
D-2400 Lübeck, FRG

K. Engelbart

Hoechst AG
D-6230 Frankfurt 80, FRG

R. Erny

Gynecologie-Obstetrique
Hospital Michael Levy, Annexe Conception
84a rue de Lodi
13281 Marseille Cedex 6, France

N. Farah

Department of Urology
Central Middlesex Hospital
London, UK

S. Finnie

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

J. Fleming

Department of Urology
Central Middlesex Hospital
London, UK

R. Fleming

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

J.A. Foekens

Division of Endocrine Oncology
Rotterdam Cancer Institute
The Dr. Daniel den Hoed Cancer Center,
Groene Hilledijk 301
3075 EA Rotterdam, The Netherlands

I. Fogelman

Department of Nuclear Medicine
Guy's Hospital
London, UK

Karl Folkers

Institute for Biomedical Research
The University of Texas at Austin
Austin, TX 78712, USA

C. Fouprie

Foundation de Recherche en
Hormonologie
PB110
94268 Fresnes Cedex, France

R. Francois

Foundation de Recherche en
Hormonologie
PB110
94268 Fresnes Cedex, France

Andrew J. Friedman

Brigham and Women's Hospital
Fertility and Endocrine Unit
75 Francis Street
Boston, MA 02115, USA

Barrington J.A. Furr

Bioscience I. Department
ICI Pharmaceuticals plc
Alderley Park, Macclesfield
Cheshire SK10 4TG, UK

F. Geisthövel

Department of Obstetrics and Gynecology
Division of Clinical Endocrinology
University of Freiburg
D-7800 Freiburg-im-Breisgau, FRG

B. Gilks

Department of Pediatrics and Pathology
University of British Columbia
BC's Children's Hospital
Vancouver BC V6H 3V4, Canada

L. Glode

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

A. Golan

Department of Obstetrics and Gynecology
Assaf Harofe Medical Center
Zerefin, Israel

Jessie C. Goodpasture

Institute of Biological Sciences
Syntex Research
3401 Hillview Avenue
Palo Alto, CA 94304, USA

R. Gordon

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

Jonathan Greer

Pharmaceutical Discovery Division
Abbott Laboratories
Abbott Park, IL 60064, USA

Melvin M. Grumbach

Department of Pediatrics
University of California San Francisco
San Francisco, CA 94943, USA

F. Hadziselimovic

Children's Hospital Basle
Römergasse 8
CH-4005 Basle, Switzerland

D.W. Hahn

Ortho Pharmaceutical Corporation
Route 202
Raritan, NJ 08869, USA

M. Hahn

Hoechst AG
D-6230 Frankfurt 80, FRG

H. Halkin

Institute of Endocrinology
The Chaim Sheba Medical Center
Tel-Hashomer 52621, Israel

Janet E. Hall

Reproductive Endocrine Unit
Vincent Memorial Research Laboratories
Boston, MA 02114, USA

M.P.R. Hamilton

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

H. Harvey

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

Fortuna Haviv

Pharmaceutical Discovery Division
Abbott Laboratories
Abbott Park, IL 60064, USA

M.J. Haxton

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

D.L. Healey

Medical Research Center
Monash Medical Center
Prince Henry's Hospital Campus
Melbourne, Australia

A. Herman

Department of Obstetrics and Gynecology
Assaf Harofe Medical Center
Zerefin, Israel

P. Heuschen

Department of Obstetrics and Gynecology
Staetdische Kliniken
Grafenstrasse 9
D-6100 Darmstadt, FRG

P.C. Ho

Department of Obstetrics and Gynecology
University of Hong Kong
Hong Kong

Gary D. Hodgen

Jones Institute for Reproductive Medicine
Eastern University Medical School
Norfolk, VA 23507, USA

J.C. Huber

1st Department of Gynecology and
Obstetrics
A-1090 Vienna, Austria

Magdalen E. Hull

Division of Reproductive Endocrinology
Department of Obstetrics and Gynecology
State University of New York at Stony
Brook School of Medicine
Stony Brook, NY 11794-8091, USA

R. Hummelink

Sophia Children's Hospital
160 Gordelweg
3038 GE Rotterdam, The Netherlands

F.G. Hutchinson

Pharmaceutical Department, ICI
Pharmaceutical plc
Alderley Park, Macclesfield
Cheshire SK10 4TG, UK

Vaclav Insler

Endocrinology Laboratory, Soroka
Medical Center and
Division of Obstetrics and Gynecology
Clinical Biochemistry Unit, Faculty of
Health Sciences
Ben-Gurion University of the Negev
Beer-Sheba 84101, Israel

Joseph Itskovitz

Department of Obstetrics and Gynecology
Eastern University Medical School
Norfolk, VA 23507, USA

M.E. Jamieson

University Department of Obstetrics and
Gynaecology
Glasgow Royal Infirmary
Glasgow G31 2ER, UK

G. Jerabek-Sandow

Hoechst AG
D-6230 Frankfurt 80, FRG

Edwin S. Johnson

Pharmaceutical Discovery Division
Abbott Laboratories
Abbott Park, IL 60064, USA

W. Jäger

Department of Gynecology and Obstetrics
University of Erlangen-Nürnberg
8520 Erlangen, FRG

Themis Kamilaris

Division of Endocrinology
Vanderbilt University School of Medicine
Nashville, TN 37232, USA

Selma L. Kaplan

Department of Pediatrics
University of California San Francisco
San Francisco, CA 94943, USA

R. Kauli

Institute of Pediatric and Adolescent
Endocrinology
Sackler Faculty of Medicine
Tel Aviv University
Tel Aviv, Israel

Daniel Kenigsberg

Division of Reproductive Endocrinology
Department of Obstetrics and Gynecology
State University of New York at Stony
Brook School of Medicine
Stony Brook, NY 11794-8091, USA

F. Keuppens

A.Z. Middelheim
Antwerp 2020, Belgium

I. Khan

Center for Reproductive Medicine
Medical Campus, Vrije Universiteit Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

Ludwig Kiesel

Division of Gynecological Endocrinology
Department of Obstetrics and Gynecology
University of Heidelberg
D-6900 Heidelberg, FRG

S. Kille

Hoechst AG
D-6230 Frankfurt 80, FRG

H. Kitson

Department of Pediatrics and Pathology
University of British Columbia
BC's Children's Hospital
Vancouver BC V6H 3V4, Canada

Jan G.M. Klijn

Division of Endocrine Oncology
Rotterdam Cancer Institute
The Dr. Daniel den Hoed Cancer Center,
Groene Hilledijk 301
3075 EA Rotterdam, The Netherlands

Jasna Knezvic

Department of Pediatrics
University of California San Francisco
San Francisco, CA 94943, USA

R. Knuppen

Klinik für Biochemische Endokrinologie
Medizinische Universität zu Lübeck
D-2400 Lübeck, FRG

B. Krauss

Hoechst AG
D-6230 Frankfurt 80, FRG

Renee Kreuter

Department of Pediatrics and Genetics
Division of Biology of Growth and
Reproduction
University of Geneva Medical School
1211 Geneva 4, Switzerland

Fernand Labrie

Department of Molecular Endocrinology
Laval University Medical Center
Quebec G1V 4G2, Canada

Yves Lacourciere

Laval University Medical Center
Quebec G1V 4G2, Canada

N. Lahlou

Foundation de Recherche en
Hormonologie
PB110
94268 Fresnes Cedex, France

N. Lang

Department of Gynecology and Obstetrics
University of Erlangen-Nürnberg
8520 Erlangen, FRG

Zvi Laron

Beilinson Medical Center
Petah Tikva 49100, Israel

André Lemay

Endocrinology of Reproduction
Hospital Saint-Francoise D'Asise
Quebec G1L 3L5, Canada

David Levran

Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

Joseph Levy

Endocrinology Laboratory, Soroka
Medical Center and
Division of Obstetrics and Gynecology
Clinical Biochemistry Unit, Faculty of
Health Sciences
Ben-Gurion University of the Negev
Beer-Sheba 84101, Israel

L. Levy

Tel Aviv University
Sackler School of Medicine
Tel Aviv, Israel

G. Leyendecker

Department of Obstetrics and Gynecology
Städtische Kliniken, Grafenstrasse 9
D-6100 Darmstadt, FRG

D.F.H. Li

Department of Obstetrics and Gynecology
University of Hong Kong
Hong Kong

V. Lichtenburg

UKE Frauenklinik
Division of Endocrinology
Martinistrasse 52
2000 Hamburg 20, FRG

S.L. Lightman

Charing Cross and Westminster Medical
School
London, UK

Ch. Lindner

UKE Frauenklinik
Division of Endocrinology
Martinistrasse 52
2000 Hamburg 20, FRG

Kathleen Link

Department of Medicine and Pediatrics
Harvard Medical School
Boston, MA 02115, USA

Schlomo Lipitz

Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

A. Lipton

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

P. Lopes

IVF Department
CHU Nantes
44035-Nantes Cedex 01, France

M. Luckhardt

UKE Frauenklinik
Division of Endocrinology
Martinistrasse 52
2000 Hamburg 20, FRG

Bruno Lunenfeld

Institute of Endocrinology
Sheba Medical Center and Bar Ilan
University
Remat Gan 52621, Israel

E. Lunenfeld

Division of Obstetrics and Gynecology
Soroka Medical Center and Ben-Gurion
University of the Negev
Beer-Sheba, Israel

J.L. McGuire

Ortho Pharmaceutical Corporation
Route 202
Raritan, NJ 08869, USA

Georgia I. McRae

Institute of Biological Sciences
Syntex Research
Palo Alto, CA 94304, USA

Gilles Manhes

Laval University Medical Center
Quebec G1V 4G2, Canada

Andrea Manni

Department of Medicine/Endocrinology
The Milton S. Hershey Medical Center
The Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

Joan Mansfield

Reproductive Endocrine Unit
Massachusetts General Hospital
Rear Blossom Street
Boston, MA 02114, USA

Shlomo Maschiach

Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

W.H.M. Matta

Academic Department of Obstetrics and
Gynaecology
Royal Free Hospital
London NW3 2QG, UK

Devorah T. Max
Abbott Laboratories
Abbott Park, IL 60064, USA

P. Merat
Pharmaceutical Division
Hoechst Canada Inc.
4045 Cote Vertu
Montreal, Quebec H4R 1R6, Canada

L. Mettler
Department of Obstetrics and Gynecology
University of Kiel
2300 Kiel, FRG

E. Milliet
Gynecologie-Obstetrique
Hopital Michael Levy, Annexe Conception
84a rue de Lodi
13281 Marseille Cedex 6, France

H.W. Minne
Department of Internal Medicine
University of Heidelberg
Heidelberg, FRG

Gerard Monfette
Laval University Medical Center
Quebec G1V 4G2, Canada

A. Morini
First Institute of Obstetrics and
Gynecology
University "La Sapienza"
Rome, Italy

H. Nachum
Department of Obstetrics and Gynecology
Assaf Harofe Medical Center
Zerefin, Israel

John J. Nestor Jr.
Institute of Bio-Organic Chemistry
Syntex Research
3401 Hillview Avenue
Palo Alto, CA 94304, USA

J. Neulen
Department of Obstetrics and Gynecology
Division of Clinical Endocrinology
University of Freiburg
D-7800 Freiburg-im-Breisgau, FRG

D. Newling
A.Z. Middelheim
Antwerp 2020, Belgium

Eberhard Nieschlag
Max Planck Clinical Research Unit for
Reproductive Medicine
Institute of Reproductive Medicine of the
University
D-4400 Münster, FRG

F. Oberheuser
Klinik für Frauenheilkunde und
Geburtshilfe
Institute für Biochemische Endokrinologie
Medizinische Universität zu Lübeck
D-2400 Lübeck, FRG

A.J.H. Odink
Sophia Children's Hospital
160 Gordelweg
3038 GE Rotterdam, The Netherlands

E.P.N. O'Donohue
Department of Urology
Central Middlesex Hospital
London, UK

P. Onegana
A.Z. Middelheim
Antwerp 2020, Belgium

W. Oostdijk
Sophia Children's Hospital
160 Gordelweg
3038 GE Rotterdam, The Netherlands

G.S. Pahwa
Klinik für Frauenheilkunde und
Geburtshilfe
Institute für Biochemische Endokrinologie
Medizinische Universität zu Lübeck
D-2400 Lübeck, FRG

Christopher A. Palabrica
Pharmaceutical Discovery Division
Abbott Laboratories
Abbott Park, IL 60064, USA

I. Papadopoulos
Urologische Universitätsklinik Kiel
Arnold-Heller Strasse 7
D-2300 Kiel 1, FRG

H. Parmar

Department of Oncology
Westminster Hospital
London SW1 2AP, UK

C.J. Partsch

Sophia Children's Hospital
160 Gordelweg
3038 GE Rotterdam, The Netherlands

Spyros N. Pavlou

Division of Endocrinology
Vanderbilt University School of Medicine
Nashville, TN 37232, USA

Peg Pepping

Section of Reproductive
Endocrinology/Infertility
Department of Obstetrics and Gynecology
Rush Medical Center
Chicago, IL 60612, USA

A. Phillips

Ortho Pharmaceutical Corporation
Route 202
Raritan, NJ 08869, USA

R.H. Phillips

Westminster Hospital
London SW1 2AP, UK

F. Piccinno

First Institute of Obstetrics and
Gynecology
University "La Sapienza"
Rome, Italy

G. Potashnik

Division of Obstetrics and Gynecology
Soroka Medical Center and Ben Gurion
University of the Negev
Beer Sheva, Israel

P. Puttemans

Laboratory for Gynaecological
Pathophysiology
Gasthuisberg
KU Leuven, Belgium

Ewa Radwanska

Section of Reproductive
Endocrinology/Infertility
Department of Obstetrics and Gynecology
Rush Medical College
Chicago, IL 60612, USA

R. Rappaport

Unit of Pediatric Endocrinology and
Diabetes
Hopital des Enfants Malades
75015 Paris, France

Tommie W. Redding

Veterans Administration Medical Center
1601 Perdido Street
New Orleans, LA 71046, USA

D.M. Ritchie

Ortho Pharmaceutical Corporation
Route 202
Raritan, NJ 08869, USA

Jean Rivier

Salk Institute for Biological Studies
La Jolla, CA 92037, USA

A. Rodin

Department of Gynaecology
Guy's Hospital
London, UK

M. Roger

Foundation de Recherche en
Hormonologie
PB110
94268 Fresnes Cedex, France

T. Rohner

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

R. Ron-El

Department of Obstetrics and Gynecology
Assaf Harofe Medical Center and
Zerefin, Israel

Edwina Rudak

Interdepartmental Unit of Human
Reproduction
Department of Obstetrics and Gynecology
The Chaim Sheba Medical Center and
Sackler School of Medicine
Tel-Hashomer 52621, Israel

Benno Runnebaum

Division of Gynecological Endocrinology
Department of Obstetrics and Gynecology
University of Heidelberg
D-6900 Heidelberg, FRG

D. Sala

First Institute of Obstetrics and
Gynecology
Rome, Italy

Lynda M. Sanders

Institute of Pharmaceutical Sciences
Syntex Research
Palo Alto, CA 94304, USA

Jurgen Sandow

Department of Pharmacology
Hoechst AG
D-6230 Frankfurt 80, FRG

R. Santen

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

Carlos A. Schaffenburg

5480 Wisconsin Avenue 1014
Chevy Chase, MD 20815, USA

Andrew V. Schally

Endocrine Polypeptide and Cancer
Institute
Veterans Administration Medical Center
and Section of Experimental Medicine
Department of Medicine, Tulane University
School of Medicine
New Orleans, LA 70146, USA

S.H. Scharla

Department of Internal Medicine
University of Heidelberg
Heidelberg, FRG

F. Schaumkell

Urologische Universitätsklinik Kiel
Arnold-Heller Strasse 7
D-2300 Kiel 1, FRG

H. Schillinger

Department of Obstetrics and Gynecology
Division of Clinical Endocrinology
University of Freiburg
D-7800 Freiburg-im-Breisgau, FRG

T. Schlotfeld

UKE Frauenklinik
Division of Endocrinology
Martinistrasse 52
2000 Hamburg 20, FRG

R. Scholler

Foundation de Recherche en
Hormonologie
PB110
94268 Fresnes Cedex, France

John H. Seely

Abbott Laboratories
Abbott Park, IL 60064, USA

Tzuria Segal

Endocrinology Laboratory, Soroka
Medical Center and
Division of Obstetrics and Gynecology
Clinical Biochemistry Unit, Faculty of
Health Sciences
Ben Gurion University of the Negev
Beer-Sheba 84101, Israel

Yoav Sharoni

Endocrinology Laboratory, Soroka
Medical Center and
Division of Obstetrics and Gynecology
Clinical Biochemistry Unit, Faculty of
Health Sciences
Ben Gurion University of the Negev
Beer-Sheba 84101, Israel

R.W. Shaw

Academic Department of Obstetrics and
Gynecology
Royal Free Hospital School of Medicine
London NW3, UK

R. Shiffl

Department of Internal Medicine
University of Heidelberg
Heidelberg, FRG

Karol Sikora

Department of Clinical Oncology
Hammersmith Hospital
DuCane Road
London W12 0HS, UK

M. Simmonds

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

W.G. Sippell

Sophia Children's Hospital
160 Goodelweg
3038 GE Rotterdam, The Netherlands

Pierre C. Sizonenko

Department of Pediatrics and Genetics
Division of Biology of Growth and
Reproduction
University of Geneva Medical School
1211 Geneva 4, Switzerland

P.H. Smith

A.Z. Middelheim
Antwerp 2020, Belgium

J. Smitz

Center for Reproductive Medicine
Medical Campus, Vrije Universiteit Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

Y. Soffer

Department of Obstetrics and Gynecology
Assaf Harofe Medical Center
Zerefin, Israel

C. Staesson

Center for Reproductive Medicine
Medical Campus, Vrije Universiteit Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

David Stephure

Department of Pediatrics
University of California San Francisco
San Francisco, CA 94943, USA

R. Strum

Klinik für Frauenheilkunde und
Geburtshilfe
Institut für Biochemische Endokrinologie
Medizinische Universität zu Lübeck
D-2400 Lübeck, FRG

Linda J. Swanson

Abbott Laboratories
Abbott Park, IL 60064, USA

R. Sylvester

A.Z. Middelheim
Antwerp 2020, Belgium

G.P. Taylor

Departments of Pediatrics and Pathology
University of British Columbia
BC's Children's Hospital
Vancouver, BC V6H 3V4, Canada

J.E. Toublane

Foundation de Recherche en
Hormonologie
PB110
94268 Fresnes Cedex, France

Ian Tummon

Section of Reproductive
Endocrinology/Infertility
Department of Obstetrics and Gynecology
Rush Medical College
Chicago, IL 60612, USA

W.J. Tze

Department of Pediatrics and Pathology
University of British Columbia
BC's Children's Hospital
Vancouver, BC V6H 3V4, Canada

Wylie Vale

Salk Institute for Biological Studies
La Jolla, CA 92037, USA

A.N. van Geel

Department of Surgery
The Rotterdam Cancer Institute
Rotterdam, The Netherlands

A.C. Van Steirteghen

Center for Reproductive Medicine
Medical Campus, Vrije Universiteit Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

L. Van Waesberghe

Center for Reproductive Medicine
Medical Campus, Vrije Universiteit Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

Brian H. Vickery

Institute of Biological Sciences
Syntex Research
Palo Alto, CA 94304, USA

W. von Rechenberg

Hoechst AG
D-6230 Frankfurt 80, FRG

S. Waibel

Department of Obstetrics and Gynecology
Städtische Kliniken
Grafenstrasse 9
D-6100 Darmstadt, FRG

H. Wand

Urologische Universitätsklinik Kiel
Arnold-Heller Strasse 7
D-2300 Kiel 1, FRG

Jonathan Waxman

Department of Clinical Oncology
Hammersmith Hospital
DuCane Road
London W12 OHS, UK

Gerhard F. Weinbauer

Max Planck Clinical Research Unit for
Reproductive Medicine
Institute of Reproductive Medicine of the
University
D-4400 Münster, FRG

Z. Weinraub

Department of Obstetrics and Gynecology
Assaf Hasofe Medical Center
Zerefin, Israel

J. Wettlaufer

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

D. White-Hershey

Department of Medicine/Endocrinology
Milton S. Hershey Medical Center
Pennsylvania State University
PO Box 850, Hershey, PA 17033, USA

L. Wildt

Department of Gynecology and Obstetrics
University of Erlangen-Nürnberg
8520 Erlangen, FRG

G. Williams

Department of Urology
Hammersmith Hospital
DuCane Road
London W12 OHS, UK

A. Wisanto

Center for Reproductive Medicine
Medical Campus, Vrije Universiteit Brussel
Laarbeeklaan 101
1090 Brussels, Belgium

Arnon Wiznitzer

Endocrinology Laboratory, Soroka
Medical Center and
Division of Obstetrics and Gynecology
Clinical Biochemistry Unit, Faculty of
Health Sciences
Ben-Gurion University of the Negev
Beer-Sheba 84101, Israel

C. Wüster

Department of Internal Medicine
University of Heidelberg
Heidelberg, FRG

Attila Zalatnai

Veterans Administration Medical Center
1601 Perdido Street
New Orleans, LA 71046, USA

R. Ziegler

Department of Internal Medicine
University of Heidelberg
Heidelberg, FRG

INTRODUCTION

Brian H. VICKERY and Bruno LUNENFELD

Institute of Biological Sciences, Syntex Research, Palo Alto,
CA 94304, USA and Institute of Endocrinology, Sheba Medical Center and
Bar Ilan University, Ramat Gan 52621, Israel

Since 1971 and the isolation, identification and synthesis of mammalian GnRH by Drs. Schally and Guillemin the pace of research in this subject area has been explosive and involved many disciplines. Synthetic efforts have produced several thousand analogues; seven of the most potent agonists are under commercial sponsorship and are either marketed or in late stages of clinical trial around the world. Prostatic cancer, precocious puberty and endometriosis patients now have a valuable new therapy to alleviate their condition and progress. A range of other applications is summarized in Chapter 1 and Volumes 2-4.

Basic Aspects

The chemistry of the nona- and decapeptide agonist analogues (Chapter 2) appears to have matured and no advances in potency have been reported since the early 1980s. An intriguing new direction for chemistry may follow the reports (Chapter 3) on new reduced size analogues which may be tailored to agonism or antagonism, at least in vitro. If high potency could be achieved in vivo these hexapeptides could potentially lead to the first orally administered agents, assuming stability to gastro-intestinal degradation. We are finally beginning to understand the molecular biology of GnRH (Chapter 4) and this in turn may suggest new chemical directions. The controversy over extrapituitary receptors and sites of action of GnRH in animals and man continues (Chapters 5 & 10); early reports of direct inhibitory effects of GnRH agonists on breast cancer cells seem to have given way to evaluation of the antagonists for this purpose. We have progressed through several generations of GnRH competitive antagonist analogues (Chapters 6-8) and in this iterative fashion may finally be approaching molecules which retain sufficient potency, while minimizing adverse effects, to be clinically evaluable. As noted, presently available analogues of GnRH are not deliverable orally but great progress has been made with alternatives to daily injection, first by intranasal insufflation

and then the potentially more compliance-compatible 1-3 month depot injectable formulations (Chapters 9, 11-12).

Reproduction and Gynecology

The utility of GnRH agonists in the management of endometriosis and their superior acceptability to existing approved therapy appears amply demonstrated, at least in so far as pain and size/extent of lesions are concerned (Chapters 13-18). Whether this, or indeed any other form of management, increases fertility outcome is still open to debate. To the extent that the GnRH agonists shut down ovarian function they are useful in symptomatic relief of dysfunctional bleeding and polycystic ovarian disease (Chapters 19 and 25). Correction of these conditions beyond the period of treatment however is not likely. A triphasic regimen, consisting of a GnRH agonist prior to and/or in conjunction with exogenous gonadotropins is being used in primary, idiopathic infertility, particularly in clomiphene failure and in secondary infertility such as polycystic ovarian disease, oligomenorrhea and premature menopause induced by chemotherapy. In these cases the suppression of endogenous gonadotropins with the GnRH analogue makes it easier to control follicular development and also eliminates the problems of premature and asynchronous luteinization (Chapters 22-24). This regimen also is being evaluated for oocyte harvest in IVF/ET programs (Chapters 28, 29).

Management of Tumors of the Reproductive System

A wealth of evidence now indicates the GnRH agonists to be equivalent to orchiectomy in palliation of metastatic prostatic cancer (Chapters 40, 43). Problems associated with the flare or initial stimulatory effect of the analogues and possibly to continued adrenal androgen secretion, a concept initially championed by Labrie, are receiving wide geographical attention (Chapters 39-42, 44). It seems that "total androgen blockade" by combination of a GnRH agonist and an antiandrogen may slightly delay time to progression of disease but do little for survival figures, with the notable exception of the results from one group. Breast cancer, ovarian cancer, uterine leiomyomata and BPH are all receiving attention although the numbers of cases so far reported are few (Chapters 31-37, 45-48). Interestingly, on the basis of animal studies, some tumor types such as pancreatic acinar and osteosarcomas may need to be redefined as to their hormonal dependency. Animal studies also suggest that coadministration of GnRH analogues with representatives of another new peptide series, the somatostatin analogues, may be beneficial for treatment of certain tumors (Chapter 49).

Precocious Puberty

The relatively rare but relatively devastating syndrome of central or true isosexual precocious puberty may now be treatable on more than a symptomatic level. Several studies with the GnRH agonists are now of sufficient duration to suggest that a major, positive influence on final adult height may result from long term treatment (Chapters 53-57).

Safety

As always, with any new class of therapeutic agents, safety considerations must be paramount. Potent endocrine drugs such as the GnRH analogues should not be expected to be without "side effects". These unwanted effects will include sequelae of the mechanism of action and reflect the induction of a hypogonadotropic hypogonadism, such as vasomotor symptoms, in both sexes and osteoporotic changes in women, together with changes in gonadotropin subunit secretion (Chapters 51, 58). Clearly in this situation, benefit - to - risk considerations, as far as they can be assessed, will play a role (Chapters 67, 68). In so far as established therapy already includes extirpation of the gonads, no additional risk would appear to ensue from the use of GnRH agonists. The situation with the antagonistic analogues of GnRH thus far is more complex, involving the additional factor of an anaphylactoid response. Additional chemical efforts are required before these agents are to be considered for widespread trial, let alone therapeutic candidates.

The geographic and linguistic diversity of the contributors to this book together with constraints of early publication resulted in a somewhat autocratic editorial policy. The contributors are to be congratulated on their hard work; any inadvertent omissions or changed emphasis or context are the responsibility of the editors.

ANALOGS OF LHRH: THE PRESENT AND THE FUTURE

**Andrew V. SCHALLY, Sandor BAJUSZ, Tommie W. REDDING,
Attila ZALATNAI and Ana Maria COMARU-SCHALLY**

Veterans Administration Medical Center, 1601 Perdido Street,
New Orleans, LA 71046, USA

INTRODUCTION

Seventeen years have passed since our laboratory accomplished the isolation, determination of structure and synthesis of the hypothalamic hormone controlling the secretion of both LH and FSH from the anterior pituitary gland [1-6]. This hypothalamic hormone is known as the LH and FSH releasing hormone, (LHRH/FSHRH) or gonadotropin-releasing hormone (GnRH) [6, 7]. While LHRH is also established as the main FSH releasing hormone, for reasons of convenience and historical continuity, the abbreviation LHRH is generally used for naming its analogs.

Many studies demonstrated that LHRH is the main link between the brain and the pituitary gland insofar as reproductive function is concerned [5-7]. Various clinicians, particularly our collaborators, established clinical uses of LHRH [8-13]. In 1971, we also postulated that LHRH or its analogs might be useful for the control of fertility by disrupting the menstrual cycle and preventing the ovulatory LH surge [14]. We suggested that replacement of one or more amino acids in LHRH might result in analogs possessing features required for binding to receptors, but devoid of intrinsic activity. Since 1972, systematic work has been proceeding to synthesize agonistic and antagonistic analogs of LHRH. A strong interest in medical applications of LHRH derivatives stimulated this undertaking. In the past 16 years, more than 2000 analogs of LHRH have been synthesized. Agonistic analogs more potent than the parent hormone have been made. Many of these analogs are being used clinically (Table 1) and the list of their applications is steadily expanding.

Various investigators have determined that large doses of superactive stimulatory analogs of LHRH cause paradoxical antifertility effects in animals and human being. Extensive work is in progress to apply these effects to the development of new contraceptive methods. The phenomena of pituitary desensitization and inhibition of sex steroid levels by LHRH agonists are being used for treatment of true idiopathic precocious puberty, endometriosis, leiomyomas and benign prostatic hyperplasia (BPH). Chronic administration of LHRH agonists is being utilized to

induce the regression of endocrine-dependent malignant neoplasms, especially prostate and breast cancer and more recently, ovarian carcinoma. LHRH agonists are also being tried in IVF-ET, GIFT and polycystic ovarian disease.

TABLE I
LH-RH AGONISTS IN CLINICAL USE

	STRUCTURE										RELATIVE POTENCY	ROUTE OF ADMINISTRATION
	1	2	3	4	5	6	7	8	9	10		
LH-RH	p- Glu - His - Trp - Ser - Tyr - Gly - Leu - Arg - Pro - Gly-NH ₂											
Buserelin (Hoechst)	D-Ser(TBU)					Ethylamide					100	SC, IN, DEPOT
Nafarelin (Syntex)	D-(2-Nal)										200	SC, IN, DEPOT
Leuprolide (Abbott-Takeda)	D-Leu					Ethylamide					50	SC, IN
Lutrelin (Wyeth)	D-Trp-7-N-Me-Leu					Ethylamide					100	
Goserelin (ICI)	D-Ser(TBU)					Az-Gly-NH ₂					50	SC, DEPOT
Histrelin (Ortho)	D-His(Bzl)					Ethylamide					100(?)	
Decapeptyl (Debiopharm)	D-Trp										100	SC OR DEPOT

SC, subcutaneous; IN, intranasal; DEPOT-IM, intramuscular for once a month microcapsules or for implants

Potent inhibitory analogs of LHRH, which block ovulation in laboratory animals, have been synthesized. Several antagonists of LHRH have been tested in men and women and shown to be active enough for practical use. The approach based on inhibitory analogs of LHRH has clearly been proved feasible for the development of new methods of birth control, although the exact clinical regimens are still lacking. In addition, it is possible that the LHRH antagonists could be useful for treatment of precocious puberty, endometriosis, and also for management of breast and prostate carcinoma. Thus, analogs of LHRH have various practical applications including the treatment of hormone sensitive tumors. We will review now some selected experimental and clinical findings on the agonistic and antagonistic analogs of LHRH, principally those obtained by us and our collaborators.

LHRH AGONISTS

Initially it was thought that LH releasing activity and ovulation-

inducing effects of LHRH observed in laboratory animals could be applied to the treatment of male and female infertility [15, 16]. However, the half-life of LHRH is very short [17, 18] and more potent and longer-acting analogs were considered to be necessary for clinical applications. A greater binding affinity and enzymatic resistance to proteolysis [19] became the goal for the rational design of all the LHRH analogs. Potent agonists were synthesized, but it was discovered that following chronic administration, they exert anti-fertility effects and clinical results aimed at fertility stimulation were disappointing.

Several LHRH analogues substituted in positions 6, 10 or both are much more active than LHRH (Table 1) and also possess prolonged activity [7, 20-27].

Paradoxical Antifertility Effects

Although an acute injection of superactive agonists of LHRH induces a marked and prolonged release of LH and FSH, paradoxically, chronic administration produces dramatic inhibitory effects through a process of "down regulation" of pituitary receptors for LHRH, desensitization of the pituitary gonadotropes and reduction in gonadal receptors for LH and FSH [7, 25-30]. In male animals this is manifested by a fall in the weights of testes, seminal vesicles, and prostate, a decrease in plasma testosterone levels, and a reduction in testicular LH/hCG receptors [7, 26, 30-32]. In men, a persistent suppression of Leydig cell function, manifested by a fall in serum testosterone and dihydrotestosterone levels, has been observed after chronic administration of LHRH agonists [33-35].

Chronic administration of agonistic analogs of LHRH induces paradoxical inhibitory effects on pituitary gonadotropes and gonads in female animals analogous to those reported in males. Among the effects reported in female rats are: delay in vaginal opening; reduction in ovarian and uterine weight; interference with mating and implantation; termination of gestation; and a fall in ovarian receptors for LH/hCG [7, 15, 28, 36, 37]. Antifertility effects, after administration of LHRH agonists, have also been documented in females of other species, including primate and human [15, 37, 38]. Mice are more resistant to the inhibitory effects of LH-RH agonists [39], but our recent results indicate that gonadal suppression is eventually induced [40].

Attempts are being made to utilize antifertility effects of LHRH agonists in men and women for the development of contraceptive methods [41, 42]. Continuous administration of high doses of LHRH agonists causes a suppression of circulating levels of LH (bioactive but not always radioimmunoactive) [43] and of ovarian estrogen levels [44]. The inhibition of ovarian steroidogenesis and LH forms the basis for the therapeutic use of LHRH agonists in diseases and conditions which result from inappropriate hormone levels or which can be treated by suppression of estrogens. These applications include endometriosis, polycystic ovarian disease, uterine leiomyomas, dysfunctional uterine bleeding, and hirsutism [44-47].

Therapeutic Uses

Because of potential risk of down-regulation phenomena, synthetic LHRH itself is being used for induction of ovulation, treatment of cryptorchidism and hypogonadism. Within the past three years, using automatic portable pumps designed to deliver pulses of synthetic LHRH, several investigators have demonstrated the effectiveness and safety of this mode of ovulation induction [48-50]. Treatment of cryptorchidism in prepubertal boys by intranasal administration of synthetic LHRH has also been reported [51, 52].

Female Contraception

Since the inhibition of pituitary-gonadal functions induced by chronic administration of LHRH agonists is reversible, attempts are being made to develop contraceptive methods based on the use of LHRH agonists [16, 38, 42, 53, 54]. Contraceptive action of LHRH agonists was revealed in studies on inhibition of ovulation in women when repeated administration was used. Nillius and collaborators reported that chronic subcutaneous or intranasal administration of [D-Ser(Bu^t)⁶,Pro⁹-NHet]LHRH (HOE766) to regularly menstruating women could inhibit ovulation [38, 41]. More recently, Gudmundsson reported a study in which forty seven women used intranasal nafarelin for contraception over a period of six months. Ovulation was inhibited during 261 of 262 treatment months and no pregnancy occurred [55]. However, some problems may exist with this approach. Spotting, amenorrhea, unpredictable follicular maturation, effects of unopposed estradiol secretion on the endometrium as well as hypoestrogenism leading to osteoporosis [42, 54-58] must be more carefully investigated before this method could be accepted for long term contraception even with progestins as supplements.

Other approaches to female contraception based on LHRH agonists may also be linked with major problems. Although LHRH and its superactive agonists induce luteolysis in women and in monkeys if administered in mid luteal phase [59], the practical application of this finding for prevention or interception of implantation is highly limited because of the careful timing required for administration of LHRH agonists [60]. It is improbable that LHRH agonists can be used as abortifacients or interceptives in women [61, 62].

Male Contraception

It is also unlikely that LHRH agonists will be suitable as male contraceptives. Linde et al, [63] reported a fall in testosterone levels and inhibition of spermatogenesis in 8 normal men after subcutaneous treatment with [D-Trp⁶,Pro⁹-NHEE]LHRH for 6-10 weeks. However, 5 men became impotent and had to discontinue the treatment [63]. Replacement therapy with long acting testosterone preparations would prevent the decline in libido and forestall the hot flashes seen after LHRH agonists [64]. Bhasin et al., [65]

and Doelle and colleagues [66] noted only a partial suppression of spermatogenesis using a combined regimen of LHRH agonists with testosterone. Moreover, long-term administration of exogenous testosterone in methods intended for male contraception might lead to an increased incidence of prostate cancer and/or BPH in men [67]. The suitability of this method for male contraception is questionable.

Endometriosis

That LHRH agonists may be useful in the management of endometriosis was first shown by Meldrum et al., [68] and Lemay and Quesnel [69]. Meldrum et al., [68] treated 5 women with proven pelvic endometriosis with agonist [D-Trp⁶,Pro⁹-NHet] LHRH for four weeks and achieved a "medical oophorectomy".

In our study [70], 8 menstrually cycling women with endometriosis were treated with a delayed-release preparation of [D-Trp⁶]LHRH in biodegradable microcapsules for a period of 3 to 5 months. A sustained hypogonadal state, characterized by low estradiol levels was induced in all patients in 2 weeks. In all patients, estradiol levels fell to zero after the second or third injection but gonadotropins remained in the normal range. Hot flashes and severe dyspareunia were the main side effects. Resumption of normal pituitary-ovarian activity was documented in all patients 40-60 days after the last injection of microcapsules. All patients showed a clinical improvement. Other studies resulted in similar findings [71].

Polycystic Ovarian Disease

Chang et al. [44] investigated the effects of chronic administration of LHRH agonists in polycystic ovary syndrome. The concentration of estradiol declined throughout treatment. Complete suppression of ovarian androgen production may be due to the marked reduction of bioactive LH levels [43, 44].

Ayalon et al. [72] carried out a study in which 17 patients with polycystic ovarian disease (PCOD) and severe ovarian hyperstimulation syndrome (OHSS) during therapy with HMG/HCG were treated with [D-Trp⁶]LHRH daily SC or in microcapsules until medical gonadectomy was attained. Under the suppressive therapy with the LHRH agonist, ovulation was induced with hFSH/HCG. In 15 of 17 patients, ovulatory cycles were obtained and 7 patients conceived. Most patients demonstrated symptoms of mild OHSS and only one patient developed severe OHSS. LHRH agonist administration for several months, to reduce ovarian androgen production and ameliorate the ovarian cystic disease process, followed by exogenous gonadotropin treatment may allow successful ovulation induction [72, 73].

In Vitro Fertilization and Embryo Transfer (IVF-ET) and Gamete Intra Fallopian Transfer (GIFT)

In IVF-ET and GIFT programs, LHRH analogs can be used to suppress

ovarian function early in the follicular phase and then an hMG followed by hCG regimen is employed to induce ovulation [74-76]. The ova may be harvested for in vitro manipulation and transfer or GIFT. The elimination of interference from endogenous gonadotropins appears to reduce the risk of ovarian hyperstimulation [74-76].

Leiomyomas

Filicori et al. [77] and Healy et al. [78] reported that the size of uterine leiomyomata could be reduced in patients by long-term treatment with LHRH agonists. Maheux and Lemay [47] employed buserelin for therapy of patients with uterine tumors. They found that this treatment resulted in tumor shrinkage, presumably because of the induction of a state of hypoestrogenism.

Coddington et al. [79] evaluated the rate at which uterine leiomyomas shrink in response to the induction of a state of hypoestrogenism by the administration of an LHRH analog. Even in patients with relatively small uterine leiomyomata, they observed at least a 50% reduction in uterine volume. Similar results were obtained by Van Leusden [80].

Perl et al. [81] used chronic administration of [D-Trp⁶] LHRH to suppress the secretion of ovarian steroids for treatment of leiomyomata uteri. Ten menstruating women, presenting with a total of 20 uterine leiomyomas, were treated for three months by daily subcutaneous injections. The volume of seven uterine leiomyomas diminished by more than 80%, while 8 leiomyomas decreased partially. After an initial reduction, 5 leiomyomas re-enlarged during the last month of treatment. The agonist was well tolerated and few side effects were observed. It was concluded that therapy with LHRH agonists offers a good alternative in the management of some uterine leiomyomas [81]. Similar findings were obtained by George et al. [82]. These results suggest that LHRH agonist-induced hypoestrogenism may be a useful method for management of uterine leiomyomata, as a primary therapy or as an adjunct to leiomyomectomy.

Ovarian Cancer

Gonadotropins might increase the incidence of cancers of the ovary in animals and humans [31]. The effects of chronic administration of agonists of LHRH are being investigated in murine models and in nude mice bearing transplanted human ovarian epithelial tumors. Suppression of the secretion of gonadotropins produced by LHRH agonists appears to inhibit the growth of ovarian tumors [31, 42]. Parmar et al. [83] reported that chronic treatment with [D-Trp⁶]LHRH microcapsules induced the regression of an inoperable, bilateral, serous cystadenocarcinoma of the ovary in a 78 year old woman. More recently, these findings were confirmed and extended in a trial of 39 women with epithelial ovarian cancer [84]. Eleven patients showed remission or stabilization of disease.

Prostate Cancer

A marked inhibition of pituitary and gonadal function that occurs after chronic administration of agonists of LHRH, made possible a new approach for the treatment of sex hormone-dependent tumors [30, 31, 42]. In our initial studies, we have shown that chronic administration of [D-Trp⁶]LHRH suppressed tumor growth in rats with Dunning R3327H prostate cancers and reduced serum levels of testosterone [85]. Subsequent clinical trials documented marked improvement in patients with stage C or D prostate carcinoma after treatment with [D-Trp⁶]LHRH, buserelin, leuprolide or zoladex [86-95]. Initially, superagonists of LHRH were given daily by the subcutaneous or intranasal route [86-95]. Subsequently, we developed a long-acting delivery system for [D-Trp⁶]LHRH in microcapsules of poly (DL-lactide-co-glycolide) designed to release a controlled dose of the peptide over a 30-day period [96, 97].

Microcapsules of [D-Trp⁶]LHRH that can be injected once a month make the treatment of patients with prostate carcinoma and other neoplasms more convenient. The efficacy of the slow-release formulation of [D-Trp⁶]LHRH microcapsules in the treatment of advanced prostatic carcinoma was demonstrated in animal experiments and in clinical trials [96, 98, 99]. In a trial with microcapsules in men with prostate cancer carried out with Parmar, Lightman and Phillips, we obtained 87% objective response in the microcapsule treated group vs. 81% for total orchiectomy [99].

The existing results warrant continuation of clinical trials with LHRH agonists for the treatment of advanced prostate carcinoma. However, it is possible that the therapeutic response could be improved by combining LHRH agonists with other compounds including peptides such as somatostatin analogs and various chemotherapeutic agents [100-105].

The duration of remission in patients with prostate cancer may be limited as hormonal manipulations do not prevent the ultimate growth of hormone-independent cells [106, 107]. Combination of hormonal therapy with chemotherapy could forestall this phenomenon and prolong survival [107]. The effect of combining hormonal treatment consisting of microcapsules of [D-Trp⁶]LHRH with various chemotherapeutic agents has been investigated in the Dunning prostate cancer model. The combination of cyclophosphamide (Cytoxan) with the microcapsules was much more effective than the single agents in reducing tumor growth [103]. Novantrone (Mitoxantrone), an anthracenedione similar to adriamycin, but less toxic, was also used in combination with microcapsules of [D-Trp⁶]LHRH [104]. Again, the combination with Novantrone led to a better inhibition of prostate cancer than [D-Trp⁶]LHRH microcapsules alone and in fact, arrested tumor growth [104]. Pathological examination of the tumors showed that the combination therapy inhibited the growth of tumor cells and that these cells were replaced by connective tissue [108].

Prolactin could be involved in prostate cancer as a cofactor [101, 102, 109]. Somatostatin analogs inhibit prolactin and

growth hormone secretion and have direct antiproliferative effects on cells [101, 102, 105]. The reduction in prolactin levels produced by the administration of somatostatin analogs, combined with the decrease in serum testosterone resulting from chronic treatment with LHRH agonists, may inhibit growth of prostate tumors better than LHRH agonists alone [101, 102, 105].

The fall in GH levels induced by somatostatin analogs could, through mechanisms involving endogenous growth factors, be even more important for the inhibition of tumor growth than the reduction in prolactin [105]. GH stimulates local production of insulin-like growth factor 1 (IGF-1) [105]. IGF polypeptides (also called somatomedins) and various growth factors including epidermal growth factor (EGF), platelet-derived growth factor (PDGF), transforming growth factor (TGF) and others, appear to be involved in the proliferation of both normal and neoplastic cells or phenotypic transformation of cells [105]. In cancer cell lines, somatostatin can reverse the stimulatory effect of EGF on the phosphorylation of the tyrosine kinase portion of the EGF receptor [100, 110]. Some of our superactive analogs of somatostatin such as RC-160 and RC-121 exhibit even higher activity on dephosphorylation of EGF receptor [110].

Thus somatostatin analogs might inhibit prostate cancers by reducing release of GH and prolactin and interfering with the action, signal transmission or secretion of endogenous growth factors [105]. Modern superactive octapeptide analogs of somatostatin, including RC-121, significantly decreased the weight and volume of Dunning R3327H prostate cancers and, when given in combination with once-a-month [D-Trp⁶]LHRH microcapsules, potentiated the effects of the latter [105]. Microcapsules of RC-160 designed for a controlled release of this analog for 30 days also inhibited the growth of Dunning prostate tumors when given alone. Combination of microcapsules of [D-Trp⁶]LHRH with microcapsules of RC-160 resulted in a synergistic potentiation of the inhibition of prostate tumors [105]. Serum GH and PRL levels in treated groups were decreased. Somatomedin-C levels in extracted serum of rats treated for 84 days with somatostatin analog RC-160, alone or in combination with [D-Trp⁶]LHRH, were significantly reduced as compared to control. Histopathological evaluation of prostate cancers in the group of rats which received the combination treatment showed an inhibitory effect on tumors nearly equivalent to that induced by the combination of novantrone and [D-Trp⁶]LHRH [111].

LHRH analogs have also been used for the treatment of benign prostate hyperplasia (BPH) and shown to cause a regression of the prostate [112, 113].

Breast Cancer

About 30% of breast cancers in women are estrogen dependent [114]. Various experimental studies suggest that analogs of LHRH might be useful for treatment of estrogen-dependent breast cancer [30, 31, 115, 116]. In rats bearing the MT/W9A mammary adenocarcinoma, chronic administration of [D-Trp⁶]LHRH

injected daily or once a month as microcapsules, or the antagonist [Ac-D-p-Cl-Phe¹, D-Trp³, D-Arg⁶, D-Ala¹⁰]LHRH decreased tumor weight and volume [115]. Regression of mammary tumors in rats and mice in response to agonists and antagonists of LHRH suggested that these compounds should be considered for a new hormonal therapy for breast cancer in women.

In clinical trials carried out so far, regression of tumor mass and disappearance of metastases in premenopausal and postmenopausal women with breast cancer treated with [D-Trp⁶]LHRH, buserelin or leuprolide have been described [30, 31, 117-122]. Some direct antitumor effects of LHRH analogs on mammary carcinomas are also possible since several groups, including ours, found LHRH receptors in human breast cancers [123, 124]. The use of microcapsules or depot implants of LHRH agonists makes the treatment of breast cancer with LHRH agonists more convenient and ensures patient compliance. It is also possible that treatment with LHRH agonists can be combined with chemotherapy or with peptides that inhibit prolactin release such as somatostatin analogs or prolactin release-inhibiting factor (PIF) [101]. Since prolactin and GH may act as promoters in the development or growth of mammary tumors, the reduction in prolactin and GH levels induced by somatostatin analogs and interference with growth factors, may potentiate inhibition of mammary tumors obtained with LHRH alone [101]. In rats bearing estrogen- and prolactin-dependent MT/W9A mammary adenocarcinoma, daily injections of somatostatin analogs inhibited the growth of this tumor [101]. Microcapsules of somatostatin analog RC-160 also significantly inhibited the growth of the mammary tumor. A significant synergism between the LHRH agonist and the somatostatin analog in the inhibition of tumor growth was demonstrated when both peptides were given together. At present, somatostatin analogs are being tried alone or as adjuncts together with agonistic analogs of LHRH in the treatment of breast cancer in women, depending upon the status of receptors [101].

Pancreatic Cancer

Our experimental studies [31, 101, 125-127] and those of others [128-132] are consistent with the view that the exocrine pancreatic carcinomas are sensitive to G.I. hormones, sex steroids and growth factors. Some clinical observations support the hypothesis that pancreatic cancers might be, in part, sex-hormone dependent [133, 134]. This subject is discussed in detail in the accompanying article [135]. Using animal models of transplanted pancreatic acinar and ductal cancers, we investigated the effect of analogs of LHRH and somatostatin on the growth of these tumors [125]. [D-Trp⁶]LHRH, given daily or injected once a month as microcapsules, significantly decreased tumor weight and volume in both models and suppressed serum levels of testosterone [125]. Some somatostatin analogs also inhibited tumor growth [125]. Similar inhibition of tumor growth by [D-Trp⁶]LHRH and somatostatin analog was obtained in female golden hamsters in which ductal pancreatic cancers were induced by administration of

nitrosamine [126, 127]. This suggests that pancreatic cancers may be sex hormone-sensitive. [D-Trp⁶]LHRH may decrease the growth of pancreatic carcinomas by eliminating the effects of sex steroids. On the basis of these experimental observations that administration of [D-Trp⁶]LHRH inhibits the growth of the ductal and acinar pancreatic cancers, this analog was tried clinically in a small number of patients with inoperable pancreatic cancer [134]. The trial was carried out in collaboration with Gonzalez-Barcena in Mexico City. Some patients showed clinical improvement, reduction in tumor mass and increased survival rate. Gastrin, cholecystokinin and secretin produce hyperplasia and hypertrophy of the exocrine pancreas and might influence the growth of the malignant cells of the pancreas as well [31, 101, 132].

In Syrian hamsters, modern somatostatin analogs, such as D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Trp-NH₂ (RC-160), inhibited the growth of ductal pancreatic tumors [126, 127]. These findings suggest that pancreatic adenocarcinoma may be sensitive to both gastrointestinal and sex hormones. The growth of MIA PaCa-2 human pancreatic cell line in nude mice was also inhibited by RC-160 and [D-Trp⁶]LHRH. The combination of both types of analogs appears to exert the greatest inhibitory effect on pancreatic cancers. Somatostatin analogs reduce the growth of pancreatic ductal and acinar cancers, probably by inhibiting the release and/or stimulatory action of gastrointestinal hormones on tumor cells [31, 101] and by interfering with the action and secretion of growth factors [127]. The stimulatory effect of EGF on the growth of human pancreatic MIA PaCa-2 cells in vitro can be abolished by somatostatin-14 [136]. Somatostatin analogs and [D-Trp⁶]LHRH should be considered for the development of a new hormonal therapy for cancer of the pancreas.

LHRH ANTAGONISTS

While repeated chronic administration of LHRH agonists is required to inhibit LH and FSH release and reduce the levels of sex steroids, similar effects can be obtained with a single administration of LHRH antagonist [31, 42, 102, 137].

Antagonistic analogs of LHRH were developed for contraception [137]. Modern antagonists possess modifications in positions 1, 2, 3, 6, 10 and others [138]. These antagonists act on the same receptor sites as LHRH and cause an immediate inhibition of the release of gonadotropins and sex steroids. Since 1972, hundreds of LHRH antagonists have been synthesized and assayed in animals. Some of the more potent antagonists were also tested in human beings [137-139]. [D-Phe²,D-Trp³,D-Phe⁶]LHRH was the first inhibitory analog found to be active in human beings [140]. The development of LHRH antagonists has been reviewed previously [137,138]. Insertion of D-arginine in the position 6 of LHRH antagonists increases antiovarulatory activity [141]. [Ac-D-p-C1-Phe^{1,2},D-Trp³,D-Arg⁶,D-Ala¹⁰]LHRH (ORG 30276) exhibited antiovarulatory activity at a dose of 1-3µg

[141]. High in vivo potency of D-Arg⁶ analogs can be further increased by the introduction of the very hydrophobic residue, [D-Nal(2)], at position 1 [138-139].

Various studies established that LHRH antagonists can decrease serum gonadotropins, and block ovulation in rats, hamsters and rabbits [42, 137, 142]. In male rats, chronic administration of LHRH antagonists inhibits LH, FSH and testosterone levels and suppresses spermatogenesis [143, 144]. In a series of extensive studies Asch et al, [145, 146] demonstrated inhibitory effects of modern LHRH antagonists on gonadotropin production and ovulation in monkeys.

[N-Ac-D-p-Cl-Phe^{1,2}, D-Trp³, D-Arg⁶, D-Ala¹⁰]LHRH has also been shown to depress serum levels of LH, FSH and testosterone in normal male rhesus monkeys [147]. Several LHRH antagonists have been tested in men and women and demonstrated to be active and powerful enough for clinical use [137, 148, 149].

Clinical Trials with Antagonists of LHRH

In 1977 we demonstrated that large doses of [D-Phe², D-Trp³, D-Phe⁶]LH-RH diminished the gonadotropin response to LHRH in normal men (140). In normal ovulatory women, this antagonist disrupted the menstrual cycle and ovulation did not occur [148]. In oophorectomized or postmenopausal women, we found that the administration of [D-Phe², D-Trp³, D-Phe⁶]LHRH decreased LH and FSH levels up to 24 hours after injection [150].

In another study a still more powerful antagonist, [N-Ac-D-p-Cl-Phe^{1,2}, D-Trp³, D-Phe⁶, D-Ala¹⁰]LHRH, administered on day 12 in a dose of 5mg, blocked ovulation in 6 out of 9 normal women [137]. No breakthrough bleeding occurred. [N-Ac-D-p-Cl-Phe^{1,2}, D-Trp³, D-Phe⁶, D-Ala¹⁰]LHRH was then tested in smaller doses in 10 normally menstruating ovulatory women [149]. Intramuscular administration of the antagonist in doses of 2mg on day 12 of the menstrual cycle inhibited the midcycle surge of LH and FSH and ovulation in 6 out of 10 women. Serum progesterone levels and urinary pregnanediol values in these women were consistently low, and corresponded to anovulatory cycles. These women menstruated 12 to 26 days after injection of the antagonist. Two patients in whom the analog did not abolish the LH mid-cycle surge showed a short luteal phase and premature menstruation. An early decline in progesterone levels suggested that luteolysis took place [149].

Other groups also reported clinical studies on LHRH antagonists [151]. A powerful LHRH antagonist, [Ac-dehydro-Pro¹, D-p-F-Phe², D-Trp^{3,6}]LHRH, administered intravenously in doses of 80µg/kg, (about 5mg/patient), lowered serum levels of LH and FSH within 5 hours in hypergonadotropic postmenopausal women, and decreased the pulses of both gonadotropins. It was also shown that single subcutaneous injection of this antagonist suppresses serum gonadotropin and testosterone levels in normal men without any side effects [152]. Similarly, [N-Ac-D-Nal(2)¹, D-pCl-Phe², D-Trp³, D-hArg (Et)₂₆, D-Ala¹⁰]LHRH (Detirelix; RS-68439) lowered serum LH, FSH and testosterone and caused no systemic side

effects after s.c. injection in normal men in doses of 5-20mg [153].

However, side effects have been observed with most LHRH antagonists with D-Arg in the 6 position. Thus, several such LHRH antagonists in doses of 1.25mg/kg (s.c.) caused a transient allergic reaction in rats (swollen tail, legs, face, ears), suggesting a local and general edema. This reaction is probably due to histamine liberation [154-157]. This phenomenon is species specific: mice, rabbits, dogs or rhesus monkeys do not show this phenomenon [154-157].

In preliminary human tolerance studies in Europe in women, an erythema upon the skin at the side of the subcutaneous injection was observed in some cases after administration of ORG 30276 in doses of 9.5mg[144]. This reaction could be due to histamine liberation. These side effects delayed clinical use of some LHRH antagonists in humans.

Recent LHRH Antagonists

In order to eliminate the undesirable edematogenic effect of the LHRH antagonists containing basic D-amino acids at position 6, exemplified by [Ac-DpCl-Phe^{1,2},D-Trp³,D-Arg⁶,D-Ala¹⁰] LHRH, new analogs with D-ureidoalkyl amino acids such as D-Cit, D-Hci, at position 6 were synthesized in our laboratory and tested in several in vitro and in vivo systems [158]. HPLC analysis revealed that the overall hydrophobicity of the new D-Cit/D-Hci⁶ analogs was similar to that of the basic D-Arg⁶ antagonists. In vitro, most of the new analogs completely inhibited LHRH-mediated LH release in perfused rat pituitary cell systems at an antagonist to LHRH ratio of 5:1. In vivo, the most active peptides, e.g. [Ac-D-Nal(2)¹,D-pCl-Phe²,D-Trp³,D-Cit⁶,D-Ala¹⁰]LHRH caused 100% inhibition of ovulation in cycling rats in doses of 3µg and suppressed the LH level in ovariectomized female rats for 47 hours when administered at levels of 25µg[158]. Characteristically, these peptides did not exert any edematogenic effects even at a dose of 1.5mg/kg. These properties of the D-Cit/D-Hci⁶ antagonists may make them useful clinically.

Other groups have also reported different structural modifications which diminish anaphylactoid toxicity, in particular the histamine releasing activity of the D-Arg⁶ antagonists [159, 160]. In the first approach, Arg was transposed from position 6 to 5 and a hydrophobic D-residue was introduced into position 6 to yield analogues such as [Ac-D-Nal(2)¹,D-pCl-Phe²,D-Pal(3)³,Arg⁵,D-Abu(AA)⁶,D-Ala¹⁰]LHRH [159]. In the other approach, moderately basic acylated L-Lys and D-Lys were introduced into positions 5 and 6, respectively, and Arg⁸ was replaced with alkylated Lys or Orn to obtain peptides such as [Ac-D-Nal(2)¹,D-pCl-Phe²,D-Pal(3)³,Lys(Nic)⁵,D-Lys(Nic)⁶,Lys(Ipr)⁸,D-Ala¹⁰]LH-RH [160]. Both the peptides showed diminished histamine releasing potency.

In view of successful clinical use of LHRH agonists in the treatment of prostate cancer [86-99], breast cancer [117-122], leiomyomas [77-81], endometriosis [68-71] and precocious puberty

[161, 162] as well as experimental efficacy of LHRH antagonists in animal models of prostate cancer [163] and breast cancer [115], the development of LHRH antagonists should continue. The advantage of the antagonists would be based on the fact that they inhibit LH, FSH and sex steroids secretion from the start of the administration. The use of antagonistic analogs of LHRH for the treatment of cancer and other conditions would avoid the transient stimulation of the release of gonadotropins and sex steroids which occurs initially in response to LHRH agonists, thus preventing the temporary clinical "flare-up" of the disease. In addition to contraception, LHRH antagonists could find applications for the treatment of hormone dependent cancers as well as conditions such as endometriosis, leiomyomas and precocious puberty, where the depression of pituitary-gonadal function is desirable. The development of long-acting delivery systems of the LHRH antagonists for once a month or even less frequent administration, will be essential for a practical contraceptive method and other clinical uses. These formulations may be similar to those developed for LHRH agonists [96, 97, 164].

GONADAL PROTECTION AGAINST CHEMOTHERAPY AND RADIATION

Actively dividing cells are more sensitive than resting cells to some chemotherapeutic agents and to radiation [165-167]. Therefore, we administered the agonist [D-Trp⁶]LHRH or the powerful LHRH antagonist [Ac-D-pCl-Phe^{1,2}, D-Trp³, D-Arg⁶, D-Ala¹⁰]LHRH to rats or subhuman primates for 2-3 months before radiation or chemotherapy, in order to suppress the pituitary-gonadal function and maintain inactive testes [165-167]. The baboon pretreated with [D-Trp⁶]LHRH before 4 months of therapy with cytoxan showed a recovery of spermatogenesis, but the control animal given only cytoxan showed azoospermia [166]. Similar results were obtained in rats pretreated with [D-Trp⁶]LHRH or the antagonist before administration of procarbazine (matulane) [167]. When x-radiation was used to inflict gonadal damage, rats pretreated with LHRH antagonist showed a complete recovery of testicular weights and spermatogenesis 3 months after 415 rads and partial recovery after 622 rads; LH and FSH levels returned to normal in both of these groups [165].

These results indicate that radiation and chemotherapy damage the testes temporarily inhibited by LH-RH analogs to a lesser extent than the unsuppressed gonads, and the recovery of reproductive function is accelerated in animals treated with the analogs [165-167]. Our findings have clinical implications and suggest the merit of continued exploration of LHRH analogs as gonadal protectors against chemotherapy and radiation.

CONCLUSIONS

Many clinical approaches are being developed based on analogs of LHRH. The phenomena of pituitary desensitization and inhibition of sex steroid levels by LHRH agonists are being used for treat-

ment of precocious puberty, endometriosis and benign prostate hyperplasia. LHRH agonists are also being tried in contraception and employed in polycystic ovarian disease and IVF-ET or GIFT. Inhibition of the pituitary-gonadal axis forms the basis for oncological applications of LHRH agonists. Various endocrine-dependent or hormone-sensitive tumors can be treated with LHRH analogs. The use of delayed delivery systems based on microcapsules or implants makes the treatment more practical and efficacious. A successful utilization of agonistic analogs of LHRH for the treatment of androgen-dependent prostate cancer has been documented in hundreds of patients. Agonists or antagonists of LHRH might be also beneficial for treatment of breast cancer in premenopausal and postmenopausal women. The development of LHRH antagonists free of edematogenic effects is continuing. Work is in progress on the application of LHRH analogs for the treatment of ovarian cancer, neoplasms of the female genital tract such as leiomyomas, endometrial carcinoma and for protection against gonadal damage during chemotherapy and radiation. Methods based on the analogs of LHRH or somatostatin might supplement or in some cases replace conventional procedures for the treatment of hormone-sensitive cancers.

It is gratifying to see that the discovery of LHRH has led to so many practical clinical uses. The list of these applications might increase further, particularly for the antagonists. Additional new classes of antitumor drugs might be developed based on LHRH.

ACKNOWLEDGEMENTS

The experimental work described in this paper was supported by National Institutes of Health Grants DK7467, CA 40003, CA 40077 and CA 40004 and by the Medical Research Service of the Veterans Administration.

REFERENCES

- Schally, AV, Nair, RMG, Redding, TW and Arimura, A (1971). Isolation of the luteinizing hormone and follicle-stimulating hormone-releasing hormone from porcine hypothalami. *J Biol Chem*, 246, 7230
2. Schally, AV, Arimura, A, Baba, Y, Nair, RMG, Matsuo, H, Redding, TW, Debeljuk, L and White, WF (1971). Isolation and properties of the FSH and LH-releasing hormone. *Biochem Biophys Res Commun*, 43, 393
3. Matsuo, H, Baba, Y, Nair, RMG, Arimura, A and Schally, AV (1971). Structure of the porcine LH and FSH-releasing hormone. Part 1. The proposed amino acid sequence. *Biochem Biophys Res Commun*, 43, 1334
4. Matsuo, H, Baba, Y, Nair, RMG, Arimura, A and Schally, AV (1971). Synthesis of the porcine LH and FSH releasing hormone by the solid-phase method. *Biochem Biophys Res Commun*, 43, 822

5. Schally, AV, Kastin, AJ and Arimura, A (1971). Hypothalamic follicle stimulating hormone (FSH) and luteinizing hormone (LH)-regulating hormone: structure, physiology and clinical studies. *Fertil Steril*, 22, 703
6. Schally, AV, Arimura, A, Kastin, AJ, Matsuo, H, Baba, Y, Redding, TW, Nair, RMG, Debeljuk, L and White, WF (1971). The gonadotropin-releasing hormone: one polypeptide regulates the secretion of luteinizing and follicle stimulating hormones. *Science*, 173, 1036
7. Schally, AV, Coy, DH and Meyers, CA (1978). Hypothalamic regulatory hormones. *Ann Rev Biochem*, 47, 89
8. Kastin, AJ, Schally, AV, Gual, C, Midgley, AR, Jr., Bowers, CY and Diaz-Infante, A, Jr (1969). Stimulation of LH release in men and women by hypothalamic LH-releasing hormone purified from porcine hypothalami. *J Clin Endocrinol Metab*, 29, 1046
9. Kastin, AJ, Schally, AV, Gual, C, Midgley, AR, Jr, Bowers, CY and Gomez-Perez, E (1970). Administration of LH-releasing hormone to selected human subjects. *Am J Obstet Gynecol*, 108, 177
10. Kastin, AJ, Schally, AV, Gual, C and Arimura, A (1972). Release of LH and FSH after administration of synthetic LH-releasing hormone. *J Clin Endocrinol Metab*, 34, 753
11. Zarate, A, Canales, ES, Schally, AV, Ayala-Valdes, L and Kastin, AJ (1972). Successful induction of ovulation with synthetic luteinizing hormone-releasing hormone in anovulatory infertility. *Fertil Steril*, 23, 672
12. Zarate, A, Canales, ES, Soria, J, Gonzalez, A, Schally, AV, Kastin, AJ (1974). Further observations on the therapy of anovulatory infertility with synthetic luteinizing hormone-releasing hormone. *Fertil Steril*, 25, 3
13. Kastin, AJ, Gual, C and Schally, AV (1972). Clinical experience with hypothalamic releasing hormones. II. Luteinizing hormone-releasing hormone and other hypophysiotropic hormones. *Rec Prog Horm Res*, 28, 201
14. Schally, AV and Kastin, AJ (1971). Stimulation and inhibition of fertility through hypothalamic agents. *Drug Ther.*, 1, 29
15. Schally, AV, Coy, DH and Arimura, A (1980). LH-RH agonists and antagonists. *Int J Gynaecol Obstet*, 18, 318
16. Schally, AV, Arimura, A and Coy, DH (1980). Recent approaches to fertility control based on derivatives of LH-RH. In: Munson, P.L., Diczfalussy, E, Glover, J and Olson, RE (eds.). *Vitamins and Hormones*. 38, pp. 258-323. (New York: Academic Press)
17. Redding, TW, Kastin, AJ, Gonzalez-Barcena, D, Coy, DH, Coy, EJ, Schalch, DS and Schally, AV (1973). The half-life, metabolism and excretion of triated luteinizing hormone (LHRH) in man. *J. Clin Endocrinol Metab*, 37, 626
18. Barron, JL, Millar, RP and Searle, D (1982). Metabolic clearance and plasma half-disappearance time of D-Trp⁶ and exogenous LHRH. *J Clin Endocrinol Metab*, 54, 1169
19. Koch, Y, Baram, T, Chobseing, P and Fridkin, M (1974). Enzymatic degradation of luteinizing hormone-releasing hormone (LHRH) by hypothalamic tissue. *Biochem Biophys Res Commun*, 61, 95

20. Nestor, JJ Jr (1984). Development of agonistic LHRH analogs. In: Vickery, BH, Nestor, JJ Jr and Hafez, ESE (eds.). "LHRH and Its Analogs". p3. (Lancaster: MTP Press)
21. Coy, DH, Coy, EJ, Schally, AV, Vilchez-Martinez, JA, Hirotsu, Y and Arimura, A (1974). Synthesis and biological properties of [D-Ala⁶, desGly-NH₂¹⁰]-LH-RH ethylamide, a peptide with greatly enhanced LH- and FSH-releasing activity. *Biochem Biophys Res Commun*, 57, 335
22. Coy, DH, Vilchez-Martinez, JA, Coy, EJ and Schally, AV (1976). Analogs of luteinizing hormone-releasing hormone with increased biological activity produced by D-amino acid substitutions in position 6. *J Med Chem*, 19, 423
23. Fujino, M, Fukuda, T, Shinagawa, S, Kobayashi, S, Yamazaki, I, Nakayama, R, Seely, JH, White, WF and Ripell, RH (1974). Synthetic analogs of luteinizing hormone-releasing hormone (LH-RH) substituted in position 6 and 10. *Biochem Biophys Res Commun*, 60, 406
24. Fujino, M, Yamazaki, I, Kobayashi, S, Fukada, T, Shinagawa, S, Nakayama, R, White, WF and Rippel, RH (1974). Some analogs of luteinizing hormone-releasing hormone (LH-RH) having intense ovulation-inducing activity. *Biochem Biophys Res Commun*, 57, 1248
25. Sandow, J (1976). Hypothalamic control of anterior pituitary hormone secretion: physiological aspects. In: Charro-Salgado, A, Fernandez-Durango, R and Lopez del Campo, JG (eds.). "Basic Applications and Clinical Uses of Hypothalamic Hormones". p 113. (Amsterdam: Excerpta Medica)
26. Sandow, J, von Rechenberg, W, Jerzabek, G and Stoll, W (1978). Pituitary gonadotropin inhibition by a highly active analog of luteinizing hormone-releasing hormone. *Fertil Steril*, 20, 205
27. Schally, AV, Kastin, AJ, Coy, DH (1976). LH-releasing hormone and its analogues: recent basic and clinical investigations. *Int J Fertil*, 21, 1
28. Corbin, A (1982). From contraception to cancer: a review of the therapeutic applications of LH-RH analogues as antitumor agents. *Yale J Biol Med*, 55, 27
29. Johnson, B, Gendrich, RL, and White, WF (1976). Delay of puberty and inhibition of reproductive processes in the rat by a gonadotropin-releasing hormone agonist analog. *Fertil Steril*, 27, 853
30. Schally, AV, Redding, T and Comaru-Schally, AM (1984). Potential use of luteinizing hormone-releasing hormones in the treatment of hormone-sensitive neoplasms. *Cancer Treat Rep*, 68, 281
31. Schally, AV, Comaru-Schally, AM and Redding, T (1984). Antitumor effects of analogs of hypothalamic hormones in endocrine-dependent cancers. *Proc Soc Exp Biol Med*, 175, 259
32. Auclair, C, Kelly, PA, Labrie, R, Coy, DH and Schally, AV (1977). Inhibition of testicular luteinizing hormone receptor level by treatment with a potent luteinizing hormone-releasing hormone agonist or human chorionic gonadotropin. *Biochem Biophys Res Commun*, 75, 855

33. Jacobi, GH and Wenderoth, UK (). Gonadotropin releasing hormone analogues for prostate cancer: untoward side effects of high-dose regimens acquire a therapeutical dimension. *Eur J Urol*, 8, 129
34. Borgmann, V, Hardt, W, Schmidt-Gollwitzer, M, Adenauer, H and Nagel, R (1982). Sustained suppression of testosterone production by the luteinizing hormone-releasing hormone agonist Buserelin in patients with advanced prostate carcinoma. *Lancet*, i, 1097
35. Bergquist, C, Nillius, SJ, Bergh, T, Skarin, G and Wide, W (1979). Inhibitory effects on gonadotropin secretion and gonadal function in men during chronic treatment with a potent stimulatory luteinizing hormone-releasing hormone analogue. *Acta Endocrinol*, 91, 601
36. Maynard, PV and Nicholson, RI (1979). Effects of high doses of a series of new luteinizing hormone-releasing hormone analogues in intact female rats. *Brit J Cancer*, 39, 274
37. Corbin, A, Beattie, CW, Tracy, J, Jones, R, Foell, TJ, Yardley, J and Rees, RWA (1978). The anti-reproductive pharmacology of LH-RH and antagonistic analogues. *Int J Fertil*, 23, 81
38. Bergquist, C, Nillius, SJ and Wide, L (1979). Intranasal gonadotropin-releasing hormone agonist as a contraceptive agent. *Lancet*, 2, 215
39. Bex, FJ, Corbin, A and France, E (1982). Resistance of the mouse to the antifertility effects of LH-RH agonists. *Life Sci*, 30, 1263
40. Bokser, L, Zalatnai, A and Schally, AV (1988). Inhibition of pituitary-gonadal axis in mice by long term administration of D-Trp-6-LH-RH microcapsules. *Peptides*, (submitted)
41. Nillius, SJ, Bergquist, CH and Wide, L (1978). Inhibition of ovulation in women by chronic treatment with a stimulatory LHRH analogue- a new approach to birth control? *Contraception*, 17, 537
42. Schally, AV (1981). LH-RH analogs in contraception and cancer. In Labrie, F, Belanger, A and Dupont, A (eds.). "LHRH and its Analogues". p. 3. (Amsterdam: Elsevier Science)
43. Meldrum, DR, Tsao, Z, Monroe, SE, Braunstein, GD, Sladek, J, Lu, JKH, Vale, W, Rivier, J, Judd, HL and Chang, RJ (1984). Stimulation of LH fragments with reduced bioactivity following GnRH agonist administration in women. *J Clin Endocrinol Metab*, 58, 755
44. Chang, RJ, Laufer, LH, Meldrum, DR, de Fazio, J, Lu, JKH, Vale, WW, Rivier, JE and Judd, HL (1983). Steroid secretion in polycystic ovarian disease after ovarian suppression by a long-acting gonadotropin-releasing hormone agonist. *J Clin Endocrinol Metab*, 56, 897
45. Andreyko, J, Monroe, S, Blumenfeld, Z and Jaffe, R (1985). Treatment of hirsutism with a GnRH agonist. Program of the 32 Meeting of the Society for Gynecologic Investigation, Phoenix, Arizona, p. 41 (Abstract)
46. Shaw, RW and Fraser, HM (1984). Use of a superactive luteinizing hormone releasing-hormone (LHRH) agonist in the treatment of menorrhagia. *Brit J Obstet Gynaecol*, 91, 913

47. Maheux, R, Guilloteau, C, Lemay, A, Bastide, A, Fazekas, ATA (1984). Regression of leiomyomata uteri following hypoestrogenism induced by repetitive LHRH agonist treatment: preliminary report. *Fertil Steril*, 42, 644
48. Crowley, WF and McArthur, JW (1980). Stimulation of the normal menstrual cycle in Kallmann's syndrome by pulsatile administration of luteinizing hormone-releasing hormone (LH-RH). *J Clin Endocrinol Metabol*, 51, 173
49. Reid, RL, Leopold, GR and Yen, SSC (1981). Induction of ovulation and pregnancy with pulsatile luteinizing hormone releasing factor: dosage and mode of delivery. *Fertil Steril*, 36, 553
50. Miller, DS, Reid, RR, Cetel, NS, Rebar, RW and Yen, SC (1983). Pulsatile administration of low-dose gonadotropin-releasing hormone: ovulation and pregnancy in women with hypothalamic amenorrhea. *J Am Med Assoc*, 250, 2937
51. Happ, J, Kollmann, F, Krawehl, C, Neubauer, M, Krause, U, Demisch, K, Sandow, J, von Richenberg, W and Bever, J (1978). Treatment of cryptorchidism with pernasal gonadotropin-releasing hormone therapy. *Fertil Steril*, 29, 546
52. De Muninck Keizer-Schrama, SMPF, Hazebroek, FWJ, Matroos, AW, Drop, SLS, Molenaar, JC and Visser, HKA (1986). Double-blind, placebo-controlled study of luteinizing-hormone-releasing hormone nasal spray in treatment of undescended testes. *Lancet*, 1, 876
53. Corbin, A and Bex, FJ (1981). In: Zatuchni, GI, Shelton, JD and Sciarra, JJ (eds.). "LHRH Peptides as Female and Male Contraceptives". p66. (Philadelphia: Harper and Row)
54. Monroe, SE, Blumenfeld, Z, Andreyko, JL, Schriock, E, Henzl, MR and Jaffe, RB (1986). Dose-dependent inhibition of pituitary-ovarian function during administration of a gonadotropin-releasing hormone agonistic analog (nafarelin). *J Clin Endocrinol Metab*, 63, 1334
55. Gudmundsson, JA (1986). Peptide contraception in women, a clinical and metabolic study. Doctoral thesis Uppsala University
56. Schmidt-Gollwitzer, M, Hardt, W, Schmidt-Gollwitzer, K, von der Ohe, M and Nevinny-Stickel, J (1981). Influence of the LH-RH analog busserlin on cyclic ovarian function and on endometrium. A new approach to fertility control? *Contraception*, 23, 187
57. Schmidt-Gollwitzer, M, Hardt, W, Schmidt-Gollwitzer, K, von der Ohe, M (1981). In: Zatuchni, GI, Shelton, JD and Sciarra, JJ (eds.). "LHRH Peptides as Female and Male Contraceptives" p. 199. (Philadelphia: Harper and Row)
58. Bergquist, C, Nillius, SJ, Wide, L and Lindgren, A (1981). Endometrial patterns in women on chronic luteinizing hormone-releasing hormone agonist treatment for contraception. *Fertil Steril*, 36, 339
59. Casper, FR and Yen, SSC (1979). Induction of luteolysis in the human with a long-acting analog of luteinizing hormone-releasing hormone factor. *Science*, 205, 408
60. Sheehan, KL and Yen, SSC (1981). In: Zatuchni, GI, Shelton, JD, Sciarra, JJ (eds.). "LHRH Peptides as Female and Male Contraceptives". p.237. (Philadelphia: Harper and Row)

61. Casper, RF, Sheehan, KL and Yen SCC (1980). Chorionic gonadotropin prevents LRF-agonist-induced luteolysis in the human. *Contraception*, 21, 471
62. Tolis, G, Comaru-Schally, AM, Mehta, AE and Schally, AV (1981). Failure to interrupt established pregnancy in humans by D-tryptophan-6-luteinizing hormone-releasing hormone. *Fertil Steril*, 36, 241
63. Linde, R, Doelle, GC, Alexander, N, Kirchner, F, Vale, W, Rivier, J and Rabin, D (1981). Reversible inhibition of testicular steroidogenesis and spermatogenesis by a potent gonadotropin-releasing hormone agonist in normal man. *N Eng J Med*, 305, 663
64. Heber, D, Swerdloff, RS (1980). Male Contraception: synergism of gonadotropin-releasing hormone analog and testosterone in suppressing gonadotropin. *Science*, 209, 9936
65. Bhasin, S, Steiner, BS, Heber, D, Handelsman, D and Swerdloff, RS (1985). Hormonal effects of GnRH agonist in the human male: II. Effects of long term combined treatment with GnRH agonist and androgen. *J Clin Endocrinol Metab*, 60, 998-1003
66. Doelle, GC, Alexander, NA, Evans, RM, Linde, R, Rivier, J, Vale, W and Rabin, D (1983). Combined treatment with LHH agonist and testosterone in man: reversible oligozoospermia without impotence. *J Androl*, 4, 298
67. Schally, AV and Comaru-Schally, AM (1987). Male contraception involving testosterone supplementation: possible increased risks of prostate cancer? *Lancet*, 1, 418
68. Meldrum, DR, Chang, RJ, Lu, J, Vale, W, Rivier, J and Judd, HL (1982). Medical oophorectomy using a long-acting GnRH agonist: a possible new approach to the treatment of endometriosis. *J Clin Endocrinol Metabol*, 54, 1081
69. Lemay, A and Quesnel, G (1982). Potential new treatment of endometriosis: reversible inhibition of pituitary ovarian function by chronic intranasal administration of a luteinizing hormone releasing hormone (LH-RH) agonist. *Fertil Steril*, 38, 376
70. Zorn, JR, Tanger, CH, Roger, M, Grenier, J, Comaru-Schally, AM and Schally, AV (1980). Therapeutic hypogonadism induced by a delayed-release preparation of microcapsules of D-Trp⁶-luteinizing hormone-releasing hormone: a preliminary study in eight women with endometriosis. *Int J Fertil*, 31, 11
71. Henzl, MR, Corson, SL, Moghissi, K, Buttram, VC, Bergquist C and Jacobson, J (1988). Administration of nasal nafarelin as compared with oral danazol for endometriosis: a multicenter double-blind comparative clinical trial. *Eng Med*, 318, 485
72. Ayalon, D, Ben-David, M, Wohl, R, Jaffe, R, Vagman, I, Eckstein, N, Limor, R, Comaru-Schally, AM and Schally, AV (1988). Induction of ovulation with D-Trp-6-LH-RH combined with purified FSH in patients with polycystic ovarian disease. *J Clin Endocrinol Metab*, (in press)
73. Fleming R, Haxton, MJ, Hamilton, MPR, McCune, GS, Black, WP, MacNaughton, MC and Coutts, JRT (1985). Successful treatment of infertile women with oligomenorrhoea using a combination of an LHRH agonist and exogenous gonadotrophin. *Brit J Obstet Gynaecol*, 92, 369

74. Porter, RN, Smith, W, Craft, IL, Abdulwabid, NA and Jacobs, HS. Induction of ovulation for in vitro fertilization using Buserelin and gonadotropins. *Lancet*, 2, 1284
75. Wildt, L, Diedrich, K, van der Ven, H, Hasanis, AL, Hubner, H, Klazen, R (1986). Ovarian hyperstimulation for in vitro fertilization controlled by GnRH agonist administered in combination with human menopausal gonadotrophins. *Human Reprod*, 1, 15
76. Zorn, JR, Boyer, P and Guichard, A (1987). Never on a Sunday: programming for IVF-ET and GIFT. *Lancet*, 2, 385
77. Filicori, M, Hall, DA, Loughlin, JS, Rivier, J, Vale, W and Crowley, WF (1983). A conservative approach to the management of uterine leiomyoma: pituitary desensitization by luteinizing hormone-releasing hormone analogue. *Am J Obstet Gynecol*, 147, 726
78. Healy, DL, Fraser, HM and Lawson, SL (1984). Shrinkage of a uterine fibroid after subcutaneous infusion of a LHRH agonist. *Brit Med J*, 289, 1267
79. Coddington, CC, Collins, RL, Shawker, TH, Anderson, R, Loriaux, DL and Winkel, CA (1986). Long-acting gonadotropin hormone-releasing hormone analog used to treat uteri. *Fertil Steril*, 45, 624
80. Van Leusden, HAIM (1986). Rapid reduction of uterine myomas after short-term treatment with microencapsulated D-Trp⁶-LH-RH. *Lancet*, 2, 1213
81. Perl, V, Marquez, J, Schally, AV, Comaru-Schally, AM, Leal, G, Zacharias, S and Gomez-Kira, C (1987). Treatment of leiomyomata uteri with D-Trp-6-LH-RH. *Fertil Steril*, 48, 383
82. George, M, Lhomme, C, Lefort, J, Gras, C, Comaru-Schally, AM and Schally, AV (1989). Long term use of an LH-RH agonist in the management of uterine leiomyomas: a study of 17 cases. *Int J Fertil*, 34, 19
83. Parmar, H, Nicoll, J, Stockdale, A, Cassoni, A, Phillips, RH, Lightman, SL and Schally, AV (1985). Advanced ovarian carcinoma: response to the agonist D-Trp-6-LH-RH. *Cancer Treat Rep*, 69, 1341
84. Parmar, H, Phillips, RH, Rustin, G, Hanham, IW, Schally, AV and Lightman, SL (1988). Response to D-Trp-6-Lh-RH (Decapeptyl) microcapsules in advanced ovarian cancer. *Brit Med J*, 296, 1229
85. Redding, TW and Schally, AV (1981). Inhibition of prostate tumor growth in two rat models by chronic administration of D-Trp analogue of luteinizing hormone-releasing hormone. *Proc Natl Acad Sci USA*, 78, 6509
86. Tolis, G, Ackman, D, Stellos, A, Mehta, A, Labrie, F, Fazekas, ATA, Comaru-Schally, AM and Schally, AV (1982). Tumor growth inhibition in patients with prostatic carcinoma treated with luteinizing hormone-releasing hormone agonists. *Proc Natl Acad Sci USA*, 79, 1658
87. Warner, B, Worgul, TJ, Drago, J, Demers, L, Dufau, M, Max, D and Santen, RJ (1983). Effect of high dose D-Leucine⁶-Gonadotropin-releasing hormone proethylamide on the hypothalamic-pituitary testicular axis in patients with prostatic cancer. *J Clin Invest*, 17, 1842

88. Borgmann, V, Nagel, R, Al-Abadi, H and Schmidt-Gollwitzer, M (1983). Treatment of prostatic cancer with LH-RH analogs. *Prostate*, 99, 553
89. Mathe, G, VoVan, ML, Duchier, J, Missett, JL, Morin, P, Keiling, R, Schwarzenberg, L, Kerbrat, P, Achille, E, Tronc, JC, Machover, D, Fendler, JP, Pappo, E, Metz, R, Prevot, G, Comaru-Schally, AM and Schally, AV (1984). An oriented phase-II trial of D-Trp-6-LH-RH in patients with prostatic carcinoma. *Med Oncol Tumor Pharmacother*, 1, 119
90. Waxman, JH, Wass, JAH, Hendry, WF, Whitfield, HN, Besser, GM, Malpas, JS and Oliver, RTD (1983). Treatment with gonadotrophin releasing hormone analog in advanced prostatic cancer. *Brit Med J*, 286, 1309
91. Walker, KJ, Nicholson, RI, Turkes, AO, Turkes, A, Robinson, M, Crispin, Z and Dris, S (1983). Therapeutic potential of the LH-RH agonist, ICI 118630, in the treatment of advanced prostatic carcinoma. *Lancet*, 2, 413
92. Ahmed, SR, Brooman, PJC, Shalmet, SM, Howell, A, Blacklock, NJ and Rickards, D (1983). Treatment of advanced prostatic cancer with LH-RH analog ICI 118630: clinical response and hormonal mechanisms. *Lancet*, 2, 415
93. Allen, JM, O'Shea, JP, Mashiter, K, Williams, G and Bloom, SR (1983). Advanced carcinoma of the prostate: treatment with a gonadotrophin releasing hormone agonist. *Brit Med J*, 286, 1607
94. Gonzalez-Barcena, D, Perez-Sanchez, P, Ureta-Sanchez, S, Dominguez, HB, Graef-Sanchez, A, Morales, MB, Comaru-Schally, AM and Schally, AV (1985). Treatment of advanced prostatic carcinoma with D-Trp-6-LH-RH. *Prostate*, 7, 21
95. Mathe, G, Schally, AV, Comaru-Schally, AM, Mauvernay, RY, VoVan, ML, Machover, D, Missett, JL, Court, B, Bochard, P, Duchier, J, Morin, P, Keiling, R, Schwarzenberg, L, Kerbrat, P, Achille, E, Tronc, JC, Fendler, JP, Pappo, E, Metz, R and Prevot, G (1986). A phase II trial with D-Trp-6-LH-RH in prostatic carcinoma: a comparison with other hormonal agents. *Prostate*, 9, 327
96. Redding, TW, Schally, AV, Tice, TR and Meyers, WE (1984). Long-acting delivery systems for peptides: inhibition of rat prostate tumors by controlled release of (D-Trp-6) luteinizing hormone-releasing hormone from injectable microcapsules. *Proc Natl Acad Sci USA*, 81, 5845
97. Mason-Garcia, M, Vigh, S, Comaru-Schally, AM, Redding, TW, Somogyvari-Vigh, A, Horvath, J and Schally, AV (1985). Radioimmunoassay for D-Trp⁶ analog of luteinizing hormone-releasing hormone: measurement of serum levels after administration of long-acting microcapsule formulation. *Proc Natl Acad Sci USA*, 82, 1547
98. Roger, M, Duchier, J, Lahlou, N, Nahoul, K and Schally, AV (1985). Treatment of prostatic carcinoma with D-Trp-6-LH-RH: plasma hormone levels after daily subcutaneous injections and periodic administration of delayed release preparation. *Prostate*, 7, 271

99. Parmar, H, Lightman, SL, Allen, L, Phillips, RH, Edwards, L and Schally, AV (1985). Randomised controlled study of orchidectomy vs long-acting D-Trp-6-LH-RH microcapsules in advanced prostatic carcinoma. *Lancet*, 2, 1202
100. Schally, AV, Cai, RZ, Torres-Aleman, I, Redding, TW, Szoke, B, Fu, D, Hierowski, MT, Colaluca, J and Kopturek, S (1986). Endocrine, gastrointestinal and antitumor activity of somatostatin analogs: Washington Spring Peptide Symposium. In: Moody TW (Ed.). "Neural and Endocrine Peptides and Receptors". p.73. (New York: Plenum)
101. Schally, AV, Redding, TW, Cai, RZ, Paz, JI, Ben-David, M and Comaru-Schally, AM (1987). Somatostatin analogs in the treatment of various experimental tumors. In: Klijn, JM, Paridaens, R and Foekens, JA (eds). "International Symposium on Hormonal Manipulation of Cancer: peptides, growth factors and new (anti) steroidal agents". p.431. (New York: Raven Press)
102. Schally, AV, Redding, TW, Paz-Bouza, JI, Comaru-Schally, AM and Mathe, G (1987). Current concept for improving treatment of prostate cancer based on combination of LH-RH agonists with other agents. In: Murphy, GP, Khoury, S, Kuss, R, Chatelain, C and Denis, L (eds.). "Prostate cancer part A: research, endocrine treatment and histopathology". p.173. (New York: Alan R. Liss)
103. Schally, AV and Redding, TW (1985). Combination of long-acting microcapsules of the D-tryptophan-6 analog of luteinizing hormone-releasing hormone with chemotherapy: investigation in the rat prostate cancer model. *Proc Natl Acad Sci USA*, 82, 2498
104. Schally, AV, Kook, AI, Monje, E, Redding, TW and Paz-Bouza, JI (1986). Combination of a long-acting delivery system for luteinizing hormone-releasing hormone agonist with Navantrone chemotherapy: increased efficacy in the rat prostate cancer model. *Proc Natl Acad Sci USA*, 83, 8764
105. Schally, AV, and Redding, TW (1987). Somatostatin analogs as adjuncts to agonists of luteinizing hormone-releasing hormone in the treatment of experimental prostate cancer. *Proc Natl Acad Sci USA*, 84, 7275
106. Isaacs, JT and Coffey, DS (1981). Adaptation versus selection as the mechanism responsible for the relapse of prostatic cancer to androgen ablation therapy studied in the Dunning R-3327-H adenocarcinoma. *Cancer Res*, 41, 5070
107. Isaacs, JT (1984). The timing of androgen ablation therapy and/or chemotherapy in the treatment of prostatic cancer. *Prostate*, 5, 1
108. Paz-Bouza, JI, Schor, NA, Monje, E, Redding, TW and Schally, AV (1987). Histological findings in the rat prostate cancer model during treatment with a luteinizing hormone-releasing hormone agonist and Novantrone. *Prostate*, 10, 291
109. Assimos, D, Smith, C, Lee, C and Grayhack, JT (1984). Action of prolactin in regressing prostate: independent of action mediation by androgen receptors. *Prostate*, 5, 589
110. Hierowski, MT, Liebow, C, duSapin, K and Schally, AV (1985). Stimulation by somatostatin of dephosphorylation of

membrane proteins in pancreatic cancer MIA PaCa-2 cell line. FEBS Lett, 179, 252

111. Zalathnai, A, Paz-Bouza, JI, Redding, TW and Schally, AV (1988). Histological changes in the rat prostate cancer model after treatment with somatostatin analogs and D-Trp-6-LH-RH. Prostate (in press)

112. Gabrilove, JL, Levine, Ac, Kirschenbaum, A and Droller, M (1987). Effect of a GNRH analogue (leuprolide) on benign prostatic hypertrophy. J Clin Endocrinol Metab, 64, 1331

113. Peters, CA and Walsh, PC (1987). The effect of nafarelin acetate, a luteinizing-hormone-releasing hormone agonist, on benign prostatic hyperplasia. N Engl J Med, 317, 599

114. McGuire, WL (1979). Steroid Hormone receptors and disease: breast cancer. Proc Soc Exp Biol Med, 162, 22

115. Redding, TW and Schally, AV (1983). Inhibition of mammary tumor growth in rats and mice by administration of agonistic and antagonistic analogs of luteinizing hormone-releasing hormone. Proc Natl Acad Sci USA, 80, 1459

116. Nicholson, RI and Maynard, PV (1979). Anti-tumor activity of ICI 118630, a new potent luteinizing hormone-releasing hormone agonist. Brit J Cancer, 39, 268

117. Klijn, JGM and DeJong, FH (1982). Treatment with a luteinizing-hormone releasing hormone analogue (Buserelin) in premenopausal patients with metastatic breast cancer. Lancet, 2, 1213

118. Harvey, HA, Lipton, A and Max, D (1983). LH-RH analogs for human mammary carcinoma. In: Vickery, BH, Nestor, JJ Jr. and Hafez, ESE (eds.). "LHRH and Its Analogs - Contraceptive and Therapeutic Applications". p.329. (Lancaster: MTP Press)

119. Manni, A, Santen, R, Harvey, H, Lipton, A and Max, D (1986). Treatment of breast cancer with gonadotropin-releasing hormone. Endocr Rev, 7, 89

120. Walker, KJ, Turkes, A, Williams, MR, Blamey, RW and Nicholson, RI (1986). Preliminary endocrinological evaluation of a sustained-release formulation of the LH-releasing hormone agonist D-Ser(Bu^t)⁶Azgly¹⁰LH-RH in premenopausal women with advanced breast cancer. J Endocrinol, 111, 349

121. Plowman, PN, Nicholson, RI and Walker, KJ (1986). Remission of postmenopausal breast cancer during treatment with the luteinising hormone releasing hormone agonist ICI 118630. Brit J Cancer, 54, 903

122. Williams, MR, Walker, KJ, Turkes, A, Blamey, RW and Nicholson, RI (1986). The use of an LH-RH agonist (ICI 118630, Zoladex) in advanced premenopausal breast cancer. Brit J Cancer, 53, 629

123. Miller, WR, Scott, WN, Morris, R, Fraser, HM and Sharpe, RM (1985). Growth of human breast cancer cells inhibited by a luteinizing hormone-releasing hormone agonist. Nature, 313, 231

124. Eidne, KA, Flanagan, CA and Millar, RP (1985). Gonadotropin-releasing hormone binding sites in human breast carcinoma. Science, 229, 989

125. Redding, TW and Schally, AV (1984). Inhibition of growth of

- pancreatic carcinomas in animal models by analogs of hypothalamic hormones. *Proc Natl Acad Sci USA*, 81, 248
126. Paz-Bouza, JI, Redding, TW and Schally, AV (1987). Treatment of nitrosamine-induced pancreatic tumors in hamsters with analogs of somatostatin and luteinizing hormone-releasing hormone. *Proc Natl Acad Sci USA*, 84, 1112
127. Zalatnai, A and Schally AV (1989). Responsiveness of the hamster pancreatic cancer to treatment with microcapsules of D-Trp-6-LH-RH and somatostatin analog RC-160: histological evidence of improvement. *Int J Pancreatol*, 4, 149
128. Greenway, BA (1987). Carcinoma of the exocrine pancreas: a sex hormone responsive tumour? *Brit J Surg*, 74, 441
129. Greenway, B, Duke, D, Pym, B, Iqbal, MJ, Johnson, PJ and Williams, R (1982). The control of human pancreatic adenocarcinoma xenografts in nude mice by hormone therapy. *Brit J Surg*, 69, 595
130. Sandberg, A and Rosenthal, H (1979). Steroid receptors in exocrine glands: the pancreas and the prostate. *J Steroid Biochem*, 11, 293
131. Chester, JF, Gaissert, HA, Ross, JS and Malt, RA (1986). Pancreatic cancer in the Syrian hamster induced by N-Nitrobis (2-oxopropyl) amine: cocarcinogenic effect of epidermal growth factor. *Cancer Res*, 46, 2954
132. Johnson, LR (1981). Effects of gastrointestinal hormones on pancreatic growth. *Cancer*, 47, 1640
133. Theve, NO, Pousette, A and Carlstrom, K (1983). Adenocarcinoma of the pancreas- a hormone sensitive tumor? A preliminary report on Nolvadex treatment. *Clin Oncol*, 9, 193
134. Gonzalez-Barcena, D, Rangel-Garcia, NE, Nerez-Sanchez, PL, Gutierrez-Damperio, C, Garcia-Carrasco, F, Comaru-Schally, AM and Schally, AV (1986). Response to D-Trp-6-LH-RH in advanced adenocarcinoma of pancreas. *Lancet*, 2, 154
135. Comaru-Schally, AM and Schally, AV (1988). LH-RH agonists as adjuncts to somatostatin analogs in the treatment of pancreatic cancer. In: Vickery, BH and Lunenfeld, B (eds.). "GnRH Analogues in Cancer and Human Reproduction". (In press)
136. Liebow, C, Hierowski, M and DuSapin, K (1986). Hormonal control of pancreatic cancer growth. *Pancreas*, 1, 44
137. Schally, AV (1983). Current status of antagonistic analogs of LH-RH as a contraceptive method in the female. *Research Frontiers in Fertility Regulation (PARFR)*, 2, p.1, Northwestern University Press
138. Karten, MJ and Rivier, JE (1986). Gonadotropin-releasing hormone analog design. Structure-function studies toward the development of agonists and antagonists: rationale and perspective. *Endocr Rev*, 7, 44
139. Vickery, BH (1986). Comparison of the potential for therapeutic utilities with gonadotropin-releasing hormone agonists and antagonists. *Endocr Rev*, 7, 115
140. Gonzalez-Barcena, D, Kastin, AJ, Coy, DH, Nikolics, K and Schally, AV (1977). Suppression of gonadotropin release in man by an inhibitory analogue of luteinizing hormone-releasing hormone. *Lancet*, 2, 997

141. Coy, DH, Horvath, A, Nekola, MV, Coy, EJ, Erchegyi, J and Schally, AV (1982). Peptide antagonists of LH-RH: large increases in antioviulatory activities produced by basic D-amino acids in the six position. *Endocrinology*, 110, 1445
142. Schally, AV and Coy, DH (), Stimulatory and inhibitory analogs of LH-releasing hormone: basic and clinical studies. In: McCann, SM and Dhindsa, DS (eds.). "Role of Peptides and Proteins in Control of Reproduction". p.89. (New York: Elsevier Biomedical)
143. Rivier, C, Rivier, J and Vale, W (1981). Effect of a potent GnRH antagonist and testosterone propionate on mating behavior and fertility in the male rat. *Endocrinology*, 108, 1998
144. Debeljuk, L, and Schally, AV (1986). Antifertility Effects of a potent LH-RH antagonist in male and female rats. *Int J Fertil*, 31, 284
145. Borghi, MR, Niesvisky, R, Coy, DH, Balmaceda, JP, Schally, AV and Asch, RH (1983). Administration of agonistic and antagonistic analogues of LH-RH induce anovulation in *Macaca fascicularis*. *Contraception*, 27, 619
146. Asch, RH (1984). The effects of LH-RH inhibitory analogues in the non-human primate. Basic studies in contraception and their direct application for future clinical uses in both sexes. In: Labrie, F, Belanger, A and Dupont, A (eds.). "LHRH and its Analogues". p.188. (Amsterdam: Elsevier Science)
147. Burgos-Briceno, LA, Schally, Av, Bartke, A and Asch, RH (1984). Inhibition of serum luteinizing hormone and testosterone with an inhibitory analogue of luteinizing hormone-releasing hormone in adult male rhesus monkeys. *J Clin Endocr Metab*, 59, 601
148. Canales, ES, Montvelinsky, H, Zarate, A, Kastin, AJ, Coy, DH and Schally, AV (1980). Suppressive effect of an inhibitory LHRH analog on the gonadotropin response to LHRH in normal women. *Int J Fertil*, 25, 190
149. Zarate, A, Canales, ES, Sthory, I, Coy, DH, Comaru-Schally, AM and Schally, AV (1981). Anovulatory effect of a LHRH antagonist in women. *Contraception*, 24, 315
150. Gonzalez-Barcena, D, Trevino-Ortiz, H, Gordon, F, Kastin, AJ, Coy, DH and Schally, AV (1980). Influence of LH-RH agonists and antagonists on gonadotropin release in humans. *Int J Fertil*, 25, 185
151. Cetel, NS, Rivier, J, Vale, W and Yen, SSC (1983). The dynamics of gonadotropin inhibition in women induced by an antagonistic analog of gonadotropin-releasing hormone. *J Clin Endocr Metab*, 57, 62
152. Pavlou, N, Debold, C, Islan, P, Wakefield, G, Rivier, J, Vale, W and Rabin, D (1986). Single subcutaneous doses of a luteinizing hormone-releasing hormone antagonist suppresses serum gonadotropin and testosterone levels in normal men. *J Clin Endocrinol Metab*, 63, 303
153. Pavlou, SN, Wakefield, GB, Island, DP, Hoffman, PG, LePage, ME, Chan, RL, Nerenberg, CA and Kovacs, WJ (1987). Suppression of pituitary-gonadal function by a potent new luteinizing hormone-releasing hormone antagonist in normal men. *J Clin Endocrinol Metab*, 64, 931

154. Schmidt, F, Sundaram, K, Thau, RB and Bardin, CW (1984). (Ac-D-Nal(2)¹, 4FD-Phe², D-Trp³, D-Arg⁶)-LRH, potent antagonist of LHRH, produces transient edema and behavioral changes in rats. *Contraception*, 29, 282
155. Hahn, DW, McGuire, JL, Vale, WW and Rivier, J (1984). Reproductive/endocrine and anaphylactoid properties of an LHRH-antagonist, ORF 18260 (Ac-DNAL¹(2), 4FDPhe², D-Trp³, D-Arg⁶)-GnRH. *Life Sci*, 37, 505
156. Morgan, JE, O'Neil, CE, Coy, DH, Hocart, SJ and Nekola, MV (1986). Antagonistic analogs of luteinizing hormone-releasing hormone are mast cell secretagogues. *Int Arch All Appl Immunol*, 80, 70
157. Hook, WA, Karten, M and Siraganian, RP (1985). Histamine release by structural analogs of LHRH. *Fed Proc*, 44, 1323
158. Bajusz, S, Kovacs, M, Gazdag, M, Bokser, L, Karashima, T, Csernus, VJ, Janaky, T, Guoth, J and Schally, AV (1988). Highly potent antagonists of luteinizing hormone-releasing hormone (LHRH) free of edematogenic effects. *Proc Natl Acad Sci USA*, 85, 1637
159. Rivier, JE, Porter, J, Rivier, CL, Perrin, M, Corrigan, A, Hook, WA, Siraganian, RP and Vale WW (1986). New effective gonadotropin releasing hormone antagonists with minimal potency for histamine release in vitro. *J Med Chem*, 29, 1864
160. Ljungqvist, A, Feng, D-M, Tang, P-FL, Kubota, M, Okamoto, T, Zhang, Y, Bowers, Cy, Hook, WA and Folkers, K (1987). Design, synthesis and bioassays of antagonists of LH-RH which have high antioviulatory activity and release negligible histamine. *Biochem Biophys Res Commun*, 148, 849
161. Laron, Z, Kauli, R, Ben, Zeev, Z, Comaru-Schally, AM and Schally, AV (1981). D-Trp-6-analogue of luteinizing hormone releasing hormone in combination with cyproterone acetate to treat precocious puberty. *Lancet*, 2, 955
162. Roger, M, Chaussain, JL, Berlier, P, Bost, M, Canlorbe, P, Colle, M, Francois, R, Garandeau, P, Lahlou, N, Morel, Y and Schally, AV (1986). Long term treatment of male and female precocious puberty by periodic administration of a long acting preparation of D-Trp-6-LH-RH microcapsules. *J Clin Endocrinol Metab*, 62, 670
163. Schally, AV, Redding, TW, and Comaru-Schally, Am (1983). Inhibition of prostate tumors by agonistic and antagonistic analogs of LH-RH. *Prostate*, 4, 545
164. Vickery, BH, McRae, GI, Sanders, LH, Kent, JS and Nestor, JJ Jr (1984). In vitro assessment of long-acting formulations of LHRH analogs. In: Zatuchni, GI, Goldsmith, A, Shelton, JD and Sciarra, JJ (eds.). "Long-Acting Contraceptive Delivery Systems". p.180. (Philadelphia: Harper and Row)
165. Schally, AV, Paz-Bouza, JI, Schlosser, JV, Karashima, T, Debeljuk, L, Gandle, B and Sampson, M (1987). Protective effects of analogs of luteinizing hormone-releasing hormone against x-radiation-induced testicular damage in rats. *Proc Natl Acad Sci USA*, 84, 851
166. Lewis, RW, Dowling, KJ and Schally, AV (1985). D-Tryptophan-6 analog of luteinizing hormone-releasing hormone as

a protective agent against testicular damage caused by cyclophosphamide in baboons. Proc Natl Acad Sci USA, 82, 2975.
167. Karashima, T, Zalatnai, A and Schally AV (1988). Protective effects of analogs of luteinizing hormone-releasing hormone against chemotherapy-induced testicular damage in rats. Proc Natl Acad Sci, 85, 2329

CHEMISTRY OF GnRH ANALOGUES

Anand S. DUTTA

Pharmaceuticals Division, Imperial Chemical Industries plc, Mereside,
Alderley Park, Macclesfield, Cheshire, UK

INTRODUCTION

The mammalian gonadotrophin releasing hormone structure was first published by Schally et al., in 1971 from porcine hypothalamus [1] and soon afterwards by Guillemin et al., from ovine hypothalamus [2]. Due to the enormous clinical potential of this peptide a large number of agonist and antagonist analogues have been synthesised in the last seventeen years. The structure-activity relationships of all these analogues cannot be summarised in the limited space available here. The main purpose of this review is, therefore, to identify and discuss only those changes which have been incorporated into the agonist and antagonist analogues which are either in the market or in advanced stages of development. Detailed structure-activity studies of agonist and antagonist analogues, and their clinical and biological studies have been reviewed elsewhere [3-7].

Until 1980 GnRH variants from other species had not been isolated and sequenced. Only recently the structures of chicken, salmon and lamprey GnRH have been published and these are shown in Table 1 along with the structure of the porcine and ovine GnRH [8]. Additional GnRH-like peptides have also been found in alligator, skink and lizard. The alligator and skink GnRH were similar to the chicken GnRH but of the three GnRH sequences identified in lizard one was similar to salmon and the other two were novel [8].

Due to the absence of structural data on the other naturally occurring GnRH-like peptides before 1980 all of the early work on the agonist analogues was based on the original porcine/ovine sequence. Some of the new information is however presently being used in the design of antagonist analogues.

STRUCTURE-ACTIVITY RELATIONSHIPS OF THE AGONIST ANALOGUES

The agonist analogues of GnRH which are presently being marketed, or are in advanced stages of clinical trials, are listed in Table 2. Of the analogues listed in the table goserelin,

Table 1. Structures of the GnRH molecules isolated from various species.

Origin	Sequence									
	1	2	3	4	5	6	7	8	9	10
Porcine,	Glu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-NH ₂									
Ovine										
Chicken I	-----							Gln	-----	
Chicken II	-----					His	Trp-Tyr		-----	
Salmon	-----						Trp-Leu	-----		
Lamprey	-----		Trp	-----		Leu-Glu-Trp-Lys	-----			

Table 2. Structures of GnRH analogues undergoing further evaluation.

Agonist Structure	Name (Company)
[D-Ser(Bu ⁺) ⁶ , Azgly ¹⁰]-LHRH	'Zoladex' (ICI)
[D-Ser(Bu ⁺) ⁶ , des-Gly-NH ₂ ¹⁰]-LHRH(1-9)NHE+	Buserelin (Hoechst)
[D-Trp ⁶]-LHRH	Tryptorelin (Debiopharm) (Decapeptyl)
[des-Gly-NH ₂ ¹⁰]-LHRH(1-9)-NHE+	Fertirelin (Takeda)
[D-His(Bzl) ⁶ , des-Gly-NH ₂ ¹⁰]-LHRH(1-9)NHE+	Histreltin (Ortho)
[D-Leu ⁶ , des-Gly-NH ₂ ¹⁰]-LHRH(1-9)NHE+	Leuprolide (Abbott)
[D-Trp ⁶ , MeLeu ⁷ , des-Gly-NH ₂ ¹⁰]-LHRH(1-9)NHE+	Lutreltin (Wyeth)
[D-Nal(2) ⁶]-LHRH	Nafarelin (Syntex)

buserelin, tryptorelin and leuporelin are already being marketed for the treatment of prostate cancer and histrelin and nafarelin are in advanced stages of clinical trials. These six analogues have either one or two changes from the parent peptide. In tryptorelin and nafarelin only the Gly⁶ residue has been replaced by a D-amino-acid residue but in the remaining four the C-terminal glycine residue has also been modified. Goserelin has an azaglycine residue in position 10 but buserelin, histrelin and leuporelin have an ethylamide group in place of the C-terminal Gly-NH₂. The significances of these position 6 and 10 changes are summarized below.

Replacement of the C-terminal Gly-NH₂ by an ethylamide group

Elimination of the C-terminal glycine residue gave a nonapeptide with about 10% biological activity (in vitro LH release). Further reduction in the chain length from the C-terminus led to inactive compounds (<0.01% GnRH). When the Gly-NH₂ was replaced with various linear, branched or cyclic alkylamines, e.g. a C-terminal ethylamide, propylamide or isopropylamide, some of the resulting analogues, were 2-3 times more potent than GnRH in releasing LH in pituitary cell cultures [9,10]. Other analogues with C-terminal methyl, dimethyl or butylamide groups were somewhat less potent. These early results formed the basis of all the agonists containing a C-terminal ethylamide group, e.g. buserelin, histrelin and leuporelin.

Replacement of the C-terminal Gly-NH₂ by an α -aza-amino-acid

α -Aza-amino-acid replacements in peptides are likely to induce conformational changes which may lead to increased receptor affinity and at the same time may also stabilise the peptide bonds to enzymic degradation. Detailed conformational studies on α -aza-peptides have not yet been reported but the enzymic stability of model azapeptides to various enzymes has been investigated [11]. The aza-residue was found to stabilise the peptide bonds (formed by the amino and the carboxyl groups of these residues) to aminopeptidase, carboxypeptidase and thermolysin. The effects on stability were comparable to those of the D-amino-acid residues.

The analogues containing a single aza-residue, e.g. [Azgly¹⁰]-GnRH and [Azala¹⁰]-GnRH, were as potent as GnRH in inducing ovulation in androgen-sterilised, constant oestrous rats [12, 13]. In contrast, [Ala¹⁰]-GnRH and [Pro¹⁰]-GnRH had earlier been shown to be much less potent (<5% GnRH) [10]. The 25-fold difference between the activities of [Azala¹⁰]-GnRH and [Ala¹⁰]-GnRH probably reflects the advantages of the conformational change induced by an aza-alanine residue.

Replacement of Gly⁶ with various L and D-amino-acid residues

Substitution of another L-amino-acid residue in position 6, e.g. Ala, Ile, Val, Pro, or some D-amino-acid residues, e.g. D-Val,

D-Pro, resulted in much less potent analogues (0.1 to 32% GnRH). In contrast replacement with a number of other D-amino-acid residues, e.g. D-Ala, D-Leu, D-Phe, D-Trp, D-Nal(2), D-Arg, D-hArg(Et₂), D-Ser(Bu^t), D-Tmp, D-Dmb, led to analogues which were 30 to 200-fold more potent than GnRH [3,14-19]. Because of this enormous increase in potency the D-amino-acid residues in position 6 play an essential role in all of the agonists presently in development.

Replacements in position 6 and 10

Various combinations of substitutions in positions 6 and 10 have been attempted in order to improve the potency even further but this approach has not been universally successful. The C-terminal ethylamide analogues of [D-Ala⁶]-GnRH, [D-Leu⁶]-GnRH (Leuprolide; leuprORElin) and [D-Ser(Bu^t)⁶]-GnRH (buserelin) were more potent than the corresponding Gly-NH₂¹⁰ analogues, but [D-Nal(2)⁶]-GnRH and [D-Nal(2)⁶,N-MeLeu⁷]-GnRH were more potent than the corresponding ethylamide analogues. In the position 10 aza-amino-acid series of analogues [D-Phe⁶, Azgly¹⁰]-, [D-Tyr(me)⁶,Azgly¹⁰]-, [D-Ser(Bu^t)⁶, Azgly¹⁰]- (Zoladex; goserelin), [D-Nal(2)⁶,Azgly¹⁰]- and [D-Dmb⁶ Azgly¹⁰]-GnRH were extremely potent [20-22].

STRUCTURE-ACTIVITY RELATIONSHIPS OF GnRH ANTAGONIST ANALOGUES

First antagonists of GnRH, discovered within a year of the discovery of GnRH itself, were obtained either by eliminating the whole of the histidine residue or just its side-chain [4, 23, 24]. Although [des-His²]- and [Gly²]-GnRH were weak antagonists (active only in *in vitro* test systems) their discovery pointed to the key to antagonism, and led to the synthesis of a large number of position 2 modified analogues. The majority of these, e.g. Ser, Thr, Gln, Arg, Leu, Phe, Tyr, Trp, D-Arg, D-Leu and D-Val² analogues, were only weak agonists. Only [D-Phe²]- and [D-Trp²]-GnRH were more potent antagonists than [des-His²]-GnRH.

Since changes in positions 6 and 10 had resulted in significant enhancement in potency in the agonist series of analogues these changes were also made in the antagonist series of analogues along with position 2 changes. All the des-His² and D-Phe² analogues with D-amino-acid residues in position 6, e.g. [des-His²,D-Ala⁶]-, [des-His²,D-Lys⁶]-, [des-His², D-Orn⁶]-, [D-Phe²,D-Ala⁶]-, [D-Phe²,D-Phe⁶]-, and [D-Phe²,D-Trp⁶]-GnRH, were more potent antagonists of GnRH than the parent [des-His²]- or [D-Phe²]-GnRH [25,26].

The C-terminal ethylamide substitution was not successful in the antagonists. Most of the ethylamide analogues of position 2 or positions 2 and 6 modified compounds, e.g. [des-His², D-Lys⁶, des-Gly-NH₂¹⁰]-GnRH(1-9)NHet, [D-Phe²,D-Ala⁶, des-Gly-NH₂¹⁰]-GnRH(1-9)NHet, were much less potent than the corresponding Gly¹⁰ decapeptides in inhibiting ovulation in rats

Table 3 Amino-acid Substituents Leading to potent LHRH Antagonists

1	2	3	4	5	6	7	8	9	10
Glu	His	Trp	Sar	Tyr	Gly	Leu	Arg	Pro	Gly
$\text{Ac-N-CH}_2\text{-CO-}$ (Pro)	D-Phe (p-Cl) D-Phe (p-F) D-Phe (3,4-di-Cl)	D-Trp D-Nal D-Phe	D-Trp D-Nal D-Phe	CH_3 CH_2 CH NH-CH-CO- (Ile)	CH_2 CH NH-CH-CO- 	$(\text{CH}_2)_3\text{-CH}_3$ NH-CH-CO- CH_2 NH-CH-CO- 	$(\text{CH}_2)_4\text{-NH}_2$ NH-CH-CO- NH-CH-CO-CH_2 $(\text{CH}_2)_4$ NH-CH-CO-	Pro	Gly NH-NH-CO- NH-CH-CO-
Ac-NH-CH-CO- 	D-Trp D-Nal D-Phe	D-Trp D-Nal D-Phe	D-Trp D-Nal D-Phe	CH_3 CH_2 CH NH-CH-CO- (Ile)	CH_2 CH NH-CH-CO- 	$(\text{CH}_2)_3\text{-CH}_3$ NH-CH-CO- CH_2 NH-CH-CO- 	$(\text{CH}_2)_4\text{-NH}_2$ NH-CH-CO- NH-CH-CO-CH_2 $(\text{CH}_2)_4$ NH-CH-CO-	Pro	Gly NH-NH-CO- NH-CH-CO-
Ac-NH-CH-CO- 	D-Trp D-Nal D-Phe	D-Trp D-Nal D-Phe	D-Trp D-Nal D-Phe	CH_3 CH_2 CH NH-CH-CO- (Ile)	CH_2 CH NH-CH-CO- 	$(\text{CH}_2)_3\text{-CH}_3$ NH-CH-CO- CH_2 NH-CH-CO- 	$(\text{CH}_2)_4\text{-NH}_2$ NH-CH-CO- NH-CH-CO-CH_2 $(\text{CH}_2)_4$ NH-CH-CO-	Pro	Gly NH-NH-CO- NH-CH-CO-
Ac-NH-CH-CO- 	D-Trp D-Nal D-Phe	D-Trp D-Nal D-Phe	D-Trp D-Nal D-Phe	CH_3 CH_2 CH NH-CH-CO- (Ile)	CH_2 CH NH-CH-CO- 	$(\text{CH}_2)_3\text{-CH}_3$ NH-CH-CO- CH_2 NH-CH-CO- 	$(\text{CH}_2)_4\text{-NH}_2$ NH-CH-CO- NH-CH-CO-CH_2 $(\text{CH}_2)_4$ NH-CH-CO-	Pro	Gly NH-NH-CO- NH-CH-CO-

(All D)

[25,27]. In contrast to the results with the C-terminal ethylamide analogues the results with an azaglycinamide substitution in position 10 were more promising. [des-His², Azgly¹⁰]- and [des-His², D-Phe⁶, Azgly¹⁰]-GnRH blocked ovulation completely at a dose of 250 µg/rat and partially at 62.5 µg/rat. [D-Phe^{2,6}, Azgly¹⁰]-GnRH blocked ovulation completely at a dose of 15 µg/rat and was about 10-fold more potent than [D-Phe^{2,6}]-GnRH [20,28]. Similar enhancement in antagonist potency was also observed in a number of other analogues with multiple substitutions [4].

Further improvement in potency was then sought by modifications in various other positions. This led to potent GnRH antagonists which had 5-8 of the original amino-acids replaced by unnatural amino-acid residues. The preferred amino-acid residues in each position of GnRH (for high antagonists potency) are summarised in Table 3 and some of the more potent antagonists incorporating these changes are listed in Table 4 [29-34]. All of these antagonists inhibited ovulation in constant cycling rats at doses below 10 µg/kg.

TABLE 4: Potent antagonists of LHRH

LHRH	Glu	His	Trp	Ser	Tyr	Gly	Leu	Arg	Pro	Gly	NH ₂
	Cl	Cl									
	I	I									
ORG 30093	Ac-D-Phe	D-Phe	D-Trp			D-Phe				D-Ala	
	Cl	Cl									
	I	I									
ORG 30276	Ac-D-Phe	D-Phe	D-Trp			D-Arg				D-Ala	
		F									
		I									
ORF 18260	Ac-D-Nal	D-Phe	D-Trp			D-Arg					
		Cl					Et ₂				
		I					I				
RS 68439	Ac-D-Nal	D-Phe	D-Trp			D-hArg				D-Ala	
		Cl									
		I									
	Ac-D-Nal	D-Phe	D-3-Pal			D-Arg				D-Ala	
		Cl									
		I									
	Ac-D-Nal	D-Phe	D-3-Pal			D-Arg	Trp			D-Ala	
		Cl									
		I									
	Ac-D-Nal	D-Phe	D-3-Pal			Arg	D-3-Pal			D-Ala	

Table 5: Anti-ovulatory and histamine releasing activities of some potent GnRH antagonists

Compound	Anti-ovulatory activity (ED ₁₀₀ µg/rat)	In vitro histamine release (ED ₅₀ µg/ml)
[Ac-D-Nal(2) ¹ , D-Phe(p-Cl) ² , D-Trp ³ , D-Arg ⁶ , D-Ala ¹⁰]-LHRH	1.0	0.1
[Ac-D-Nal(2) ¹ , α-Me-D-Phe(p-Cl) ² , D-Trp ³ , Arg ⁵ , D-Tyr ⁶ , D-Ala ¹⁰]-LHRH	2.5	3.7
[Ac-D-Nal(2) ¹ , D-Phe(p-Cl) ² , D-3-Pal ³ , Arg ⁵ , D-Glu(AA) ⁶ , D-Ala ¹⁰]-LHRH	1.5	1.6
[Ac-D-Nal(2) ¹ , D-Phe(p-Cl) ² , D-3-Pal ³ , Arg ⁵ , D-3-Pal ⁶ , D-Ala ¹⁰]-LHRH	1.0	2.9
[Ac-D-Nal(2) ¹ , D-Phe ^{2,3} , D-Lys(x) ⁶ , Phe ⁷ , D-Ala ¹⁰]-LHRH		
X = H	3.0	0.37
- cyclopentyl	3.0	0.24
- cyclohexyl	3.0	0.32
[Ac-D-Nal(2) ¹ , D-Phe ^{2,3} , D-Arg ⁶ , Phe ⁷ , Lys(x) ⁸ , D-Ala ¹⁰]-LHRH		
X = H	>3	0.38
- neopentyl	>3	4.68
[Ac-D-Nal(2) ¹ , D-Phe(p-Cl) ² , D-3-Pal ³ , (5), (6), (8), D-Ala ¹⁰]-LHRH		
(5) (6) (8)		
Nic Nic IPr		
I I I	1.0	>300
Lys D-Lys Lys		
Nic Nic IPr		
I I I	1.0	206
Lys D-Lys Orn		
Pic Pic IPr		
I I I	0.5	93
Lys D-Lys Lys		

An examination of the types of amino-acid residues needed for high antagonist potency and also the structures of the potent antagonists shows that the overall hydrophobicity and basicity of the original GnRH sequence are significantly increased. This combination of increased hydrophobicity and basicity has led to a problem due to histamine release induced by most of the potent antagonists. This problem has previously been associated with a number of other peptides, e.g. substance P, which contain hydrophobic and basic residues [35]. Attempts have recently been made to synthesise analogues lacking in histamine releasing activity and these can broadly be categorised into three types. The first approach involved switching the residues between positions 5 and 6. The second and third approaches were based on reducing the overall hydrophobicity and shielding the side-chain basic groups. These were achieved by replacing the D-Trp³ with D-Pal³ and also by incorporating various substituents, e.g. isopropyl, nicotinyl or picolyl, on the side-chain amino groups in positions 5, 6 and 8. Examples of some of these modified analogues are shown in table 5 along with their antiovarulatory and in vitro histamine releasing activities [36-38]. In comparison to the parent peptide, [Ac-D-Nal(2)¹,D-pCl-Phe²,D-Trp³,D-Arg⁶,D-Ala¹⁰]-GnRH, the analogues containing an Arg residue in position 5 instead of a D-Arg residue in position 6 were 16-37 times less potent in releasing histamine. The Arg⁵ analogues were only marginally less potent as antiovarulatory agents. Incorporation of cyclopentyl, cyclohexyl and some other groups onto the side-chain amino group of Lys⁶ did not show any improvement in the histamine releasing activity. An analogue with a neopentyl group in position 8, [Ac-D-Nal(2)¹,D-Phe^{2,3},D-Arg⁶,Phe⁷,Lys(neopentyl)⁸,D-Ala¹⁰]-GnRH, was about 10-15 times less potent in releasing histamine than the corresponding Lys⁸ analogue but both of these analogues were also less potent as antiovarulatory agents (ED₁₀₀ >3 µg/rat). Analogues with side-chain modifications in positions 5, 6 and 8 gave the most promising results. A number of such analogues, e.g. [Ac-D-Nal(2)¹,D-p-Cl-Phe²,D-Pal(3)³,Lys(Nic)⁵,D-Lys(Nic)⁶,Lys(iPr)⁸,D-Ala¹⁰]-, [Ac-D-Nal(2)¹,D-p-Cl-Phe²,D-Pal(3)³,Lys(Nic)⁵,D-Lys(Nic)⁶,Orn(iPr)⁸,D-Ala¹⁰]- and [Ac-D-Nal(2)¹,D-p-Cl-Phe²,D-Pal(3)³,Lys(Pic)⁵,D-Lys(Pic)⁶,Lys(iPr)⁸,D-Ala¹⁰]-GnRH, were very poor in releasing histamine from the rat mast cells (ED₅₀ 90-300 µg/ml) but were extremely potent as antiovarulatory agents (ED₅₀ 0.51 µg/rat). In vivo side-effect profiles of these analogues have not yet been reported.

CONCLUSIONS

The chemical work on GnRH has led to potent agonist and antagonist analogues which have been shown to be effective in various fertility related disorders, contraception and hormone-dependent tumors. Some agonists have been formulated to give slow release of the peptide and some of these slow release formulations (including Zoladex depot) are already on the market.

The progress in the antagonist field has been slow. Considerable work still remains to be done in improving the side-effect profile and developing slow release formulations before these agents can be considered as useful drugs.

REFERENCES

1. Matsuo, H, Baba, Y, Nair, RM, Arimura, A and Schally, AV (1971). Structure of porcine LH and FSH releasing hormone I. The proposed amino-acid sequence. *Biochem Biophys Res Commun*, 43, 1334
2. Burgus, R, Butcher, M, Amoss, M, Ling, N, Monahan, M, Rivier, J, Fellows, R, Blackwell, R, Vale, W, and Guillemin, R (1972). Primary structure of the ovine hypothalamic luteinising hormone-releasing factor (LRF). *Proc Natl Acad Sci USA*, 69, 278
3. Dutta, AS (1988) Luteinizing hormone-releasing hormone (LHRH) agonists. *Drugs of the Future*, 13, 43
4. Dutta, AS (1988). Luteinising hormone-releasing hormone (LHRH) antagonists. *Drugs of the Future*, 13, in press
5. Dutta, AS and Furr, BJA (1985). Luteinizing hormone releasing hormone (LHRH) analogues. *Annu Rep Med Chem*, 20, 203
6. Bex, FJ and Corbin, A (1984). LHRH and analogues: reproductive pharmacology and contraceptive and therapeutic utility. In: Martin L and Ganong WF (eds) "Frontiers in neuro-endocrinology". p. 85. (New York: Raven Press)
7. Karten, MJ and Rivier, JE (1986). GnRH analogue design. Structure-function studies towards the development of antagonists and antagonists: rationale and perspective. *Endocr Rev*, 7, 44
8. Millar, RP and King JA (1987). Structural and functional evolution of gonadotropin-releasing hormone. *Int Rev Cytol*, 106, 149
9. Fujino, M, Kobayashi, S, Obayashi, M, Fukuda, T, Shinagawa, S, Yamazaki, I, Nakayama, R, White, WF and Rippel, RH (1972). Syntheses and biological activities of LHRH. *Biochem Biophys Res Commun*, 49, 698
10. Fujino, M, Shinagawa, S, Obayashi, M, Kobayashi, S, Fukuda, T, Yamazaki, I, Nakayama, R, White, WF and Rippell, RH (1973). Further studies on the structure-activity relationships in the C-terminal part of LHRH. *J Med Chem*, 16, 1144
11. Dutta, AS and Giles, MB (1976). Polypeptides. Part XIV. a comparative study of the stability towards enzymes and model tripeptides containing α -aza-amino-acids, L-amino-acids and D-amino-acids. *J Chem Soc Perkin*, I, 244
12. Dutta, AS, Furr, BJA, Giles, MB and Morley (1976). Synthesis and biological activity of α -azapeptides: α -aza-analogues of LHRH. *Clin Endocrinol*, 5(Suppl), 291S
13. Dutta, AS, Furr, BJA, and Giles, MB (1979). Polypeptides. Part 15. Synthesis and biological activity of α -aza-analogues of luliberin modified in positions 6 and 10. *J Chem Soc Perkin*, I, 379
14. Monahan, MW, Amoss, MS, Anderson HA and Vale W (1973). Synthetic analogues of LHRH with increased agonist or antagonist properties. *Biochem*, 12, 4646

15. König, W, Sandow, J and Geiger, R (1975). Structure-function relationships of LHRH/FSH-RH. In: Walter R and Meinenhofer J (eds). "Peptides: Chemistry, Structure and Biology". p. 888. (Ann Arbor: Science Publishers Inc)
16. Nestor JJ Jr, Ho, TL, Simpson, RA, Horner, BL, Jones, GH, McRae, GI and Vickery, BH (1982). Synthesis and biological activity of some very hydrophobic superagonist analogues of LHRH. *J Med Chem*, 25, 795
17. Nestor, JJ Jr, Horner, BL, Ho, TL, Jones, GH, McRae, GI and Vickery, BH (1984). Synthesis of a novel class of heteroaromatic amino-acids and their use in the preparation of analogues of LHRH. *J Med Chem*, 27, 320
18. Coy, DH, Labrie, F, Savary, M, Coy, EJ and Schally AV (1975). LH-releasing activity of potent LHRH analogues in vitro. *Biochem Biophys Res Commun*, 67, 576
19. Nestor, JJ Jr, Tahiramani, R, Ho, TL, McRae, GI and Vickery BH (1985). Potent LHRH agonists containing NG, NG'-dialkyl-D-homoarginines. In: Deber, CM, Hruby VJ and Kopple KD (eds). "Peptides: Structure and Function". p. 557. (Illinois: Pierce Chemical Company)
20. Dutta, AS, Furr, BJA, Giles, MB, Valcaccia, B and Walpole, AL (1978). Potent agonist and antagonist analogues of luliberin containing an azaglycine residue in position 10. *Biochem Biophys Res Commun*, 81, 382
21. Dutta, AS, Furr, BJA, Giles, MB and Valcaccia, B (1978). Synthesis and biological activity of highly active α -aza-analogues of luliberin. *J Med Chem*, 21, 1018
22. Ho, TL, Nestor, JJ Jr, McRae, GI and Vickery, BH (1984). Hydrophobic azaglycine analogues of LHRH. *Int J Peptide Protein Res*, 24, 79
23. Vale, W, Grant, G, Rivier, J, Monahan, M, Amoss, M, Blackwell, R, Burgus, R and Guillemin, R (1972). Synthetic polypeptide antagonists of the hypothalamic luteinizing hormone releasing hormone releasing factor. *Science*, 176, 933
24. Folkers, K, Humphries, J and Bowers, CY (1982). Evolution of design and achievement of inhibitors of the luteinising hormone-releasing hormone as inhibitors of ovulation. *Z. Naturforsch*, 37b, 246
25. Ling, NC, Rivier, JE, Monahan, MW and Vale, WW (1976). Analogues of LHRH modified at positions 2, 6 and 10. *J Med Chem*, 19, 937
26. Rivier, JE and Vale, WW (1978). [D-Glu¹, D-Phe², D-Trp^{3,6}]-LRF. A potent luteinising hormone releasing factor antagonist in vitro and inhibitor of ovulation in the rat. *Life Sci*, 23, 869
27. Yardley, JP, Foell, TJ, Beattie, CW and Grant, NH (1975). Antagonism of luteinising hormone release and ovulation by an analogue of LHRH. *J Med Chem*, 18, 1244
28. Dutta, AS, Furr, BJA and Giles MB (1979). Part 16. Synthesis and biological activity of α -aza-analogues of luliberin with high antagonist activity. *J Chem Soc Perkin*, I, 389

29. Coy, DH, Horvath, A, Nekola, MV, Coy, EJ, Erchegeyi, J and Schally, AV (1982). Peptide antagonists of LHRH: large increases in antioovulatory activities produced by basic D-amino-acids in six position. *Endocrinology*, 110, 1445
30. Nestor, JJ Jr, Tahilramani, R, Ho, TL, McRae, GI, Vickery, BH and Bremner, WJ (1983). New luteinizing hormone-releasing factor antagonists. In: Hruby VJ and Rich DH (eds). "Peptides, Structure and Function". p. 861. (Illinois; Pierce Chemical Company)
31. Erchegeyi, J, Coy, DH, Nekola MV, Coy, EJ, Schally, AV, Mezo, I and Teplan, I (1981). LHRH analogues with increased antioovulatory activity. *Biochem Biophys Res Commun*, 100, 915
32. Folkers, K, Bowers, CY, Kubiak, T and Stepinski, J (1983). Antagonists of LHRH with pyridylalanines which completely inhibit ovulation at nanogram dosage. *Biochem Biophys Res Commun*, 111, 1089
33. Folkers, K, Bowers, CY, Shieh, H-M, Yin-Zeng, L, Shao-bo, X, Tang, P-F, L and Ji-Yu, C (1984). Antagonists of LHRH with emphasis on the Trp⁷ of the salmon and chicken II LHRH's. *Biochem Biophys Res Commun*, 123, 1221
34. Folkers, K, Bowers, C, Shao-bo, X, Tang, P-F, L, and Kubota, M (1986). Increased potency of antagonists of LHRH which have D-3-Pal in position 6. *Biochem Biophys Res Commun*, 137, 709
35. Dutta, AS (1987). Agonists and antagonists of substance P. *Drugs of the Future*, 12, 781
36. Karten, MJ, Hook, WA, Siraganian, RP, Coy, DH, Folkers, K, Rivier, JE and Roeske, RW (1987). In vitro histamine release with LHRH analogs. In: Vickery, BH and Nestor, JJ, Jr (eds). "LHRH and Its Analogs. Contraceptive and Therapeutic Applications. Part 2". p. 179. (Lancaster; MTP Press Ltd)
37. Hocart, SJ, Nekola, MV and Coy, DH (1987). Effect of reductive alkylation of lysine in position 6 and/or 8 on the histamine-releasing activity of LHRH antagonists. *J Med Chem*, 30, 1910
38. Ljungqvist, A, Feng, D-M, Tang, PL, Kubota, M, Okamoto, T, Zhang, Y, Bowers, CY, Hook, WA and Folkers, K (1987). Design, synthesis and bioassay of antagonists of LHRH which have high antioovulatory activity and release negligible histamine. *Biochem Biophys Res Commun*, 148, 849

ACTIVE REDUCED SIZE ANALOGUES

**Fortuna HAVIV, Christopher A. PALABRICA, Eugene N. BUSH,
Gilbert DIAZ, Edwin S. JOHNSON and Jonathan GREER**

Pharmaceutical Discovery Division, Abbott Laboratories, Abbott Park,
Illinois 60064, USA

INTRODUCTION

Although a large number of GnRH analogues have been synthesized, little has been reported on reduction of the size of these peptides. In addition, there are very few reports in the literature about active fragments of GnRH. J. Sandow and co-workers [1] tested fragments of buserelin using ovulation induction in the rat. They examined fragments wherein amino acids were removed sequentially from the N-terminus and they found that the (3-9)-buserelin was the sole fragment which still retained significant activity. It had 23% of GnRH activity. Although this activity represents only 0.2% of buserelin potency, we chose the (3-9)-peptide as a starting point for the design of reduced size GnRH analogues.

METHODS

Chemistry

All of the peptides were synthesized using solid phase methodology. All of the products were purified using HPLC and characterized using fast atom bombardment (FAB) mass spectroscopy, and amino acid analysis.

Biology

Receptor binding

A modified procedure from the literature [2] was utilized for the receptor binding assay. A rat pituitary plasma membrane fraction is prepared as a source of GnRH receptors, and ^{125}I -(Tyr⁵)-leuprolide is made via chloramine-T oxidation and purified via ion exchange chromatography. Several concentrations of test compounds are coincubated for 120 minutes at 4°C with ^{125}I -(Tyr⁵)-leuprolide and receptors until equilibrium binding, then bound tracer is separated from free via centrifugation. Data are reported as pK_I , the negative logarithm of the equilibrium

dissociation constant.

LH release

The ability of compounds to release luteinizing hormone (LH) from cultured pituitary cells [3, 4] was utilized for establishment of their activities. Pituitaries are dispersed with collagenase and hyaluronidase, and the cells are plated in culture plates for 72 hours. The cells are incubated with test compounds for 3 hours at 37°C, then the medium is harvested for determination of released LH via radioimmunoassay (RIA). The RIA utilizes ^{125}I -LH (ovine) as a tracer, immunoaffinity purified anti-rat-LH IgG from Dr. P.M. Conn, and rat LH (NIH-RP2) standard. Bound and free tracer are separated with immobilized protein A. The results for agonists are reported as a pD_2 which is defined as the negative logarithm of the concentration of agonist which results in 50% of the maximum release of LH produced by the compound. Antagonist results are reported as a pA_2 , the negative logarithm of the concentration of an antagonist which shifts the LH release dose-response produced by leuprolide two-fold to the right.

RESULTS AND DISCUSSION

As an initial screen we selected rat pituitary receptor binding [2] (pK_1) and rat pituitary cell culture LH release [3, 4] (pD_2 for agonist, pA_2 for antagonist). In these assays, GnRH had a pK_1 of 8.90 and a pD_2 of 9.23 (Table 1).

Table 1. Rat pituitary receptor binding and LH release or inhibition by (3-9) GnRH analogues and indole-(4-9) GnRH analogues. Modifications of position 6.

X ³ -Ser-Tyr-Z ⁶ -Leu-Arg-ProNHet					
No.	X ³	Z ⁶	pK ₁	pD ₂	pA ₂
1	Trp	DLeu	7.07		6.06
2	Trp	DTrp	8.33		7.24
3	(1)-Nal ^a	DLeu	7.30		
4	none	DLeu	5.36	NT ^b	NT
5	3-Indolepropionyl	DLeu	6.71	6.93	
6	3-Indolepropionyl	DTrp	8.34	7.35	
7	3-Indolepropionyl	D-(2)Nal ^a		8.77	
8	3-Indolepropionyl	DPhe	7.78		6.07
9	3-Indolepropionyl	DCha ^a	7.38		6.44
GnRH			8.90	9.27	
D-4-Cl-Phe ^{1,2} DTrp ³ DArg ⁶ DAla ¹⁰			10.74		9.44


^a(1)-Nal = 1-naphthyl-3-alanine; D-(2)-Nal = D-2-naphthyl-3-alanine; D-Cha = D-3-cyclohexylalanine. ^bNT = not tested.

(3-9)-Leuprolide (1) had a pK_I of 7.07 and did not release LH, but antagonized its release with a pA_2 of 6.06. Similarly, D-Trp⁶-(3-9)-leuprolide (2) was also a potent antagonist with pA_2 of 7.24. These results were surprising to us since (3-9)-buserelin was reported to be an agonist, and was active in the ovulation induction test in the rat [1]. For the (1)-Nal³-(3-9)-leuprolide, which had a pK_I of 7.30, neither pD_2 or pA_2 could be determined suggesting that the peptide was probably degraded either chemically or by some proteolytic enzymes in the pituitary cell culture medium. (4-9)-Leuprolide (4) had a very weak binding with pK_I of 5.36 and was not tested further.

The first attempted structural modification was replacement of the Trp³ of (1) with desamino-Trp or 3-indolepropionic acid to give compound 5 with virtually the same receptor binding affinity, indicating that the amino group is not essential for receptor binding. Removal of the α -amino group converts the molecule from antagonist to agonist. Affinity and potency increase as the residue at position 6 is changed from D-Leu to D-Trp and D-(2)-Nal; however, while the D-Leu (5) and D-Trp (6) forms are agonists, the D-(2)-Nal (7), as well as D-Phe (8) and D-Cha (9) are antagonists. The results indicate that not only the substituent at position 3 can influence the activity to be either agonist or antagonist, but also the substituent at position 6 has a similar effect. The highest affinity molecules of this series are about one fifth (for agonist 6) to equal (for antagonist 7) the affinity of GnRH. The affinity of antagonist 7 was one hundredth that of [D-4-Cl-Phe^{1,2}, D-Trp³, D-Arg⁶, D-Ala¹⁰] LHRH, a potent, commercially available, decapeptide antagonist.

With affinity levels as high as these in a molecule that contains only six amino acids, we decided to explore the structure activity relationships a little further. The length of the aliphatic chain, connecting the indole group at position 4 to the molecule was varied (Table 2). Shortening the chain in (5) by one

Table 2. Rat pituitary receptor binding and LH release or inhibition by indole-(4-9) GnRH analogues. Variations of the indole chain length.



No.	n	Position 3	Z ⁶	pK_I	pD_2	pA_2
10	0	3-Indoloyl	DLeu	6.69		6.72
11	1	3-Indoleacetyl	DLeu	5.87		5.56
12	2	3-Indolepropionyl	DLeu	6.71	6.93	
13	3	3-Indolebutyryl	DLeu	6.91		6.72
14	3	3-Indolebutyryl	D-(2)Nal ^a	8.86		8.07

^aD-(2)-Nal = D-2-naphthyl-3-alanine.

methylene (11) converted the compound from agonist to antagonist, however with a large loss in binding. Further shortening of the chain by an additional methylene (10) was still an antagonist, but receptor binding affinity was restored. Lengthening the chain in (5) by one methylene (12) also converted the compound from agonist to antagonist, but in this case maintaining the same binding affinity. [3-Indolebutyryl³-D-(2)-Nal⁶-Pro⁹-NHET]-(4-9)-LHRH (14) showed the highest binding affinity, which was equivalent to GnRH. Thus, once again we were able to shift between agonist and antagonist, this time solely by changing the number of methylene groups in the chain at position 3. The results suggest that there is an optimal size and shape for substituents at position 3 for potency and for agonist versus antagonist activity. These are unprecedented results, since previously antagonists have been reported by replacing residues 2, 3, and sometimes 1, and 10 with D-amino acids, or by deleting residue 2 in the decapeptide or nonapeptide GnRH analogue.

To further analyze the properties of the substituents at position 3, 1-naphthylpropionic acid, corresponding to the (1)-Nal³ substitution, was placed at this position (Table 3).

Table 3. Rat pituitary receptor binding and LH release or inhibition by naphthyl (4-9) GnRH analogues. Modification of position 6 and variations of the naphthyl chain length.

X ³ -Ser-Tyr-Z ⁶ -Leu-Arg-ProNHet					
No.	X ³	Z ⁶	pK ₁	pD ₂	pA ₂
14	1-Naphthylpropionyl	DLeu	7.60		6.60
15	1-Naphthylpropionyl	DTrp	9.28		8.74
16	1-Naphthylpropionyl	D-(2)NaI ^a	9.55		9.25
17	1-Naphthylpropionyl	DPhe	8.01		7.83
18	1-Naphthylpropionyl	DCha ^a	7.44		6.08
19	1-Naphthoyl	DLeu	6.12	6.00	
20	1-Naphthylacetyl	DLeu	7.17	6.85	
21	1-Naphthylacetyl	DTrp	8.65		8.18
22	1-Naphthylacetyl	D-(2)NaI	8.58		8.20
23	1-Naphthylacetyl	DPhe	8.01		7.83
24	1-Naphthylacetyl	DCha	7.44		6.08

^aD-(2)-Nal = D-2-naphthyl-3-alanine; D-Cha = D-3-cyclohexylalanine.

Compounds with impressive receptor binding were obtained; the antagonist [N-(1-naphthylpropionyl),Ser⁴,D-2-Nal⁶,ProNHET]-(4-9)-LHRH (16) had fifteen times higher affinity than GnRH and virtually the same antagonist potency (pA₂) as the potent standard antagonist, [D-4-Cl-Phe^{1,2},D-Trp³,D-Arg⁶,D-Ala¹⁰]LHRH, even though it contained only six amino acids (versus 10 in the reference antagonist) and only one of D configuration (versus five in the reference antagonist). When the

length of the aliphatic chain connecting the naphthyl group to the molecule was shortened by one or two methylenes (20 and 19 respectively) once again the biological response switched, this time from antagonist to agonist. Consequently, we examined the influence of changes at position 6 on the biological activity of N-naphthylacetyl-Ser⁴series. Again, as the size of the residue at position 6 was increased from D-Leu to D-Cha, D-Phe, D-Trp, or D-(2)-Nal, the molecule changed from an agonist to an antagonist with the expected enhancement in binding.

Table 4. Rat pituitary receptor binding and LH release or inhibition by naphthyl (4-9) GnRH analogues. Modification of position 3.

X³-Ser-Tyr-Z⁶-Leu-Arg-ProNHET

No.	X ³	Z ⁶	pK ₁	pD ₂	pA ₂
25	2-Naphthylacetyl	DTrp	8.43		7.80
26	1-Naphthylacrylyl	DLeu	7.30		6.69
27	1-Naphthoxyacetyl	DTrp	8.72		7.89
28	trans-O-Methoxycinnamoyl	DTrp	6.80		6.46
29	cis-O-Methoxycinnamoyl	DTrp	8.85		7.22
30	3-Pyridylpropionyl	DLeu	5.79	4.96	
31	3-Pyridylacrylyl	DLeu	5.82		5.72
32	1-Adamantylacetyl	DTrp	9.34		8.50

To further probe the properties of the side chain at position 3, we decided to examine several more radical modifications for Trp³ in the (3-9)-GnRH series. Table 4 shows that a wide range of affinities were obtained, from very weak for the 3-pyridylpropionyl (30) to quite high affinity for the adamantylacetyl (32) with binding affinity three-fold higher than GnRH.

Finally, as a byproduct of the solid phase synthesis, several analogues were obtained with an O-benzyl group on the Ser⁴ side chain (Table 5). The Ser⁴(O-Bzl) peptides had only slightly different binding affinity from their respective parents, sometimes greater, sometimes lower, indicating that the free hydroxyl of the Ser⁴ is not essential for binding. The biological activity results were more anomalous. In the case of 1-naphthylpropionyl³-D-Leu⁶ (33), the molecule was transformed from an antagonist to an agonist by the O-benzyl substitution. In another case, 1-naphthylacetyl³-D-Leu⁶, (35), the reverse effect happened; the O-benzyl group changed an agonist (20) into an antagonist (35).

measurement of LH by radioimmunoassay. Immediately thereafter ice cold methanol / 6 N HCl (1:200) solution was added (2 ml per well), and transferred to tubes with chloroform (1 ml and water (1.8 ml). Cells were scraped off using a rubber policeman. The extracts were sonified (for 5 min) and centrifuged (3000xg for 10 min). The upper water soluble phase containing IP was separated from the lower phase containing the lipid fraction (polyphosphoinositides). The radiolabeled inositol phosphates in the aqueous phase were determined by anion exchange chromatography on columns containing AG1-x8 resins (BioRad, 200-400 mesh, formate form). Free inositol was eluted in the flow-through of distilled water and glycerophosphate by disodium tetraborate 5 mM/sodium formate 60 mM. Then sequential washes with 0.1 M formic acid containing 0.2, 0.4 and 1.0 M ammonium formate progressively eluted IP₁, IP₂ and IP₃. Five ml fractions were collected and measured after addition of 10ml scintillation fluid (Amersham) for myo-(2-³H)-inositol.

Standard curves have been established using radiolabeled inositophosphates. Data are shown in counts per minute myo-(2-³H)-inositol \pm SD. Each data point consists of 3 independent control values and shows one out of two experiments. Statistical analysis was performed by Student's t test and, when for more than two means, analysis of variance was carried out.

Superfusion of cultured cells with phospholipases, arachidonic acid and leukotrienes

In contrast to static cultures, superfused cells are good for investigating substances with a short half-life or for determining the secretory profile of stimulants [21-24]. For that purpose, pituitary cells were cultured after enzymatic dispersion on cytodex beads and transferred to columns. They were allowed to recover for 90 min while being superfused with medium 199 Earle and 20 mM HEPES at a speed of 9.5 ml/min. Pituitary cells were stimulated by the addition of GnRH for 4 min dissolved in medium 199. As shown in Fig. 1, pulses of GnRH evoked a rapid release of LH with increasing amount of LH released per peak (area under the curve). GnRH was also added as a pulse at the end of other superfusions to demonstrate response of cells, and it served as a comparison for test substances. Arachidonic acid, and its lipoxygenase metabolites 5-hydroxy-6,8,11,14-eicosatetraenoic acid (5-HETE), (5s)-5-hydroperoxy-6,8,11,14-eicosatetraenoic acid (5-HPETE), 15-hydroxy-5,8,10,14-eicosatetraenoic acid (15-HETE), leukotrienes A₄, B₄, C₄, D₄ and E₄ were dissolved in ethanol after deacylation and further diluted in incubation medium prior to use. These agents were added to media for 4 min with a final concentration of 0.01% ethanol. Phospholipase A₂, melittin, quinacrine and chloroquine were added in incubation medium. No effect of 0.01% ethanol on superfused rat pituitary cells was detected. Fractions were collected every 2 min and assayed for LH by radioimmunoassay kits RP-2 provided by the Hormone and Pituitary Program, Baltimore, USA. Data are presented

results obtained that a wide range of receptor binding affinities can be achieved with these hexapeptide GnRH analogues up to and exceeding the activity of GnRH itself. Even more important, by manipulating the group at position 3, one can change from agonist to antagonist almost at will. In examining the structure-activity relationships, it appears that a very precise size and shape of side chain substituent is required at position 3 to achieve agonist activity. If the group is too large, too small, too long, or too short, the molecule becomes an antagonist. The receptor binding site for position 3 seems to be formed in such a way that even though the response switches between agonist and antagonist quite readily depending upon changes in the size and shape of the substituent that occupies the site, these changes may have little effect upon the receptor binding affinity.

A novel result has been obtained upon modifying the nature of the side chain at position 6 in order to improve potency. While still performing its usual function of modulating receptor binding affinity in much the same way that it does in the nonapeptide or decapeptide GnRH analogues, i.e. D-(2)-Nal > D-Trp > D-Leu, in the hexapeptide series, position 6 somehow feeds back to the residue at position 3 to change the compound from agonist to antagonist as the side chain gets larger.

Finally, even modification of the serine side chain at position 4 by O-substitution with a benzyl group has been observed to trigger the switch from agonist to antagonist in either direction depending on the nature of the substituent at position 3. This suggests that it is not a direct effect of the benzyl group, but once again the feeding back of a conformational change, probably small, to the 3 position which influences the agonist to antagonist transition.

ACKNOWLEDGEMENT

We are indebted to Dr. P. Michael Conn of the University of Iowa for providing the anti-rat LH antibody and for his helpful suggestions regarding biological methodology.

REFERENCES

1. Sandow, J and Konig, W (1979). Studies with fragments of a highly active analogue of luteinizing hormone releasing hormone. *J Endocrinol*, 81, 175
2. Marian, J, Cooper, RL and Conn, PM (1981). Regulation of the rat pituitary gonadotrophin-releasing hormone receptor. *Mol Pharmacol*, 19, 399
3. Jinnah, HA and Conn, PH (1985). GnRH stimulated LH release from rat anterior pituitary cells in culture- refractoriness and recovery. *Amer J Physiol*, 249, E619
4. Vale, W, Grant, G, Amoss, M, Blackwell, R and Guillemin, R (1974). Culture of enzymatically dispersed anterior pituitary cells: functional validation of a method. *Endocrinology*, 91, 562

INTRACELLULAR ASPECTS

Ludwig KIESEL and Benno RUNNEBAUM

Division of Gynecological Endocrinology, Department of Obstetrics
and Gynecology, University of Heidelberg, Voss-Strasse 9,
D-6900 Heidelberg, FRG

INTRODUCTION

The physiological actions and clinical applications of GnRH and its analogs have been extensively studied [1], but the mechanisms of GnRH and its regulation are not fully clarified yet. GnRH stimulates the biosynthesis and secretion of LH and FSH from the anterior pituitary gland. The intracellular pathway through which GnRH induces the release of pituitary gonadotropins is receptor-mediated and calcium-dependent. Cyclic nucleotides including cAMP [2,3] and later cGMP [4,5] were proposed to act as second messengers during GnRH-stimulated LH release. However, cAMP production does not correlate with GnRH-stimulated LH release and GnRH effects on LH release are not reproduced by elevation of intracellular cAMP concentrations [6]. Furthermore, while cGMP formation correlates well with the activation of LH secretion by GnRH and is also calcium-dependent, LH release has been shown to occur independently of cGMP production in GnRH-stimulated gonadotropes [7]. It is now believed that cyclic nucleotides play little if any part in the acute action of GnRH upon gonadotropin release [7,8], and that the effects of cAMP upon gonadotrope function are of a more long-term nature that may contribute to the facilitation of LH release during subsequent stimulations [9].

The stimulation of gonadotropin release by GnRH and high potassium concentrations is dependent upon the presence of extracellular calcium [10,11], raising the possibility of calcium influx being an important component of the activation process. Several recent studies have confirmed that increases in cytosolic calcium stimulate LH release [12-14], and that the actions of GnRH upon LH secretion and cGMP formation are related to the extracellular calcium concentrations [13]. The role of Ca^{2+} in GnRH action is well established. LH release is fully dependent on extracellular Ca^{2+} [14], but the binding of GnRH to its receptor is not a Ca^{2+} -dependent step [15]. The elevation of intracellular Ca^{2+} by different methods i.e. calcium ionophores, Ca^{2+} -loaded liposomes, veratridine provoked release of LH [16] and, in addition, the use of the fluorescent analog of the Ca^{2+} -specific

chelator EGTA allowed the demonstration of intracellular movement of Ca^{2+} in response to GnRH [17].

The observation that drugs which inhibit calmodulin also inhibit GnRH-stimulated gonadotropin secretion provides evidence that calmodulin may serve as an intracellular receptor for calcium following gonadotrope activation by GnRH [18]. In many tissues that are activated by extrinsic ligands changes of phospholipid metabolism have been observed which influence the availability and action of Ca^{2+} .

Various post receptor systems that may be linked to this mechanism (i.e. phospholipids and inositol phosphates, arachidonic and metabolites, protein kinase C) have been analyzed in regard to their function to mediate or regulate GnRH action on gonadotropin secretion.

METHODS FOR STUDYING POST-RECEPTOR SYSTEMS

Preparation of pituitary cells

Anterior pituitary glands of adult female Sprague-Dawley rats (200-250 g) were dispersed into a single cell suspension by trypsinization. Whole pituitaries from adult female rats were cut in small blocks and suspended in medium 199 containing 0.3% bovine serum albumin (BSA). The medium was replaced by 5 ml of 0.5% trypsin (type III, Sigma, Munich, FRG) in medium 199-0.3% BSA and incubated at 37°C for 15 min. Subsequently the medium was decanted and incubations with 5 ml DNase and soybean trypsin inhibitor (Sigma) were carried out for 1 min and 5 min, respectively. The tissue has been further incubated for 15 min each with 2 mM and 1 mM EDTA in Ca^{2+} and Mg^{2+} -free medium 199. Thereafter fragments were gently dispersed to a single cell suspension using a plastic Pasteur pipette. The cells were cultured for 2 days (3×10^5 cells/well) in medium 199 (Bio-Products, Berlin, FRG) containing 10% horse serum and 100 units penicillin and 100 μg streptomycin/ml. For certain experimental procedures pituitary cells were enriched for gonadotropes using centrifugal elutriation [19].

Measurement of inositol phosphate accumulation

GnRH-induced inositol phosphate (IP) production was measured to investigate the role of IP in the mechanism of gonadotropin secretion [20]. Two million cells per well were incubated for 4 days in Medium 199 Earle (modified, Seromed, Berlin/FRG) with penicillin 100 U/ml, streptomycin 100 mg/100 ml, 10% horse serum. On day 2 of incubation, 12×10^6 cpm myo-(2- ^3H)-inositol (10-20 Ci/mmol; Amersham) was added to each well. On day 4 medium was removed and replaced by stimulation medium (Medium 199 Earle, 20 mM HEPES, 0.01% BSA). Ten minutes prior to stimulation LiCl_2 was added, to prevent IP breakdown. Then GnRH was added at a final concentration of 10^{-7} M. The reaction was stopped by placing wells on dry ice. Medium was removed and saved for

Table 5. Rat pituitary receptor binding and LH release or inhibition by naphthylpropionyl-Ser⁴(O-Bzl) and naphthylacetyl-Ser⁴(O-Bzl) (4-9) GnRH analogues. The effect of O-substitution of Ser⁴ with benzyl.

X ³ -Ser(O-Bzl)-Tyr-Z ⁶ -Leu-Arg-PrnHET					
No.	X ³	Z ⁶	pK ₁	pD ₂	pA ₂
33	1-Naphthylpropionyl	DLeu	7.64	7.93	
34	1-Naphthylpropionyl	DTrp	8.20		7.10
35	1-Naphthylacetyl	DLeu	7.70		7.49
36	1-Naphthylacetyl	D-(2)NaI ^a	7.43		8.44
37	1-Naphthylacetyl	DPhe	8.46		7.98
38	1-Naphthylacetyl	DTrp	8.91		8.52

^aD-(2)-NaI = D-2-naphthyl-3-alanine.

CONCLUSIONS

Several novel conclusions emerge from these studies that are very important for further development of LHRH analogues, especially reduced size and nonpeptidic versions. Figure 1 shows a schematic representation of the (3-9)-GnRH molecule. It is clear from the




FIGURE 1 Schematic representation of the (3-9) GnRH hexapeptide series. The arrows represent the interactions that are believed to occur between position 6 and 3 that influence whether the molecule is an agonist or an antagonist




FIGURE 1 Stimulation of superfused pituitary cells by GnRH ($10^{-9}M$) pulses

as hormone release (ng RP-2/ml/ 2×10^7 cells per column) and show one representative experiment out of three sets.

Gonadotrope enrichment

Since the rat pituitary gland contains only about 10% gonadotropes, centrifugal elutriation (Fig. 2), a new method for enrichment of certain cell fractions [19], was used in some experiments. Thus, interference with other non-gonadotrope cell types of the pituitary could be diminished. Eight fractions were obtained by increasing the flow rate step-wise from 11.5 ml/min to 37.5 ml/min.

Fractions 6 and 7 contained 40-50% of gonadotrophs. Cells of each fraction were plated as monolayers and a GnRH challenge test was performed for 4 hours. Highest response was achieved by cells of fraction 6 and 7 (Fig. 3).




FIGURE 2 Centrifugal elutriation for preparation of gonadotrope-enriched cell populations




FIGURE 3 Cell fractions after centrifugal elutriation. GnRH (10^{-7}) challenge for 4 hours to stimulate LH and FSH secretion in initial cell suspension (ICS) and 8 fractions after elutriation. Cells were precultured for 48 hours

PHOSPHOLIPIDS, INOSITOL PHOSPHATES AND PROTEIN KINASE C

It is now recognized that activation of many tissues is accompanied by changes in the phospholipid metabolism which in turn influence the availability and actions of calcium within the target cell. One proposed sequence in ligand-activated cells attributes the earliest regulatory events to the phosphatidylinositol cycle, in which the breakdown of phosphatidylinositol (PI) to inositol phosphate and diacylglycerol is rapidly followed by the formation of phosphatidic acid (PA) and subsequently by re-synthesis of PI [26-28]. Recent findings have demonstrated the rapid change in phospholipid metabolism in rat pituitary cells stimulated by GnRH inducing the increase in ^{32}P labeling of PI [29,30]. Several of the intermediates formed during activation of the PI-cycle have the potential to act as intracellular messengers. The administration of the enzymes PL-C and A_2 which mediate the breakdown of polyphosphoinositides and the liberation of arachidonic acid, respectively, stimulates the release of LH (Figs. 4 and 5). The action of melittin (activator of PL- A_2) on




FIGURE 4 Stimulation of LH secretion in superfused pituitary cells by phospholipase C (PL-C)




FIGURE 5 Stimulation of LH secretion in superfused pituitary cells by phospholipase A₂ (PL-A₂)

gonadotropin secretion is blocked by chloroquine (Fig. 6). PA can function as a calcium ionophore in several cell types [31], and could also be involved in the actions of GnRH on rat pituitary gonadotrophs [30]. GnRH stimulates ³²P incorporation into PI and PA release and cGMP production in a calcium-dependent manner in cultured gonadotrophs [30]. Apart from its calcium ionophore properties, PA is also a source of arachidonic acid, the major precursor of potential intracellular mediators formed via the cyclo-oxygenase and lipoxygenase pathways. However, the mechanistic role of arachidonic acid in secretory granule release remains to be clarified. In earlier studies, GnRH was found to cause arachidonic acid release from cultured anterior pituitary cells, and exogenous arachidonic acid was shown to stimulate LH release in vitro [8,32,33]. Recently, emphasis has been shifted to the breakdown of phosphoinositol-1,4,5-triphosphate (PIP₃), rather than of PI, as the major source of hormonal mediators such as inositol triphosphate and diacylglycerol [20,34].




FIGURE 6 Stimulation of LH secretion in pituitary cell by melittin and the inhibition of its effect by the phospholipase A₂ blocker chloroquine

It is now evident that the diacylglycerol formed during phosphoinositide breakdown is a potent activator of the multifunctional protein kinase (protein kinase C) characterized by Nishizuka et al [35]. This calcium- and phospholipid-dependent enzyme, which is usually present in an inactive form, can be activated by the diacylglycerol that is transiently produced during ligand-stimulated phosphoinositide hydrolysis. In the presence of phosphatidylserine, the concentration of calcium required for activation of protein kinase C is markedly reduced by diacylglycerol, and the total activity of the enzyme is increased.

Protein kinase C is present in the pituitary, mostly as a soluble and partly as a particulate form [36-38]. Stimulation by GnRH induces translocation of protein kinase C from the cytosol to the membrane in the pituitary [36]. Similarly, the tumour promoter 12-O-tetradecanoyl-phorbol-13-acetate (TPA) activates protein kinase C but can elicit only a partial LH release when compared to GnRH.

In studies with myo (2-³H) inositol prelabelled pituitary cultures we [20] and others [39-42] demonstrated that GnRH stimulation enhances the turnover of polyphosphoinositides. This stimulatory effect on inositol phosphate accumulation was shown to be time- and dose-dependent in enriched gonadotropes with maximal

5- to 8-fold increase achieved after 15 min [20]. Significant rise in IP₃ production was observed within 5 sec after GnRH administration (Table 1). Measurement of LH in these experiments

Table 1. Stimulation of inositol 1,4,5-trisphosphate (IP₃) production by GnRH in rat pituitary cells

Stimulation Time	IP ₃ (cpm)		± SD
0 sec	2,429	±	147
5 sec	4,037	±	253
15 sec	3,922	±	308
30 sec	2,559	±	161
45 sec	3,220	±	173
1 min	2,840	±	224
3 min	3,313	±	508
5 min	3,150	±	144
15 min	3,304	±	138

revealed that it occurred later. The effect of GnRH on IP accumulation did not require extracellular calcium in contrast to gonadotropin secretion [20].

In further studies [43,44] the binding of inositol phosphates and the induction of Ca²⁺-release from pituitary microsomal fractions were demonstrated. Bovine anterior-pituitary microsomal fractions exhibit high-affinity, saturable and reversible binding of inositol 1,4,5-(³²P)trisphosphate; 50% of the labelled ligand is displaced by 3.5 nM-inositol 1,4,5-trisphosphate, 0.5 nM-inositol 1,4-bisphosphate and 10 μM-ATP. Inositol 1,4,5-trisphosphate induces the release of Ca²⁺ from the microsomal vesicles (half-maximal effect at 290 nM), and its action is potentiated by inositol tetrakisphosphate (half maximal effect at 4 μM).

These findings on inositol phosphate production and action suggest that they may be a mediator of GnRH-induced gonadotropin release. There are, however, reports providing evidence that inositol phosphates are not the only mediators.

ARACHIDONIC ACID AND METABOLITES

Experiments using labelled arachidonic acid demonstrated the rapid production of metabolites in enriched gonadotrope fractions [46]. In addition, inhibitors of PL-A₂ provided evidence for the role of arachidonic acid and its active metabolites in GnRH-induced

gonadotropin secretion [47]. The stimulation of LH release by arachidonic acid has been demonstrated under both static and dynamic incubation conditions. Arachidonic acid causes an increase in LH secretion in static cultures of enriched gonadotropes [8], and has similar though attenuated effects in perfused gonadotropes (Fig. 7). In static cultures, the LH




FIGURE 7 Stimulation of LH secretion in superfused pituitary cells by GnRH and arachidonic acid

response to GnRH is unimpaired following preincubation with arachidonic acid, indicating that the effects of the fatty acid are not due to non-specific toxic actions on hormone release. In the same context, addition of GnRH a few minutes after arachidonic acid in superfused gonadotrophs potentiated the LH response. The potency of arachidonic acid on gonadotropin release was about 4 orders of magnitude lower than that of GnRH, but the maximal stimulation of LH release was similar to that elicited by GnRH. Other fatty acids, including linoleic acid and oleic acid, had no effect on LH release, further indicating that the action of arachidonic acid does not reflect a non-specific effect on the gonadotrope [8].

In previous reports, GnRH action was found to include release of arachidonic acid [32,47], which serves as a precursor of several groups of intracellular mediators [32,47,48]. However, prostaglandins do not consistently stimulate LH release in vitro,

and GnRH-stimulated gonadotropin secretion is not prevented by cyclo-oxygenase inhibitors [49]. These findings suggest that cyclo-oxygenase products of arachidonic acid are not directly involved in the mechanism of GnRH action. Rather, the effect of prostaglandins on gonadotropin secretion appears to be indirect, via stimulation at the hypothalamic level [31,50].

Currently, two additional pathways of arachidonate metabolism: epoxygenase and lipoxygenase are being investigated. Epoxygenase metabolites of arachidonate, i.e. 5,6-epoxyicosatrienoic acid (5,6-EET) stimulate gonadotropin secretion in a range of 0.01-1 μM [51]. Lipoxygenase metabolites, especially leukotrienes, were found to be even more potent when added to pituitary cells in monolayer culture [32,52] and in a perfusion system [21-24]. The most effective doses were 10^{-10} and 3×10^{-11} M for LTA_4 , LTB_4 , LTC_4 and LTE_4 . In Fig. 8, the




FIGURE 8 Biphasic stimulatory effect leukotriene C_4 (LTC_4) on LH secretion in superfused pituitary cells

effect of LTC_4 is shown. LTD_4 had no stimulatory action on gonadotropin release [23]. When higher concentration was continuously administered to superfused pituitary cells, gonadotropin secretion was suppressed 50% below basal values. Similarly, simultaneous pulses of LTA_4 , LTB_4 , LTC_4 and LTE_4 partially blocked GnRH-induced LH release (Fig. 9). These




FIGURE 9 Stimulation of LH secretion in superfused pituitary cells by leukotriene C₄ (LTC₄, 10⁻¹³ M), GnRH (10⁻⁹ M) or both stimulants. The order of stimulation was changed in two different experiments. LTC₄ inhibited GnRH effect on gonadotropin secretion

findings suggest that leukotrienes can modulate ligand-stimulated hormone secretion in the pituitary [53].

The differential effect of leukotrienes may find its explanation in the following hypothesis. When leukotrienes are present in a certain narrow range of concentration (below 10⁻⁹ M) they can partially mimic the effect of GnRH and potentially induce LH secretion. Higher concentrations, however, seem to be deleterious to basal hormone release.

When NDGA is administered prior to and during leukotrienes, endogenous synthesis is blocked, intracellular concentration of these mediators is very low and therefore exogenously added leukotrienes can exert an even more potent action on gonadotropes.

The understanding of the modulation of GnRH action is, however, more complex especially because not all post-receptor events are quite defined, yet. As observed by others [54] the activation of more than one second messenger pathway seems to be required to fully mimic the GnRH-induced gonadotropin secretion, demonstrated by the addition of arachidonate and phorbol esters.

Therefore the observation made in the present publication only indicates that leukotrienes disrupt the coupling of GnRH-receptor and post-receptor mechanisms.

Others have proposed a complex interpretation of dose-response curves for LTC₄ using a hormone/receptor/effector model [55].

Apart from leukotrienes a number of other potential second messenger mechanisms have been suggested for GnRH-activation, i.e. Ca²⁺ [14], inositol phosphates [20,39-44], protein kinase C [36-38], epoxigenase metabolites of arachidonate [51]. It is up to the present yet unclear how post-receptor actions of GnRH are orchestrated, but previous data show that it is unlikely that inositol phosphates or protein kinase C alone are fully responsible for transduction of transmembrane signalling in gonadotropes [45,54]. Therefore we believe that the present observations regarding the dual mode of leukotriene action provide further evidence for the potential regulatory role of leukotrienes in pituitary hormone secretion.

CONCLUSIONS

The following mechanisms are postulated for the ligand-stimulated intracellular action of GnRH studied in this paper (Fig. 10). GnRH binds to its receptor at the membrane of gonadotropes and activates several intracellular second messengers. First, the hydrolysis of polyphosphoinositides (TPI, DPI) to PI provides inositol phosphates, especially IP₃, which serve as a signal for the mobilization of intracellular pool of Ca²⁺. Second, a further product of polyphosphoinositide breakdown, diacylglycerol, activates protein kinase C causing translocation of this enzyme from the cytosol to the membrane. This multifunctional enzyme may also modulate phospholipid hydrolysis itself. Third, diacylglycerol could serve as a substrate for PLA₂, which releases arachidonic acid. In addition, phosphorylation by protein kinase C may induce mechanisms which activate PLA₂. Together with the above mentioned messengers certain specific pathways of the arachidonic acid metabolism could supply active products which could mediate gonadotropin release via granule fusion and exocytosis. GnRH-stimulated gonadotropin secretion is biphasic and there is now evidence that the initial phase of release may be mediated by the elevation of cytosolic free-Ca²⁺ and the production of potent metabolites of arachidonic acid, i.e. leukotrienes, whereas the activation of protein kinase C could be involved in the second phase of LH release. Recent studies have shown that post-receptor mechanisms of GnRH action can be modulated which are relevant for the understanding of clinically important phenomena such as mechanism of down-regulation for the treatment of sex-hormone dependent diseases as well as the action of steroids during menstrual cycle or in hormonal contraception. The role of protein kinase C in the mechanism of gonadotrope down-regulation is controversial [45], but some data suggest that the activation of this enzyme may be involved in pituitary desensitization [56].




FIGURE 10 Summary of post-receptor actions of GnRH in pituitary cells.

G, GTP-binding protein; DG, diacylglycerol; PIP₂, phosphatidylinositol 4, 5-bisphosphate; IP₃, inositoltriphosphate; PG, prostaglandins; LT, leukotrienes; PK-C, protein kinase C; AA, arachidonic acid

In addition, leukotrienes could be responsible for an intracellular feedback mechanism as shown by the inhibitory effect of leukotrienes on GnRH-induced LH-release, whereas leukotrienes alone are very potent stimulators of gonadotropin secretion [53]. It has been suggested that leukotrienes may modulate GnRH-induced inositol phosphate accumulation.

Steroids are known modulators of gonadotrope function [57]. A detailed analysis of the inhibitory action of progestins has been performed [58] and some experimental data suggest that progestins may also interfere with post-receptor mechanisms, such as IP production [58].

ACKNOWLEDGEMENTS

This study was partially supported by grants of the Deutsche Forschungsgemeinschaft, Bonn (FRG), (Ki 283/2-1 and Ki 283/2-3 and by the Ministry of Baden-Württemberg/FRG (Forschungsschwerpunkt 18) to Dr. L. Kiesel.

REFERENCES

1. Sandow, J (1983): Clinical application of LHRH and its analogues. *Clin Endocrinol*, 18, 571
2. Borgeat, P, Chavancy, G, Dupont, A, Labrie, F, Arimura, A and Schally AV (1972). Stimulation of adenosine 3',5'-cyclic monophosphate in anterior pituitary gland in vitro by synthetic luteinizing hormone-releasing hormone. *Proc Natl Acad Sci USA*, 69, 2677
3. Kaneko, T, Saito, S, Oka, H, Oda, T and Yanaihara, N (1973). Effects of synthetic LH-RH and its analogs on rat pituitary cyclic AMP and LH and FSH release. *Metabolism*, 22, 77
4. Naor, Z, Zor, U, Meidan, R and Koch, Y (1978). Sex difference in pituitary cyclic AMP response to gonadotropin-releasing hormone. *Am J Physiol*, 235, E37
5. Naor, Z, Snyder, G, Fawcett, CP and McCann, SM (1978). A possible role for cGMP in mediating the effect of LH-RH on gonadotropin release in dispersed pituitary cells of the female rat. *J Cyclic Nucleotide Res*, 4, 475
6. Conn, PM, Morrell, DV, Dufau, ML and Catt, KJ (1959). Gonadotropin-releasing hormone action in cultured pituitary cells: independence of luteinizing hormone release and adenosine 3',5'-monophosphate production. *Endocrinology*, 104, 448
7. Naor, Z and Catt, KJ (1980). Independent actions of gonadotropin releasing hormone upon cyclic GMP production and luteinizing hormone release. *J Biol Chem*, 255, 342
8. Kiesel, L and Catt, KJ (1987). Stimulation of luteinizing hormone release and cyclic nucleotide production of arachidonic acid in cultured gonadotrophs. *Neuroendocrinol*, 46, 1
9. Cronin, MJ, Evans WS, Hewlett, EL and Thornee, MO (1984). LH release is facilitated by agents that alter cyclic AMP-generating system. *Am J Physiol*, 246, E44
10. Samli, MH and Geschwind, II (1969). Some effects of energy-transfer inhibitors and of Ca^{++} -free or K^{+} -enhanced media on the release of LH from the rat pituitary gland in vitro. *Endocrinology*, 82, 225
11. Jutisz, M and De la Llosa, MP (1970). Requirement of Ca^{++} and Mg^{++} ions for the in vitro release of follicle-stimulating hormone from rat pituitary gland and its subsequent biosynthesis. *Endocrinology*, 86, 761
12. Hopkins, C and Walker, AM (1978) Calcium as a second messenger in the stimulation of LH secretion. *Mol Cell Endocrinol*, 12, 189
13. Naor, Z, Leifer, AM and Catt, KJ (1980). Calcium-dependent actions of GnRH on pituitary guanosine 3',5'-monophosphate production and gonadotropin release. *Endocrinology*, 107, 1438
14. Conn, PM, Rogers, DC and Sandhu, FS (1979). Alteration of the intracellular calcium levels stimulates gonadotropin release from cultured rat anterior pituitary cells. *Endocrinology*, 105, 1122
15. Marian and J, Conn, PM (1980). The calcium requirement in GnRH-stimulated LH release is not mediated through a specific action on the receptor. *Life Sci*, 27, 87

16. Conn, PM and Rogers, DC (1980). Gonadotropin release from pituitary cultures following activation of endogenous ion channels. *Endocrinology*, 170, 2133
17. Clapper, D and Conn, PM (1985). GnRH stimulation of pituitary gonadotrope cells produces an increase in intracellular calcium. *Biol Reprod*, 32, 269
18. Conn, PM, Rogers, DC and Sheffield, T (1981). Inhibition of gonadotropin releasing hormone stimulated luteinizing hormone release by Pimozide: Evidence for a site of action after calcium mobilization. *Endocrinology*, 109, 1122
19. Hyde, CL, Childs, MG, Wahl, LM, Naor, Z and Catt, KJ (1982). Preparation of gonadotroph-enriched cell populations from adult rat anterior pituitary cells by centrifugal elutriation. *Endocrinology*, 111, 1421
20. Kiesel, L, Bertges, K, Rabe, T and Runnebaum, B (1986). Gonadotropin releasing hormone enhances polyphosphoinositide hydrolysis in rat pituitary cells. *Biochem Biophys Res Commun*, 134, 861
21. Kiesel, L, Przylipiak, A, Rabe, T and Runnebaum, B (1986). The role of novel arachidonic acid metabolites in GnRH action on gonadotrophin release in vitro. *Human Reprod*, 1, 349
22. Kiesel, L, Przylipiak, A, Emig, E, Rabe, T and Runnebaum, B (1987). Stimulation of gonadotropin release by arachidonic acid and its lipoxygenase metabolites in superfused pituitary cells. *Life Sci*, 40, 847
23. Kiesel, L, Przylipiak, A, Rabe, T and Runnebaum, B (1987). Leukotrienes stimulate gonadotropin release in vitro. *Gynecol Endocrinol*, 1, 25
24. Kiesel, L, Przylipiak, A, Rabe, T, Przylipiak, M and Runnebaum, B (1987). Arachidonic acid and its lipoxygenase metabolites stimulate prolactin release in superfused pituitary cells. *Human Reprod*, 2, 281
25. Knepel, W, Schwaninger, M, Helm, C and Kiesel, L (1985). Top concentrations of dynorphin-like immunoreactivity in fractions of rat anterior pituitary cells enriched in gonadotrophs. *Life Sci*, 38, 2363
26. Hokin, MR and Hokin, LE (1953). Enzyme secretion and the incorporation of ^{32}P into phospholipids of pancreas slices. *J Biol Chem*, 203, 967
27. Lapentina, EG, Billah, NM and Cuatrecasas, P. (1981). The phosphatidylinositol cycle and the regulation of arachidonic acid production. *Nature*, 292, 367
28. Michell, RH (1975). Inositol phospholipids and cell surface receptor function. *Biochem Biophys Acta*, 415, 81
29. Snyder, GD and Bleasdale, JE (1982). Effects of LHRH on incorporation of (^{32}P)-orthophosphate into phosphatidylinositol by dispersed anterior pituitary cells. *Mol Cell Endocrinol*, 28, 55
30. Kiesel, L and Catt, KJ (1984). Phosphatidic acid and the calcium-dependent actions of gonadotropin-releasing hormone in pituitary gonadotrophs. *Arch Biochem Biophys*, 321, 202
31. Putney, JW (1981). Recent hypotheses regarding the phosphatidylinositol effect. *Life Sci*, 29, 1183

32. Naor, Z and Catt, KJ (1981). Mechanism of action of gonadotropin releasing hormone. *J. Biol Chem*, 256, 2226
33. Naor, Z, Kiesel, L, Vanderhoek, J and Catt, KJ (1985). Mechanism of action of gonadotropin releasing hormone: role of lipoxygenase products of arachidonic acid in luteinizing hormone release. *J Steroid Biochem*, 23, 711
34. Berridge, M and Irvine, R (1984). Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature*, 312, 315
35. Nishizuka, Y (1984). The role of protein kinase C in cell surface signal transduction and tumor promotion. *Nature*, 308, 693
36. Hirota, K, Hirota, T, Aguilera, G and Catt KJ (1984). Hormone-induced redistribution of calcium-activated phospholipid-dependent protein kinase in pituitary gonadotrophs. *J Biol Chem*, 260, 3243
37. Catt, KJ, Loumaye, E, Wynn, P, Suarez-Quian, C, Kiesel, L, Iwashita, M, Hirota, K, Morgan, R and Chang, J (1984). Receptor-mediated activation mechanisms in the hypothalamic control of pituitary-gonadal function. In: Labrie, F and Proulx, L (eds.) "Endocrinology", p 57. (Amsterdam; Elsevier Science Publishers BV)
38. Naor, Z (1986). Phosphoinositide turnover, Ca^{2+} mobilization, protein kinase C activation and leukotriene action in pituitary signal transduction: effect of gonadotropin releasing hormone. In: Zor, U, et al (eds) "Adv Prostagl Thrombox Leukotr Res". p 225. (New York: Raven Press)
39. Schrey, MP (1985). Gonadotropin releasing hormone stimulates the formation of inositol phosphates in pituitary tissue. *Biochem J*, 226, 563
40. Martin, TJF (1986). Measurement of phospholipid turnover in cultured hormone responsive pituitary cells. *Methods Enzymol*, 124, 424
41. Naor, Z, Molcho, J, Zakut, H and Yavin, E (1985). Calcium-independent phosphatidylinositol response in gonadotropin-releasing-hormone-stimulated pituitary cells. *Biochem J*, 231, 19
42. Andrews, WV and Conn, PM (1986). Gonadotropin-releasing hormone stimulates mass changes in phosphoinositides and diacylglycerol accumulation in purified gonadotrope cell cultures. *Endocrinology*, 118, 1148
43. Kiesel, L, Lukacs, GL, Eberhardt, I, Runnebaum B and Spät, A (1987). Effect of inositol 1,4,5-triphosphate and GTP on calcium release from pituitary microsomes. *FEBS Lett*, 217, 85
44. Spät, A, Lukacs, GL, Eberhardt, I, Kiesel, L and Runnebaum, B (1987). Binding of inositol phosphates and induction of Ca^{2+} release from pituitary microsomal fractions. *Biochem J*, 244, 493
45. McCardle, CA, Huckle WR and Conn, PM (1987). Phorbol esters reduce gonadotrope responsiveness to protein kinase C activators but not to Ca^{2+} -mobilizing secretagogues. *J Biol Chem*, 262, 5028
46. Vanderhoek, JY, Kiesel, L, Naor, Z, Bailey, JM and Catt, KJ (1984). Arachidonic acid metabolism in gonadotroph-enriched pituitary cells. *Prostagl Leukotr Med*, 15, 375

47. Kiesel L, Rabe, T, Hauser, G, Przylipiak, A, Jadali, F and Runnebaum, B (1987). Stimulation of luteinizing hormone release by melittin and phospholipase A₂ in rat pituitary cells. *Mol Cell Endocrinol*, 51, 1
48. Irvine, RF (1982). How is the level of free arachidonic acid controlled in mammalian cells? *Biochem J*, 204, 3
49. Naor, Z, Koch, Y, Bauminger, S and Zor, U (1975). Action of luteinizing hormone releasing hormone and synthesis of prostaglandin in the pituitary gland. *Prostaglandins*, 9, 211
50. Hedge, GA (1977). Roles for the prostaglandins in the regulation of anterior pituitary secretion. *Life Sci*, 20, 17
51. Snyder, GD, Capdevila, J, Chacos, N, Manna, S and Falck, JR (1983). Action of luteinizing hormone-releasing hormone: Involvement of novel arachidonic acid metabolites. *Proc Natl Sci USA*, 80, 3504
52. Hulting AL, Lindgren, JA, Hokfelt, T, Eneroth, P, Werner, S, Patrono, C and Samuelsson, B (1985). Leukotriene C₄ as a mediator of luteinizing hormone release from rat anterior pituitary cells. *Proc Natl Sci USA*, 82, 3834
53. Kiesel, L, Przylipiak, A, Rabe, T, Przylipiak, M and Runnebaum, B (1988). Modulation of LH release by leukotrienes in rat pituitary cells in vitro. *Mol Cell Endocrinol* (in press)
54. Chang, JP, Graeter, J and Catt, KJ (1987). Dynamic action of arachidonic acid and protein kinase C in pituitary stimulation by gonadotropin-releasing hormone. *Endocrinology*, 120, 1837
55. Leiser, J, Conn, PM and Blum, JJ (1986). Interpretation of dose-response curves for luteinizing hormone release by gonadotropin-releasing hormone, related peptides, and leukotriene C₄ according to a hormone/receptor/effector model. *Proc Natl Acad Sci USA*, 83, 5963
56. Helm, KJ, Kiesel, L, Rabe, T, Przylipiak, A and Runnebaum, B (1988). The role of protein kinase C in desensitization of rat pituitary cells induced by gonadotrophin-releasing hormone. *Acta Endocrinol (Suppl)* (in press)
57. Kamel, F and Krey, LC (1982). Intracellular receptors mediate gonadal steroid modulation of LHRH-induced LH release. *Mol Cell Endocrinol*, 28, 471
58. Kiesel, L, Helm, K, Bertges, K, Maier, C, Rabe, T and Runnebaum, B (1987). Contraceptive progestins and gonadotropin secretion in vitro. *J Steroid Biochem*, 27, 995

EXTRAPITUITARY ACTIONS

J.G.M. KLIGN and J.A. FOEKENS

Division of Endocrine Oncology, Rotterdam Cancer Institute
 "The Dr. Daniel den Hoed Cancer Center", Groene Hilledijk 301,
 3075 EA Rotterdam, The Netherlands

INTRODUCTION

Extrapituitary actions of LHRH and its analogues *in vivo* have been suggested by the presence of LHRH-like peptides and LHRH receptors in various extrapituitary tissues and by observed biological effects *in vitro* or in hypophysectomized animals after administration of LHRH analogues.

Hypothalamic hormones are not only detected in the hypothalamus, but also in the central nervous system and in different normal and tissue tumors. LHRH-like peptides have been found in tissues derived from the pancreas, in the gonads, in cultures of Sertoli cells, in the placenta and in human biological fluids (Table 1); in addition they have been found in different human tumors including insulinomas, choriocarcinomas and breast cancers. The meaning of the presence and local production of hypothalamic peptides in peripheral tissues is not clear, but probably they have a local regulatory function by autocrine or paracrine mechanisms of action.

Table 1. LHRH-like material has been demonstrated in:

- Hypothalamic neurons
- Brain
- Spinal ganglia
- Pineal gland
- Pancreatic islet cells, insulinoma
- Ovary, testis
- Placenta, Choriocarcinoma
- Mammary carcinoma cells
- Biological fluids,
 - plasma
 - urine
 - milk
 - ovarian follicular fluid
 - semen

Specific binding sites for LHRH have been characterized in the anterior pituitary, brain, testis, ovary, placenta, adrenals and in various tumors (Table 2). In general, human extrapituitary tissues contain LHRH receptors which have a considerably lower affinity for LHRH than do the human pituitary or the rat testis and ovary [1]. It has been suggested that LHRH receptors in various tumors are induced by carcinogenic transformation of normal cells in view of the absence of these receptors in most non-neoplastic tissues [2].

Table 2. LHRH (analogue) binding sites

<u>High affinity</u> (Kd 10^{-8} - 10^{-9} M)	<u>Low affinity</u> (Kd 10^{-5} - 10^{-8} M)
- Rat pituitary	- Human corpus luteum
- Human pituitary	- Human ovarian tumors
- Rat testis (Leydig cells)	- Placenta
- Rat ovary: interstitial cells	- Adrenals
luteal cells	- Mammary carcinoma cell lines
granulosa cells	Human primary breast cancers
	- Dunning R-3327 H prostate tumor
	Prostate tumor cell lines
	- Leiomyoma

- Discrete brain regions

Direct biological effects of LHRH have been demonstrated in vitro or in hypophysectomized animals on cells from the gonads, accessory sex organs, placenta, kidney and on mammary and prostate tumor cells (Table 3). However, species variation exists concerning the direct action of LHRH and its analogues [3]. Initially, there was great interest in the extrapituitary actions on normal tissues (see reviews [1,3-6]); later, more interest was shown in the direct growth inhibitory effects on tumor cells.

TESTIS

Hsueh and Erickson [7] showed for the first time that treatment with LHRH analogues decreased the testicular weight and testicular steroidogenesis in hypophysectomized male rats. After demonstration of direct inhibition of androgen secretion by LHRH in vitro, from cultured rat testicular cells, Hsueh et al showed that the inhibitory effect of LHRH on testicular androgen production occurs at sites distal to the formation of cAMP and pregnenolone. They and others attributed the effect as being due to decreases in the activity of the enzymes 17- α -hydroxylase and 17,20-desmolase [8,9]. There was a dose-dependent inhibition

of gonadotropin-stimulated Leydig cell production of testosterone by LHRH and its agonists [8-10]. LHRH receptors have been identified by several authors in Leydig but not in Sertoli cells [10-12]. With respect to ontogeny, LHRH receptors were not detectable in rat fetal testes but were present post-natally and increased markedly with age [10]. Administration of LHRH agonists decreased testicular gonadotropin receptors [3,7], but administration of LH reduced both basal and LHRH-stimulated LHRH binding sites in cultured Leydig cells [10].

Table 3. Direct effects of LHRH analogues

1. Interstitial, luteal and granulosa cells of rat ovaries in vitro.
2. Rat testicular cells in vitro.
3. Placental tissue in vitro.
4. Mammary tumor cells in vitro.
5. Prostate tumor cells in vitro.
6. Inhibition of estrogen-induced growth of the uterus in ovariectomized hypophysectomized rats in vivo.
7. Inhibition of androgen induced growth of accessory sex organs in hypophysectomized castrated rats in vivo
8. Inhibition of androgen stimulated enzyme activity in mouse kidney in vivo.

The concentration of LHRH in peripheral blood is too low to postulate direct effects of hypothalamic LHRH on gonads. Therefore several laboratories have attempted to isolate gonadal LHRH-like peptides which may serve as the ligand for specific gonadal LHRH receptors. LHRH or LHRH-like molecules can indeed be isolated from rat testes (see review [5]). Such LHRH-like factors have been detected in extracts of rat gonads and in spent media from cultured Sertoli cells [13]. The observations of high molecular weight LHRH-like factors [14] together with the reported testicular concentration of LHRH of 1 pg/testis [15] and the existence of a precursor molecule for LHRH, alleged to be present in gonads [16], represent the available experimental evidence for intratesticular production of LHRH or LHRH-like factors [5]. Therefore a local paracrine regulatory function of gonadal LHRH or LHRH-like molecules was proposed [13]. However, in human testes LHRH-like material and receptors have not been demonstrated [17]. In addition, several groups do not find evidence for direct inhibitory effects of LHRH agonists on testicular steroidogenesis in men. Administration of LHRH agonists did not inhibit hCG-induced testosterone secretion in patients with gonadotropin deficiency [18] or hCG-induced intratesticular enzyme activities in patients with disseminated prostatic cancer [19]. It needs to be pointed out, however, that it is still conceivable that large doses of hCG/hLH may have overcome a small direct gonadal effect of the administered LHRH analogues [20]. The lack of detection of LHRH receptors in human testis could also be due to

down-regulation of LHRH receptors in the human gonads by the high circulating levels of bioactive LH [10]. Nevertheless, most evidence points to a predominant pituitary site of action of LHRH analogues in the human male.

OVARY

As for the rat testis [7], LHRH analogues inhibit FSH-induced changes in ovarian function in hypophysectomized rats [21]. In addition, LHRH and its agonists have been reported to have direct inhibitory effects on rat ovarian steroidogenesis in vitro [21,22]. Direct stimulatory and inhibitory effects of LHRH analogues on granulosa, luteal and theca cells have been demonstrated [3]. The early effects are mostly stimulatory and the later effects inhibitory [1,3]. In granulosa cells the major action of LHRH appears to be stimulation of 20 alpha-hydroxy-steroid dehydrogenase which causes increased metabolism of progesterone. LHRH also increases aromatase activity in these cells [3]. On the other hand, LHRH agonist delayed hCG-stimulated accumulation of cyclic AMP in porcine granulosa cells obtained from medium follicles [23]. This delay of cyclic AMP accumulation has been suggested to be responsible for the inhibition of progesterone production by ovarian cells [23]. In addition, treatment with high concentrations of LHRH can also induce ovulation and oocyte maturation in hypophysectomized rats, which is associated with the ability of LHH to stimulate plasminogen activator activity in cultured granulosa cells [3]. With respect to mechanism of action, recent studies have shown that LHRH action in the ovary is dependent upon calcium mobilization and probably operates through stimulation of phospholipid turnover and activation of protein kinase C [24].

In the rat ovary, specific high affinity LHRH receptors have been identified in granulosa and theca cells [3,22]. These binding sites mediate the inhibitory effects of LHRH and its agonists on gonadotropin-stimulated estrogen, progesterin and androgen biosynthesis. In addition, LHRH appeared to increase its own receptor in follicular or luteal tissue which is functionally active [25]. LHRH also stimulates prolactin receptor content in rat ovaries [24].

With respect to the effects of LHRH and its analogues on the human ovary the data are conflicting. Tureck et al [26] reported that an LHRH agonist is able to inhibit the secretion of progesterone by cultured human granulosa cells in a dose dependent manner. The effect was prevented by the addition of an LHRH antagonist. On the other hand Casper et al found no effect of LHRH or its agonist on progesterone production by granulosa cells [27]. Comparable conflicting data have been reported with respect to the presence of LHRH receptors in human ovaries. Low affinity LHRH receptors have been identified in human corpus luteum [28]. Bramley et al [29] showed marked variation of specific binding of LHRH agonist to human corpora lutea at different stages of the

luteal phase. In contrast, Clayton and Huhtaniemi [17] reported the absence of LHRH receptors in corpus luteum.

Substances with LH releasing properties in extracts from rat ovaries were first described by De Jong et al [30]. Later on, the rat ovary appeared to contain significant amounts of LHRH-like protein [31]. In contrast, Fraser et al [1] did not find an LHRH-like substance in ovaries. Recently Aten et al [31] detected equivalent amounts of LHRH-like activity in both human and rat ovaries, which was not native LHRH. The LHRH-like material in the human ovary appeared similar to that of the rat ovary. The meaning of this ovarian LHRH-like material remains speculative, but indicates that direct effects of LHRH analogues on human ovaries is possible. However, most evidence suggests that the primary antifertility effect of LHRH analogues in primates is through an action on the pituitary [1].

ACCESSORY SEX ORGANS

Uterus

LHRH agonists have been reported to have an inhibitory action upon the estrogen-stimulated growth of the uterus in ovariectomized hypophysectomized rats [32,33]. LHRH analogue treatment does not alter the estradiol receptor content either in vivo or in vitro [34]. Reddy et al [34] showed that LHRH analogues inhibited estradiol-induced ornithine decarboxylase and glucosamine-6-phosphate synthase activities in the uterus of rat. Although there is no evidence for direct effects of LHRH agonists on the human uterus, specific binding sites for LHRH in uterine leiomyoma have recently been demonstrated [35].

Prostate and seminal vesicles

LHRH agonists have been shown to have direct anti-androgenic effects by inhibition of androgen-induced growth of seminal vesicles and prostate [33]. In addition, similarly to the uterus, LHRH inhibited testosterone-stimulated ornithine decarboxylase activity in the ventral prostate of rat [34]. LHRH receptors have been demonstrated in an experimental prostate tumor [36], but thus far not in the normal human prostate or in primary human prostate tumors.

ADRENALS AND KIDNEY

Adrenals

Specific LHRH receptors have been detected in mouse [37] and rat adrenals [25]. Recently Eidne et al [38] demonstrated a 60 K molecular weight LHRH receptor in solubilized rat adrenal membranes by a ligand-immunoblotting technique.

Kidney

LHRH analogues have antisteroidal activity in the rat kidney. Androgen responsive mouse kidney beta-glucuronidase induction could be inhibited by LHRH analogues [39]. Recently, Prasad et al [40] showed that an LHRH agonist caused significant inhibition of androgen-induced renal ornithine decarboxylase and poly(A)-polymerase activity.

PLACENTA

Khodr and Siler-Khodr [41] have reported the presence of LHRH in human placenta and have also shown stimulatory effects of LHRH on the synthesis and secretion of human chorionic gonadotropin. Addition of LHRH antibodies [5] or LHRH antagonist [42] to the medium completely blocked the action of LHRH on placental hCG and progesterone secretion. Tan and Rousseau [43] demonstrated LHRH and high molecular weight LHRH-like compounds in human trophoblasts.

Specific binding sites for LHRH have been demonstrated in placental tissue [1,44-46]. It has been postulated by Iwashita et al [46] that placental low affinity binding sites may be activated by high local concentrations of a placental LHRH-like peptide. A gene for LHRH has been isolated from the human placenta [47]. It seems that the placenta is the only other tissue besides the pituitary where LHRH has probably a regulatory role in the human female. The potential clinical relevance of these findings might be the application of LHRH antagonists as post-coital contraceptives and abortifacients in humans [6].

TUMOR CELLS

Breast cancer

The presence of direct effects of LHRH analogues on mammary tumor cells is supported by the following findings: 1) presence of LHRH in milk; 2) LHRH-like immunoreactivity in mammary tumor cells; 3) specific LHRH binding sites in human mammary cell lines and human primary breast tumors; 4) direct biological effects in vitro.

Several authors found LHRH in milk [48-50]. Sarda and Nair [49] demonstrated that the LHRH concentration in milk was 6-fold that determined in plasma. They concluded that passive diffusion of LHRH may only contribute to a very small extent towards this milk LHRH. However, the values quoted in the literature for milk LHRH are relatively low.

In 1980 Seppälä et al found LHRH-like material in breast ductal carcinoma [51]. Recently Bützow et al [52] demonstrated immunoreactive LHRH in 4 mammary carcinoma cell lines. In 2 cell lines hCG was also demonstrated, but could not be stimulated by exogenous LHRH [42,45].

Specific LHRH binding sites of low affinity (K_d 10^{-5} - 10^{-6} M) have been identified in 7 mammary cancer cell lines i.e. MCF-7 [53-55], T-47-D [54], MDA-MB-231 [54,55], MDA-MB-157 [52,55], ZR-75-1 [55], SK-BR-3 [55], and UCT-Br-1 [55]. In human primary tumors, Eidne et al [56] found specific LHRH receptors in 67%. The incidence was higher in ductal (86%) than in lobular carcinoma (22%). They found no relationship with the estradiol receptor (ER), progesterone receptor (PR) or age. Binding sites for LHRH were not detected in normal breast tissue. Kaufmann et al [57] found LHRH binding in 38 out of 44 (86%) tumors. Patients with values of ≥ 3.0 fmol/mg protein (48%) were classified as LHRH receptor positive. In contrast to Eidne, these authors found higher values in ER-positive tumors than in ER-negative tumors. Miller et al also reported low affinity binding sites (K_d 10^{-5} M) in some primary tumors [1].

Several direct biological effects on mammary tumor cells in vitro have been demonstrated. In 1982 Corbin [58] and Matzuzawa and Yamamoto [59] showed direct growth inhibitory effects of LHRH agonists on mouse mammary tumor cells in vitro. In 1983 we demonstrated inhibition of estradiol stimulated growth of human breast cancer cells in vitro by the LHRH agonist buserelin [60-62]. One year later Wiznitzer and Benz [63] showed growth inhibitory effects by a LHRH agonist with respect to prolactin-stimulated growth of another cell line. In 1985 and 1986 Miller et al [53,54] observed not only inhibition of growth, but even regression in number of tumor cells by doses from 10^{-9} to 10^{-6} M buserelin. In 1986 we reported that tamoxifen partly can abolish the inhibition of estradiol stimulated growth of MCF-7 cells by buserelin [64,65]. Buserelin also inhibited the expression of prostaglandin receptors induced by estradiol [64,65]. There were no effects on the levels of cytoplasmic or nuclear ER, no change in the patterns of secreted proteins nor on the secretion of polypeptides with EGF-like activities. Both Miller et al [53,54] and our group [61,64,65] showed that concomitant administration of LHRH antagonists prevented the growth inhibitory effects of LHRH agonists on MCF-7 cells, and that the presence of estradiol concentrations higher than 30 pmol/l ($\geq 10^{-10}$ M) or insulin in the medium abolished the growth inhibiting effects of LHRH agonists. Native LHRH also appeared capable of inhibiting tumor cell growth but required higher concentrations. Furthermore, Miller et al [54] using different steroid receptor positive and negative cell lines showed parallelism between sensitivity to LHRH agonist and both responsiveness to estrogen and levels of ER. In contrast, Eidne et al [55] found no effect of LHRH agonists on various cell lines, but showed LHRH antagonists (10^{-5} to 10^{-6} M) to inhibit [3 H]thymidine-uptake within 3 hours leading to inhibition of cell growth after 6 days. Also de Launoit et al [66] showed a rapid antimitogenic effect of LHRH analogue treatment. An LHRH agonist caused a decrease of the labeling index from 12-48 hours after administration in ovariectomized hypophysectomized mice with MXT mammary tumors. On the other hand, we found no effect of LHRH agonist administration on percentage of MCF-7 cells in S-phase of

the cell cycle, when the cells were cultured in serum-free medium in the absence or presence of estradiol for up to 48 hours.

Negative results of LHRH treatment of tumor cells have also been reported. Furr and Nicholson [67] found no anti-estrogenic effects of LHRH agonists in ovariectomized immature rats. Nicholson et al [68] were not able to detect tumor growth inhibiting effects of intratumoral LHRH agonist implants in DMBA rat mammary tumors. They also did not find inhibition of estradiol supported growth of MCF-7 xenografts in ovariectomized nude mice using the LHRH agonist goserelin. Recently Wilding et al [69] reported no effect of high dose LHRH agonists (10^{-6} M) on doubling time of estradiol-stimulated MCF-7 cells. However, their negative results can be explained by the use of a high dose of estradiol (10^{-8} M), which may have overruled the suppressing effects of the LHRH agonists.

PROSTATE CANCER

In 1986 we reported inhibition of androgen (10^{-11} M) stimulated growth of LNCaP human prostate cancer cells in vitro by 0.8 μ M buserelin [65]. Schally and Redding [2] also demonstrated inhibition of this cell line by 10^{-2} to 10^{-6} M tryptorelin. In addition these authors reported inhibition of 2 other prostate tumor cell lines (DU-145, PC-3) and inhibition of the growth of the Dunning R-3327H-G8-A1 clonal prostate adenocarcinoma cell line at doses as low as 10^{-8} of the agonist D-Trp⁶-LHRH. Specific receptors for LHRH were demonstrated in the Dunning rat prostatic tumor [36]. Recently Scaletsky et al [70] were able to demonstrate LHRH receptors with low affinity (K_d of 10^{-6} M) in both LNCaP and DU145 cell lines. Interestingly, in the LNCaP line, incubation with an LHRH agonist led to a dose dependent increase in receptor binding. However, thus far in human primary prostate tumors neither LHRH-like molecules nor LHRH receptors have been found (Reubi et al, personal communication).

OVARIAN CANCER

Kullander et al [71] reported inhibition by LHRH agonist of the growth of rat ovarian tumors autografted under the splenic capsule. Recently Emons [72] demonstrated specific low affinity binding sites in 80% of human primary ovarian tumors. In some small series of patients with progressive disease after first-line chemotherapy, treatment with LHRH agonist showed growth inhibiting effects in several patients [71,73,74]. However, it is unclear whether this tumor growth inhibition was caused by direct or indirect endocrine effects. Direct inhibition on testosterone secretion from arrhenoblastoma cells cultured in vitro in the presence of an LHRH agonist has been shown [75].

CONCLUSIONS

LHRH-like peptides and specific binding sites for LHRH are demonstrable in various normal and malignant extrapituitary tissues. However, these specific binding sites in extrapituitary tissues are of low affinity with the exception of those detected in rat testis and ovary. The importance of the presence of LHRH-like molecules in different extrapituitary tissues remains unclear. Autocrine or paracrine actions have been suggested.

There exists a large species variation with respect to direct actions of LHRH and its analogues.

Although evidence exists from experimental animals for a direct antigonadal effect of LHRH agonists, such data in humans remain controversial. The demonstrated direct effects of LHRH analogues in vitro are probably of low significance in vivo, especially in humans.

REFERENCES

1. Fraser, HM, Bramley, TA, Miller, WR and Sharpe, RM (1986). Extra-pituitary actions of LHRH analogues in tissues of the human female and investigation of the existence and function of LHRH-like peptides. In: Rolland, R, Chadha, DR and Willemsen, WNP (eds.) "Gonadotropin Down-Regulation in Gynecological Practice". p.29. (New York: Alan R. Liss Inc.)
2. Schally, AV and Redding, TW (1987). Use of LH-RH analogs for the treatment of prostate cancer: combination therapy and direct effects. In: Klijn, JGM, Paridaens, R and Foekens, JA (eds.) "Hormonal Manipulation of Cancer: Peptides, Growth Factors, and New (Anti)Steroidal Agents". EORTC Monograph Series. vol. 18, p.273. (New York: Raven Press)
3. Hsueh, AJW and Schaeffer, JM (1985). Gonadotropin-releasing hormone as a paracrine hormone and neurotransmitter in extrapituitary sites. *J Steroid Biochem*, 23, 757
4. Das, C (1985). Extra-pituitary action of gonadotropin releasing hormone: possible application. *J Biosci*, 7, 203
5. Rommerts, FFG and Themmen, APN (1986). LHRH, the role of LHRH (agonists) in the regulation of gonadal function. *Acta Endocrinol. Suppl* 276, 76
6. Clayton, RN (1987). Gonadotrophin releasing hormone: from physiology to pharmacology. *Clin Endocrinol*, 26, 361
7. Hsueh, AJW and Erickson, GF (1979). Extra-pituitary inhibition of testicular function by luteinising hormone releasing hormone. *Nature*, 281, 66
8. Hsueh, AJW, Bambino, TH, Zhuang, LZ, Welsh, TH and Ling, NC (1983). Mechanism of the direct action of gonadotropin-releasing hormone and its antagonist on androgen biosynthesis by cultured rat testicular cells. *Endocrinology*, 112, 1653

9. Labrie, F, Belanger, A, Cusan, L, Seguin, C, Pelletier, G, Kelly, PA, Reeves, JJ, Lefebvre, FA, Lemay, A, Gourdeau, Y and Raynaud, JP (1980). Antifertility effects of LHRH agonists in the male. *J Androl*, 1, 209
10. Dufau, ML and Knox, GF (1985). Fetal Leydig cell culture - an in vitro system for the study of trophic hormone and GnRH receptors and actions. *J Steroid Biochem*, 23, 743.
11. Heber, D, Marshall, JC and Odell, WD (1978). GnRH membrane binding: identification, specificity and quantification in non-pituitary tissues. *Am J Physiol*, 235, E227
12. Clayton, RW, Katikinemi, M, Chan, V, Dufau, ML and Catt, KJ (1980). Direct inhibition of testicular function by gonadotrophin releasing hormone mediation by specific gonadotropin releasing hormone receptors in interstitial cells. *Proc Natn Acad Sci USA*, 77, 4459
13. Sharpe, RM, Fraser, HM, Cooper, I and Rommerts, FFG (1981). Sertoli-Leydig cell communication via an LHRH-like factor. *Nature*, 290, 785
14. Bhasin, S and Swerdloff, RS (1984). Testicular GnRH-like factors: characterization of biologic activity. *Biochem Biophys Res Commun*, 122, 1070
15. Hedger, MP, Robertson, DM, Browne, CA and De Kretser, DM (1985). The isolation and measurement of luteinizing hormone-releasing hormone (LHRH) from the rat testis. *Mol Cell Endocrinol*, 42, 163
16. Nikolics, K, Mason, AJ, Szoniy, E, Ramachandran, J and Seeburg, PH (1985). A prolactin-inhibiting factor within the precursor for human gonadotropin-releasing hormone. *Nature*, 316, 511
17. Clayton, RN and Huhtaniemi, IT (1982). Absence of gonadotropin-releasing hormone receptors in human gonadal tissue. *Nature*, 299, 56
18. Schaison, G, Brailly, S, Vuagat, P, Bouchard, P and Milgrom, E (1984). Absence of a direct inhibitory effect of the gonadotropin releasing hormone (GnRH) agonist D-Ser (TBU)⁶, des-Gly-NH₂¹⁰ GnRH Ethylamide (Buserelin) in testicular steroidogenesis in men. *J Clin Endocrinol Metab*, 58, 885
19. Rajfer, J, Sikka, SC and Swerdloff, RS (1987). Lack of a direct effect of gonadotropin hormone-releasing hormone agonist on human testicular steroidogenesis. *J Clin Endocrinol Metab*, 64, 62
20. Bhasin, S and Swerdloff, RS (1986). Mechanisms of gonadotropin releasing hormone agonist action in the human male. *Endocr Rev*, 7, 106
21. Hsueh, AJW and Erickson, GF (1979). Extrapituitary action of gonadotropin-releasing hormone: direct inhibition of ovarian steroidogenesis. *Science*, 204, 854
22. Clayton, RM, Harwood, JP and Catt, KJ (1979). Gonadotropin-releasing hormone analogue binds to luteal cells and inhibits progesterone production. *Nature*, 282, 90
23. Maruo, T, Otani, T and Mochizuki, M (1985). Antigonadotropic actions of GnRH agonist on ovarian cells in vivo and in vitro. *J Steroid Biochem*, 23, 765

24. Knecht, M, Ranta, T, Feng, P, Shinohara, O and Catt, KJ (1985). Gonadotropin-releasing hormone as a modulator of ovarian function. *J Steroid Biochem*, 23, 771
25. Pieper, DR, Richards, JS and Marshall, JC (1981). Ovarian gonadotropin-releasing hormone (GnRH) receptors: characterization, distribution, and induction by GnRH. *Endocrinology*, 108, 1148
26. Tureck, RW, Mastroianni, L, Blasco, L and Strauss, JF (1982). Inhibition of human granulosa cell progesterone secretion by a gonadotropin-releasing hormone agonist. *J Clin Endocrinol Metab*, 54, 1078
27. Casper, RF, Erickson, GF, Rebar, RW and Yen, SSC (1982). The effect of LHRH factor and its agonists on cultured human granulosa cells. *Fertil Steril*, 37, 406
28. Popkin, R, Bramley, TW, Currie, A, Shaw, RW, Baird, DT and Fraser, HM (1983). Specific binding of LHRH to human luteal tissue. *Biochem Biophys Res Commun*, 114, 750
29. Bramley, TA, Menzies, GS and Baird, DT (1985). Specific binding of gonadotrophin-releasing hormone and an agonist to human corpus luteum homogenates: characterization, properties, and luteal phase levels. *J Clin Endocrinol Metab*, 61, 834
30. De Jong, FH, Welschen, R, Hermans, WP, Smith, SD and Van der Molen, HJ (1979). Effects of factors from ovarian follicular fluid and Sertoli cell culture medium on in vivo and in vitro release of pituitary gonadotropins in the rat: an evaluation of systems for the assay of inhibin. *J Reprod Fertil*, suppl 26, 47
31. Aten, RF, Polan ML, Bayless, R and Behrman, HR (1987). A gonadotropin-releasing hormone (GnRH)-like ovarian protein of the rat. *J Clin Endocrinol Metab*, 64, 1288
32. Pedroza, E, Vilchez-Martinez, JA, Coy, DH, Arimura, A and Schally, AV (1980). Reduction of LHRH pituitary and estradiol uterine binding sites by a superactive analog of luteinizing hormone-releasing hormone. *Biochem Biophys Res Commun*, 95, 1056
33. Sundaram, K, Cao, YQ, Wang, NC, Bardin, CW, Rivier, J and Vale, W (1981). Inhibition of the action of sex steroids by gonadotropin releasing hormone (GnRH) agonists: a new biological effect. *Life Sci*, 28, 83
34. Reddy, PRK, Rao, IM, Raju, VS, Rukmini, V and Reddy, RL (1985). Direct inhibitory actions of GnRH on accessory reproductive organs of rat. *J Steroid Biochem*, 20, 819
35. Wiznitzer, A, Marbach, M, Sharoni, Y, Insher, V and Levy, J (1988). Gonadotropin-releasing hormone specific binding site in uterine leiomyoma. *Gynecol Endocrinol*, 2, Suppl.1, 184 (abstract)
36. Hierowski, MT, Altamirano, P, Redding, TW and Schally, AV (1983). The presence of LHRH-like receptors in Dunning R3327H prostate tumors *FEBS Lett*, 154, 92
37. Bernardo, LA, Petrali, JP, Weiss, LP and Sternberger, LA (1978). Adrenal luteinizing hormone releasing hormone receptors. *J Histochem Cytochem*, 26, 613

38. Eidne, KA, Hendricks, DT and Millar, RP (1985). Demonstration of a 60 K molecular weight luteinizing hormone-releasing hormone receptor in solubilized adrenal membranes by a ligand-immunoblotting technique. *Endocrinology*, 116, 1792
39. Lecompte, P, Wang, NG, Sundaram, K, Rivier, J, Vale, W and Bardin, CW (1983). The antiandrogenic action of gonadotropin-releasing hormone and its agonists on the mouse kidney. *Endocrinology*, 110, 1
40. Prasad, MSK, Vani, VV, Raju, VS and Reddy, PRK (1985). Inhibition of sex-steroid hormone induced ornithine decarboxylase and poly(A)-polymerase activities by a GnRH agonist in the rat kidney. *J Steroid Biochem*, 23, 793
41. Khodr, GS and Siler-Khodr, TM (1980). Placental luteinizing hormone-releasing factor and its synthesis. *Science*, 207, 315
42. Siler-Khodr, TM, Khodr, GS, Vickery, BH and Nestor JJ Jr. (1983). Inhibition of hCG, alpha-hCG, and progesterone release from human placental tissue in vitro by a GnRH antagonist. *Life Sci*, 32, 2641
43. Tan, L and Rousseau, P (1982). The chemical identity of the immunoreactive LHRH-like peptide biosynthesized in the human placenta. *Biochem Biophys Res Commun*, 109, 1061
44. Currie, AJ, Fraser, HM and Sharpe RM (1981). Human placental receptors of luteinizing hormone-releasing hormone. *Biochem Biophys Res Commun*, 99, 332
45. Belisle, S, Guerin, JF, Bellabarba, D. and Lehoux, JG (1984). Luteinizing hormone-releasing hormone binds to enriched human placental membranes and stimulates in vitro the synthesis of bioactive human chorionic gonadotropin. *J Clin Endocrinol Metab*, 59, 119
46. Iwashita, M, Evans, MI and Catt, KJ (1986). Characterization of a gonadotropin-releasing hormone receptor site in term placenta and chorionic villi. *J Clin Endocrinol Metab*, 62, 127
47. Seeburg, PH and Adelman, T (1984). Characterization of cDNA for precursor of human luteinizing hormone-releasing hormone. *Nature*, 311, 66
48. Baram, T, Koch, Y, Hazum, E and Fridkin, M (1977). Gonadotropin releasing hormone in milk. *Science*, 198, 300
49. Sarda, AK and Nair, RMG (1981). Elevated levels of LRH in human milk. *J Clin Endocrinol Metab*, 52, 826
50. Amarant, T, Fridkin, M and Koch, Y (1982). Luteinizing hormone-releasing hormone and thyrotropin releasing hormone in human and bovine milk. *Eur J Biochem*, 127, 647
51. Seppälä, M and Wahlström, T (1980). Identification of luteinizing hormone-releasing factor and alpha subunit of glycoprotein hormones in ductal carcinoma of the mammary gland. *Int. J Cancer*, 26, 267

52. Bützow, R, Huhtaniemi, I, Clayton, R, Wahlström, T, Anderson, LC and Seppälä, M (1987). Cultured mammary carcinoma cells contain gonadotropin-releasing hormone-like immunoreactivity, GnRH binding sites and chorionic gonadotropin. *Int J Cancer*, 39, 498
53. Miller, WR, Scott, WN, Morris, R, Fraser, HM and Sharpe, RM (1985). Growth of human breast cancer cells inhibited by a luteinizing hormone-releasing hormone agonist. *Nature*, 313, 231
54. Miller, WR, Scott, WN, Fraser, HM and Sharpe, RM (1987). Direct inhibition of human breast cancer cell growth by an LHRH agonist. In: Klijn, JGM, Paridaens, R and Foekens, JA (eds.) "Hormonal Manipulation of Cancer: Peptides, Growth Factors and New (Anti)Steroidal Agents". EORTC Monograph Series, vol 18, p.357. (New York: Raven Press)
55. Eidne, KA, Flanagan, CA, Harris, NS and Miller RP (1987). Gonadotropin-releasing hormone (GnRH)-binding sites in human breast cancer cell lines and inhibitory effects of GnRH antagonists. *J Clin Endocrinol Metab*, 64, 425
56. Eidne, KA, Flanagan, CA and Millar, RP (1985). Gonadotropin-releasing hormone binding sites in human breast carcinoma. *Science*, 229, 989
57. Kaufmann, M, Kiesel, L, Klinga, K, Runnebaum, B and Kubli, F (1987). Gonadotropin-releasing hormone receptors in human breast carcinoma. Abstract book 4th EORTC Breast Cancer Working Conference, abstr D 2.22
58. Corbin, A (1982). From contraception to cancer: a review of the therapeutic applications of LHRH analogues as antitumor agents. *Yale J Biol Med*, 55, 27
59. Matsuzawa, A and Yamamoto, T (1982). Enhanced and reversed growth in vitro of a pregnancy-dependent mouse mammary tumor (TPDMT-4) by a gonadotropin-releasing hormone agonist analog. *Europ J Cancer Clin Oncol*, 18, 495
60. Blankenstein, MA, Henkelman, MS and Klijn, JGM (1983). An analogue of LHRH antagonizes the growth-stimulating effect of estradiol on human breast cancer cells in culture (MCF-7). *J Steroid Biochem*, 19 (Suppl), 95 S
61. Blankenstein, MA, Henkelman, MS and Klijn, JGM (1985). Direct inhibitory effect of a luteinizing hormone-releasing hormone agonist on MCF-7 human breast cancer cells. *Eur J Cancer Clin Oncol*, 21, 1493
62. Klijn, JGM, de Jong, FH, Lamberts, SWJ and Blankenstein, MA (1985). LHRH-agonist treatment in clinical and experimental human breast cancer. *J. Steroid Biochem*, 23, 867
63. Wiznitzer, I and Benz, L (1984). Direct growth inhibiting effects of the prolactin antagonists buserelin and pergolide on human breast cancer. *Proc Am Assoc Cancer Res*, 25, 208
64. Foekens, JA, Henkelman, MA, Fekkink, JF, Blankenstein, MA and Klijn, JGM (1986). Combined effects of buserelin, estradiol and tamoxifen on the growth of MCF-7 human breast cancer cells in vitro. *Biochem Biophys Res Commun*, 140, 550

65. Foekens, JA, Henkelman, MS, Bolt-de Vries, J, Portengen, H, Fekkink, JF, Blankenstein, MA, Van Steenbrugge, GJ, Mulder E and Klijn, JGM (1987). Direct effects of LHRH analogs on breast and prostatic tumor cells. In: Klijn, JGM, Paridaens, R and Foekens, JA (eds.) "Hormonal Manipulation of Cancer: Peptides, Growth Factors, and New (Anti)Steroidal Agents". EORTC Monograph Series, vol 18, p.369. (New York: Raven Press)
66. De Launoit, Y, Kiss, R, Danguy, A and Paridaens, R (1987). Effect of ovariectomy, hypophysectomy and/or GnRH analog (HRF) administration on the cell proliferation of the MXT mouse hormone-dependent mammary tumor. *Eur J Cancer Clin Oncol*, 23, 1443
67. Furr, BJA and Nicholson, RI (1982). Use of analogues of luteinizing hormone-releasing hormone for the treatment of cancer. *J Reprod Fert*, 64, 529
68. Nicholson, RI, Walker, KJ, Turkes, A, Dyas, J, Gotting, KE, Plowman, PM, Williams, M, Elston, CW, and Blamey, RW (1987). The British experience with the LH-RH agonist Zoladex (ICI 118630) in the treatment of breast cancer. In: Klijn, JGM, Paridaens, R and Foekens, JA (eds.) "Hormonal Manipulation of Cancer: Peptides, Growth Factors and New (Anti)Steroidal Agents". EORTC Monograph Series, vol. 18, p.331. (New York: Raven Press)
69. Wilding, G, Chen, M and Gelmann, EP (1987). LHRH agonists and human breast cancer cells. *Nature*, 329, 770
70. Scaletsky, R, Qayum, A, Clayton, RN, Sikora, K and Waxman, J (1988). GnRH analogue and dihydrotestosterone treatment regulate expression of GnRH binding in a human hormone dependent prostatic cancer cell line. *Gynecol Endocrinol*, 2, Suppl 1, 100 (abstract)
71. Kullander, S, Rausing, A and Schally, AV (1987). LHRH agonist treatment in ovarian cancer. In: Klijn, JGM, Paridaens, R and Foekens, JA (eds.) "Hormonal Manipulation of Cancer: Peptides, Growth Factors, and New (Anti) Steroidal Agents". EORTC Monograph series, vol. 18, p.353. (New York: Raven Press)
72. Emons, G. (1988). The use of GnRH-analogues in ovarian cancer. *Gynecol Endocrinol*, 2, Suppl 1, 48 (abstract)
73. Parmar, H, Nicoll, J, Stockdale, A, Cassoni, A, Phillips, RH, Lightman, SL and Schally, AV (1985). Advanced ovarian carcinoma: response to the agonist D-Trp-6-LHRH. *Cancer Treatm Rep*, 69, 1341
74. Jäger, WH (1988). GnRH analogues in treatment of ovarian carcinoma. *Gynecol Endocrinol*, 2, Suppl 1, 51 (abstract)
75. Lamberts, SWJ, Timmers, JM, Oosterom, R, Verleun, T, Rommerts, FG and deJong, FH (1982). Testosterone secretion by cultured arrhenoblastoma cells: suppression by a luteinizing hormone releasing hormone agonist. *J Clin Endocrinol Metab*, 52, 450

FIRST GENERATION GnRH ANTAGONIST: OVULATION INHIBITION BY WEEKLY TREATMENT

Daniel KENIGSBERG and Gary D. HODGEN

The Jones Institute for Reproductive Medicine, School of Obstetrics
and Gynecology, Eastern Virginia Medical School, 700 Olney Road,
Lewis Hall, Norfolk, Virginia 23507, USA

INTRODUCTION

Although inhibition of ovulation in the form of combination oral contraceptives has been the most effective strategy for achieving reversible pharmacological fertility control in women, there is increasing interest in contraception achieved through GnRH analogs that act on the pituitary to induce a hypogonadotropic status [1]. GnRH agonists do so after a transient stimulatory phase that lasts from 1-3 weeks [2,3]. In contrast, GnRH antagonists lack stimulatory activity, causing prompt reduction of gonadotropin secretion [4].

To explore the use of a GnRH antagonist to achieve inhibition of ovulation in a practical and potentially clinically useful manner, nonhuman primates were tested in a protocol of intermittent GnRH antagonist administration during the menstrual cycle. Our initial findings prompted a second study aimed at clarifying the mechanism by which the preovulatory LH surge was blocked.

MATERIALS AND METHODS

Experimental subjects

Fifteen adult cynomolgus monkeys with regular ovulatory menstrual cycles were subjected to GnRH antagonist treatment. The methods of feeding, housing, blood collection and drug administration were described previously [5].

Experimental design

Study I. The GnRH antagonist [N-AcD-pCl-Phe¹, D-pCl-Phe², D-Trp³, D-Arg⁶, D-Ala¹⁰]GnRH was a generous gift from the Contraceptive Development Branch, NICHD. Based on previous dose studies demonstrating profound suppression of serum LH and FSH levels by this agent at a daily dose of 1 mg/kg, [4] a higher dose was chosen for weekly administration. Since the dominant follicle

was likely to be selected by cycle day 7 [6], the intermittent dose of GnRH antagonist might achieve its ablation, thereby causing initiation of a new follicular phase. The GnRH antagonist was suspended in sesame oil and injected im.

Eight monkeys were used to test ovulation inhibition. Drug administration began on day 3 after spontaneous onset of menses; a single dose of 10 mg/kg was injected. This initial injection was followed at weekly intervals with single doses of 5 mg/kg on days 10, 17, and 24. During the first 42 days after initiation of the pretreatment menstrual flow, femoral blood was obtained on alternate days. On day 43, the frequency of blood drawing was decreased to twice weekly through day 64, when the study was terminated. Throughout sera were harvested and frozen for subsequent RIA. During the 4 weeks of GnRH antagonist treatment, laparoscopies were performed at weekly intervals to assess ovarian status.

Study II. To clarify the mechanism by which the GnRH antagonist blocked estrogen-induced LH surges in study I, normal ovulatory monkeys received the GnRH antagonist daily (2 mg/kg in sesame oil) from either days 2 through 9 or days 2 through 6 of the menstrual cycle, respectively, and estradiol (E_2) benzoate (50 μ g, iv: n=4) was given on day 6. Because the weekly treatment regimen used in study I gave suppression of pituitary function followed by transient recovery in a series of four episodes, we selected a daily regimen in study II to strive for steadier, more consistent conditions for estrogen challenge and GnRH test of pituitary function. Accordingly, these estrogen or GnRH challenge tests provided an evaluation of pituitary gonadotropin secretory status during the GnRH antagonist regimen; serum LH concentrations served as the principal end point. The same test was given to three control monkeys on day 6 of the normal menstrual cycle.

RIAs for E_2 , progesterone (P_4), LH, and FSH were performed as described previously [6,7]. The monkey LH RIA had an intraassay variation of 1.9-5.0% and an interassay variation of 9.3-18.7%.

Tests for significant differences ($P<0.01$) of the intervals from the onset of menstruation to the next LH surge and/or ovulation and the intermenstrual interval were performed using the F statistic [8].

RESULTS

During the 4-week course of intermittent GnRH antagonist treatment in study I, there were no ovulation sites seen at laparoscopy. Moreover, the absence of ovulation was confirmed by the lack of serum P_4 elevations; also, no LH surges were found.

Mean peripheral serum E_2 levels immediately before and on the day after the four GnRH antagonist treatments are shown in Fig. 1. Interestingly, there was no sustained suppression of E_2 in serum.

MEAN ESTRADIOL BEFORE AND AFTER GnRH ANTAGONIST TREATMENT




FIGURE 1 Mean \pm SEM peripheral E₂ levels immediately before (hatched) and the day after (solid) the four weekly treatments with GnRH antagonist in intact adult monkeys. There was no significant suppression of E₂ levels during these intermittent treatment intervals. d₃. Day 2, etc.

Results from a representative animal with ovulation inhibition and absence of LH surges during treatment as well as resumption of an ovulatory ovarian cycle after treatment are shown in Fig. 2. After discontinuation of treatment with the GnRH antagonist, serum E₂ levels rose within 3 weeks and were followed by timely LH surges in six of eight monkeys.

Ovulation was indicated by typical P₄ elevations in serum and an ensuing luteal phase of normal length. The levels of E₂ and P₄ in the circulation were consistent with single ovulations. The mean interval from the last GnRH treatment to the subsequent preovulatory E₂ peaks and LH surges was approximately 14 days among the 6 monkeys who resumed ovulatory menstrual cycles during the study interval. The remaining 2 monkeys did not resume regular menstrual cycles until nearly a month after completion of this study. Since blood collections did not extend beyond day 64 of the study, we do not know whether these resumed menstrual cycles were ovulatory.

Figure 3 illustrates pituitary refractoriness (three of three monkeys) to the E₂ benzoate challenge test given on cycle day 6 during the daily GnRH antagonist regimen spanning days 2 and 9 (study II). Notice the initial negative feedback effects of exogenous estrogen on LH and FSH secretion, the absence of familiar preovulatory-like LH surges, and finally, the resumption of apparent ovulation about 3 weeks later.




FIGURE 2 Peripheral serum E₂, P₄, and LH levels in a monkey (six of eight) during and after weekly administration of GnRH antagonist (arrows). Note that within about 15 days after the last GnRH antagonist treatment, high serum E₂ levels and a LH surge occurred, followed by an increase in serum P₄. Day 1 was the spontaneous onset of menstruation (top line). Compared to the normal interval from menstruation to ovulation (21) the delay of apparent ovulation and next menses associated with GnRH antagonist treatment was highly significant ($P < 0.01$).

In contrast, administered GnRH (as a 50 μ g bolus dose given on cycle day 6) during the daily GnRH antagonist regimen elicited unambiguous LH responses (four of four monkeys) of seemingly normal strength and duration (Fig. 4). All responses exceeded 4-fold elevations and did not differ significantly from the response of control monkeys given the same iv bolus dose of GnRH (Fig. 5).




FIGURE 3 E₂ benzoate (E₂B) challenge tests in monkeys receiving the GnRH antagonist (2 mg/kg day, im) from menstrual cycle days 2-9. Note the absence of gonadotropin surges, indicative of pituitary refractoriness to estrogen-positive feedback. Without the inhibitory effects of the GnRH antagonist, such exogenous estrogen treatment leads to preovulatory-like LH surges within 48 h [21-23].

In six of the eight monkeys, ovulation resumed promptly after the last GnRH antagonist administration; the first post-treatment estradiol peak occurred 14.3 ± 3.8 (\pm SEM) days after the

final dose. The interval for resumption of ovulatory menstrual cycles after the final GnRH injection is summarized in Table 1.

Table 1. Post-treatment interval to first LH surge and resumption of menstruation

No. of Monkeys	Days	
	E ₂ Peak	Menses
6 of 8	14.3 \pm 3.8 ^a	30.5 \pm 3.4
2 of 8	Unknown	>64; <88

^aMean \pm SEM




FIGURE 4 Serum LH levels in four monkeys receiving a GnRH bolus dose (50 µg, iv) during concurrent administration of GnRH antagonist.




FIGURE 5 Mean \pm SE serum LH responses to a GnRH bolus dose (50 μ g, iv) in three control monkeys on day 6 of the normal menstrual cycle.

DISCUSSION

GnRH agonists effectively inhibit ovulation in women and monkeys when given in either daily or intermittent regimens [2, 3, 9-11]. Characteristically, GnRH agonists initially stimulate pituitary gonadotropin secretion and, in turn, enhance ovarian steroid production [11]. This action is transient; typically, both hypogonadotropic and hypogonadal conditions are achieved within 14-21 days [3,10,12,19]. In contrast, potent GnRH antagonists immediately reduce gonadotropin secretion [4].

Monkeys previously having regular ovarian/menstrual cycles had no ovulation (32 and 32 treatment intervals) when the GnRH antagonist was administered once weekly. After cessation of GnRH antagonist therapy, ovulation resumed promptly in 6 of 8 monkeys. That some monkeys regained ovulatory status as soon as 8-10 days

after the final dose of GnRH antagonist suggests that a weekly treatment interval may be near the upper limit for maintenance of ovulation inhibition at the dose tested here. These observations correlate well with earlier experience in monkeys [13, 14] and women [15] undergoing surgical ablation of the putative dominant follicle, in whom destruction of the largest visible follicle delayed ovulation by causing reinitiation of a new follicular phase. Perhaps each weekly dose of GnRH antagonist caused atresia of the newly selected dominant follicle. The intermittent elevations of E_2 in serum may reflect its secretion by newly recruited follicles in the next growing cohort [13].

It is not clear why two monkeys given the GnRH antagonist had longer delays before resuming regular menstrual cycles. However, their suppressed status ended less than 60 days after the cessation of treatment.

In regard to the mechanism by which the intermittent GnRH antagonist regimen prevented ovulation, the data from study II indicate that the estrogen-positive feedback for the LH surge was negated. During the normal follicular phase, E_2 benzoate will induce midcycle-like LH surges within 48 h; the GnRH antagonist clearly negated this estrogen-positive feedback [16-18]. In contrast, the response to GnRH was normal. It is possible that the iv bolus dose of GnRH displaced the GnRH antagonist from gonadotrope receptors, so that the exogenous GnRH elicited normal LH secretory responses. An alternative interpretation is that the dose of GnRH antagonist was submaximal, since previously we achieved a medical hypophysectomy-like status with only 50% of the dose used here [4].

In considering potential clinical applications derived from these studies, the avoidance of a severely hypoestrogenic milieu in these monkeys is encouraging in view of the concern that sustained reductions in E_2 levels might have undesirable sequelae, including hot flashes, urogenital atrophy, and osteoporosis in young women who might desire fertility control by a GnRH antagonist [19,20]. Even so, if clinical studies are undertaken, the effects of long-term unopposed estrogenic stimulation and the potential risk of endometrial carcinoma prompt consideration of periodic progestin supplementation during intermittent GnRH antagonist administration to accommodate sloughing of proliferated endometrium.

In summary, these studies demonstrate the potential feasibility of a weekly GnRH antagonist regimen to reliably inhibit ovulation in the primate ovarian/menstrual cycle. Further studies are warranted to assess the feasibility of intermittent GnRH antagonist therapy for reversible fertility control in women.

REFERENCES

1. Kenigsberg, D and Hodgen GD (1986). Ovulation inhibition by administration of weekly gonadotropin-releasing hormone antagonist. *J Clin Endocrinol Metab*, 62, 734

2. Monroe, SE, Henzl, MR, Martin, MC, Schriock, E, Lewis, V, Nerenberg, C and Jaffe, RB (1985). Ablation of folliculogenesis in women by a single dose of gonadotropin-releasing hormone agonist: significance of time in cycle. *Fertil Steril*, 43, 361
3. Bergquist C, Nillius, SJ and Wide, L (1979). Inhibition of ovulation in women by intranasal treatment with a luteinizing hormone-releasing hormone agonist. *Contraception*, 19, 497
4. Kenigsberg, D, Littman, BA and Hodgen, GD (1984). Medical hypophysectomy II: variability of ovarian response to gonadotropin therapy. *Fertil Steril*, 42, 116
5. Goodman, AI, Descalzi, CD, Johnson, DK and Hodgen, GD (1977). Composite pattern of circulating LH, FSH, estradiol, and progesterone during the menstrual cycle in cynomolgus monkeys. *Proc Soc Exp Biol Med*, 155, 479
6. Goodman, AI, Nixon, WE, Johnson, DK and Hodgen, GD (1977). Regulation of folliculogenesis in the rhesus monkey: selection of the dominant follicle. *Endocrinology*, 100, 155
7. Niswender, GD, Monroe, SE, Peckham, WD, Midgley, AR, Knobil, E and Reichert, LE (1971). Radioimmunoassay for rhesus monkey luteinizing hormone with anti-ovine LH serum and ovine LH⁻¹³¹I. *Endocrinology*, 88, 1327
8. Fruend, JE, Livermore, PE and Miller, I (1960). *Manual of Experimental Statistics*. Prentice-Hall, Englewood Cliffs
9. Meldrum, D, Chang, J, Lu, J, Vale, M, Rivier, J and Judd, H (1982). "Medical oophorectomy" using a long-acting gonadotropin-releasing hormone agonist: a possible new approach to the treatment of endometriosis. *J Clin Endocrinol Metab*, 54, 1081
10. Lemay, A and Quesnel, G (1982). Potential new treatment of endometriosis: reversible inhibition of pituitary-ovarian function by chronic intranasal administration of a luteinizing hormone-releasing hormone (LH-RH) agonist. *Fertil Steril*, 38, 376
11. Werlin, LB and Hodgen, GD (1983). Gonadotropin-releasing hormone agonist suppresses ovulation, menses and endometriosis in monkeys: an individualized intermittent regimen. *J Clin Endocrinol Metab*, 56, 844
12. Schmidt-Gollwitzer, M, Hardt, W, Schmidt-Gollwitzer, K, von der Ohe, M and Nevinny-Stickel, J (1981). Influence of the LHRH analog Buserelin on cycle ovarian function and on endometrium—a new approach to fertility control. *Contraception*, 23, 18
13. Goodman, AI and Hodgen, GD (1979). Between ovary interaction in the regulation of follicle growth, corpus luteum function, and gonadotropin secretion in the primate ovarian cycle. I Effects of follicle cautery and hemiovariectomy during the follicular phase in cynomolgus monkeys. *Endocrinology*, 104, 1304
14. diZerega, GS, Nixon, WE and Hodgen, GD (1980). Intercycle serum FSH elevations: significance in recruitment and selection of the dominant follicle and assessment of corpus luteum normalcy. *J Clin Endocrinol Metab*, 50, 1046
15. Nilsson, L, Wikland, M and Hamberger, L (1982). Recruitment of an ovulatory follicle in the human following follicle-ectomy and luteectomy. *Fertil Steril*, 37, 30

16. Williams, RF, Johnson and DK, Hodgen, GD (1979). Resumption of estrogen-induced gonadotropin surges in postpartum monkeys. *J Clin Endocrinol Metab*, 19, 422
17. Lundbland, EG, Williams, RF and Hodgen, GD (1980). Rapid recovery of estrogen-induced pituitary gonadotropin surges after luteectomy in rhesus monkeys. *Steroids*, 36, 349
18. Hodgen, GD (1982). The dominant ovarian follicle. *Fertil Steril*, 38, 281
19. Richelson, LS, Wahner, HW, Melton, LJ and Riggs, BL (1984). Relative contributions of aging and estrogen deficiency to postmenopausal bone loss. *N Engl J Med*, 311, 1273
20. Bush, TL, Cowan, LD, Barrett-Connor, E, Criqui, MH, Karon, JM, Wallace, RB, Tyroler, HA and Rifkind, BM (1983). Estrogen use and all-cause mortality. *JAMA*, 249, 903

SECOND GENERATION GnRH ANTAGONIST: CHARACTERIZATION OF PITUITARY RESPONSE IN MONKEYS

**Claudio F. CHILLIK, Joseph ITSKOVITZ, Do Won HAHN,
John L. McGUIRE, Douglas R. DANFORTH and Gary D. HODGEN**
Institute for Reproductive Medicine, Department of Obstetrics
and Gynecology, Eastern Virginia Medical School, 700 Olney Road,
Lewis Hall, Norfolk, Virginia 23507, USA

INTRODUCTION

Previous studies have forecast potential therapeutic uses of GnRH analogs in reproductive endocrinology [1,2,3]. Until now, most human and animal data have been obtained using agonists of GnRH [4-13]. However, the development of useful antagonists of GnRH is now approaching clinical investigation. In some circumstances, these antagonists offer more therapeutic promise than the agonists.

GnRH antagonists are analogs of GnRH having a high affinity for the pituitary GnRH receptor, but with no intrinsic biological activity to stimulate gonadotropin secretion [14]. Persistent administration of a GnRH antagonist suppresses basal levels of gonadotropins creating a reversible medical hypophysectomy [15]. Not only do GnRH antagonists almost immediately suppress basal gonadotropin secretion, they also blunt the response of the pituitary to an estrogen challenge test [16-19].

We designed the following experiment in order to evaluate: (1) dose-response characteristics of a newer generation GnRH antagonist; (2) correlation between initial basal (tonic) gonadotropin levels and days needed to achieve pituitary suppression; and (3) pituitary response to a GnRH challenge test before, during, and after GnRH antagonist administration.

MATERIALS AND METHODS

Four ovariectomized (8 weeks or more) female rhesus monkeys (*Macaca mulatta*) were assigned to this study. Animal husbandry, blood collection techniques, and assays have been previously described [20]. Reagents for FSH/LH assays were obtained from the Hormone Distribution Office, NIAMDD, Bethesda, MD. Coefficients of variation within and between assays were 7.1% and 11.8%, respectively.

DRUG ADMINISTRATION

The GnRH antagonist used in this study was [N-Ac-D-Nal(2)¹,D-pCl-Phe²,D-Pal(3)³,Arg⁵,D-Glu(AA)⁶,D-Ala¹⁰] [21]. The antagonist was dissolved in sesame oil and administered intramuscularly. During the first three days, 2 ml of vehicle was given (Fig. 1). From days 4 to 6, monkeys received 0.3 mg/kg/day of the GnRH antagonist; the dose was increased to 1.0 mg/kg/day from days 7 to 9, and to 3.0 mg/kg/day from days 10 to 12. Lastly, vehicle only was given again from days 13 to 15 during post-treatment recovery.




FIGURE 1 Experimental protocol. Four ovariectomized female monkeys received GnRH antagonist i.m. over 9 days in increasing doses of 0.3, 1.0 and 3.0 mg/kg/day. GnRH (100 µg/iv) was administered on experiment days 4, 7, 10, 13 and 16; blood samples were drawn at 0 and +30 minutes

GnRH CHALLENGE TEST

GnRH (100 µg, intravenously) was given as a bolus injection via a catheter inserted into the saphenous vein. The GnRH challenge test was performed on days 4 (control), 7, 10, 13 (treatment) and 16 (recovery). On these occasions, GnRH was injected 60 minutes before administration of the GnRH antagonist.

SERUM COLLECTION

Blood was drawn daily from the femoral vein on day 1 until experiment day 16 before the injection of GnRH antagonist, as well as at 0 and +30 minutes relative to GnRH administration. Therefore, the levels of LH and FSH reported for a certain day reflect the effect of the antagonist given on the previous day, as well as the acute effects of GnRH challenge tests given intermittently.

LH and FSH levels are expressed in ng/ml. Statistical differences were measured through analysis of variance and correlation coefficients for changes in tonic gonadotropin levels in serum and the response to GnRH challenge tests [22].

RESULTS

Tonic Gonadotropin Profiles

Initial levels of plasma gonadotropins were in the ovariectomized range. After 3 days vehicle administration, there were no meaningful changes in the tonic levels of gonadotropins. However, within 24 hours of administration of 0.3 mg/kg of the GnRH antagonist, there was a decrease in serum levels of FSH and LH. After 3 days of GnRH antagonist treatment at 0.3 mg/kg, tonic FSH and LH concentrations in serum decreased to near minimal detectable limits (Fig. 2). After 3 days of treatment with 1 mg/kg/day of the GnRH antagonist, LH and FSH levels were suppressed below assay detection limits in all monkeys. The decrease was highly significant for both FSH and LH ($p < 0.005$). As expected, when the dose of antagonist was increased to 3 mg/kg/day, gonadotropins remained suppressed. During the recovery interval, despite administration of vehicle alone, tonic serum gonadotropin levels in all monkeys remained undetectable.

Table 1. Duration of antagonist treatment before FSH/LH levels became undetectable

Monkey#	Initial FSH (ng/ml)	Days Until FSH/LH Below Detection Limits	Initial LH (ng/ml)
		FSH/LH	
S172	46	2 / 2	120
0698	88	3 / 3	140
U531	420	6 / 4	192
713H	440	6 / 5	260




FIGURE 2 Tonic serum concentrations of FSH and LH before, during, and after GnRH antagonist treatment. Each point represents the mean of 4 animals. The decrease in serum FSH and LH throughout the study was significant ($p < 0.05$)

The time needed to achieve suppression was not the same in all monkeys. Whereas some needed only two days of treatment to achieve full suppression, others required as much as 6 days of treatment with the GnRH antagonist in order to reach a 'medical hypophysectomy' state. Table 1 demonstrates a correlation between the initial tonic levels of gonadotropins and the days of antagonist administration necessary to achieve suppression. That is monkeys whose pituitaries quickly submitted to GnRH antagonist suppression had begun with lower tonic levels of FSH and LH in serum compared to females having higher initial levels of pituitary gonadotropins. Accordingly, there was a high correlation ($r = 0.992$) between initial FSH levels and the interval of treatment necessary to achieve maximal suppression ($p < 0.01$).

RESPONSE TO GnRH CHALLENGE TEST

After 3 days of treatment with 0.3 mg/kg of GnRH antagonist, bolus administration of 100 μ g of GnRH intravenously elicited the release of gonadotropins not dissimilar from responses under pretreatment conditions (Fig. 3). Upon increasing the dose of




FIGURE 3 Serum concentrations of FSH (3a) and LH (3b) before and 30 minutes after a GnRH challenge test (100 μ g/iv). Each bar represents the mean gonadotropin level \pm SEM, n=4

GnRH antagonist to 1 mg/kg, although tonic secretion of LH and FSH was markedly suppressed, acute responses to the GnRH challenge were again similar to pretreatment tests. Despite 9 days of treatment with increasing doses of the GnRH antagonist and with

gonadotropins consistently below detection limits in all monkeys, the GnRH challenge still elicited a small release of FSH ($p < 0.01$) (Fig. 3a); and the LH secretory response was only slightly inhibited ($p < 0.05$). Interestingly, as rapidly as 3 days post-treatment, although tonic gonadotropins remained suppressed, there was partial restoration of FSH secretory response to the GnRH challenge test, and acute LH release was already fully recovered.

Importantly, we found no histamine release effects either locally or systemically during 9 days of GnRH antagonist therapy at doses up to 3.0 mg/kg/day. This dose is up to 10-fold more than that required to achieve a persistent state of medical hypophysectomy.

DISCUSSION

Our data indicate that within 24 hours of a single dose of 0.3 mg/kg of GnRH antagonist, profound suppression of FSH and LH concentrations in peripheral serum was evident. Depending on the initial tonic levels of LH and FSH, gonadotropins were suppressed below detectable limits after 2 to 6 days of GnRH antagonist administration. Tonic FSH/LH levels remained suppressed during the study interval, even 3 days into the post-treatment recovery interval. These data demonstrate that this potent antagonist of GnRH has a long duration of action.

GnRH antagonists have significant therapeutic advantages over agonists in that they almost immediately inhibit gonadotropin secretion without inducing an initial rise in gonadotropins, as is characteristic of GnRH agonist during the first two to three weeks of treatment [23]. There have been relatively few clinical trials using GnRH antagonists because of the lack of a potent GnRH antagonist without side effects. One of the main side effects of GnRH antagonists was reported by Schmidt et al [24], who demonstrated that administration of certain antagonists of GnRH to rats produced transient edema of the face and extremities, including a cutaneous anaphylactoid-like reaction [25]. This side effect was due to release of histamine by the drug. Among the most potent analogs associated with this histamine release phenomenon are those having structures that include a D-arginine at position 6. The antagonist used in the present study has an arginine substitution in position 5 with a D-glutamine in position 6. Although our observations were not specifically directed at this issue, the combination of structural features avoided discernible histamine release in monkeys.

We found a strong correlation between tonic levels of gonadotropins and the days of GnRH antagonist treatment necessary to achieve full suppression. Using a different GnRH antagonist, Cetel et al [26] reported that the net decrement in gonadotropin concentrations was correlated with tonic values of LH and FSH in women. Thus, a higher dynamic state of pituitary gonadotropin secretion portends greater relative resistance to suppression by the GnRH antagonist in women and monkeys. Therefore, when

evaluating the potency of a GnRH antagonist, it may be important to take into consideration the endocrine milieu of the individual.

Although we were able to decrease the levels of gonadotropins within 24 hours after administration of a single dose (0.3 mg/kg) of GnRH antagonist and were able to achieve undetectable levels of LH and FSH after 6 days of treatment in all monkeys, even 9 days of GnRH antagonist treatment did not fully block FSH or LH secretion during a GnRH challenge test. Indeed, LH secretory responses were hardly affected. This agrees with a recent study done by Kenigsberg and Hodgen [17], but is in contrast with a study done by Balmaceda et al [19] in which bolus injections of GnRH 18 hours after administration of another GnRH antagonist elicited a blunted release of gonadotropins. A similar response was seen in adult male monkeys by other authors [18]. Importantly, we performed the GnRH challenge test 23 hours after GnRH antagonist administration. Thus, the pituitary may have had enough time to recover from the acute effects of the antagonist.

Because some recovery of pituitary FSH/LH secretion in response to the acute GnRH challenge test was seen post-treatment, we postulated that low doses of GnRH antagonist may suppress tonic gonadotropin secretion, while the synthesis and storage of LH and FSH are replenished and sustained concurrently. This would allow for maintenance of the ready releasable pool of gonadotropins, as when doses of GnRH are given. Administration of high doses of GnRH antagonist may not only suppress the secretion of basal gonadotropins, but also block the synthesis and storage of gonadotropins in the anterior pituitary.

In summary, although all monkeys were completely suppressed by treatment day 6, pituitary response to GnRH was retained. After 3 days post-treatment, although tonic gonadotropins remained suppressed, pituitary release of FSH and particularly LH in response to a GnRH challenge test had increased when compared to the test administered during antagonist treatment. None of the monkeys showed either local or general evidence of histamine release side effects. We conclude that: (1) the GnRH antagonist used in this study is potent and without discernible side effects; (2) pretreatment levels of gonadotropins (particularly FSH) are predictive of the treatment time needed to achieve full pituitary suppression of tonic gonadotropin secretion; (3) normal pituitary response to a GnRH challenge test is retained for several days; and (4) recovery of pituitary secretory capacity in response to a GnRH bolus is manifest before restoration of tonic levels of serum gonadotropins.

This GnRH antagonist, as well as others without significant histamine release effects, deserve consideration for pursuit in clinical trials. The rapidity of action of antagonists of GnRH and their absence of interim stimulatory effects (such as agonists manifest) may allow greater practical utility for both therapeutic and diagnostic applications.

REFERENCES

1. Chillik, CF, Itskovitz, J, Hahn, DW, McGuire, JL, Danforth, DR and Hodgen, GD (1987). Characterizing pituitary response to a gonadotropin releasing hormone (GnRH) antagonist in monkeys: tonic follicle-stimulating hormone/luteinizing hormone secretion versus acute GnRH challenge tests before, during and after treatment. *Fertil Steril*, 48, 480
2. Yen, SCC (1983). Clinical applications of gonadotropin-releasing hormone and gonadotropin releasing hormone analogs. *Fertil Steril*, 39, 257
3. Hodgen, GD (1983). Releasing hormones as diagnostic and therapeutic agents. *Fertil Steril*, 39, 592
4. Comite, F, Cutler, GB, Rivier, J, Vale, W, Loriaux, DL and Crowley, WF (1981). Short-term treatment of idiopathic precocious puberty with a long-acting analogue of luteinizing-releasing hormone. *N Engl J Med*, 305, 1546
5. Luder, AS, Holland, FJ, Costigan, DC, Jenner, MR, Wielgosz, G and Fazekas, ATA (1984). Intranasal and subcutaneous treatment of central precocious puberty in both sexes with a long-acting analog of luteinizing hormone-releasing hormone. *J Clin Endocrinol Metab*, 58, 966
6. Lemay, A, Maheux, R, Faure, N, Jean, C and Fazekas, ATA (1984). Reversible hyponadism induced by a luteinizing hormone-releasing hormone (LH-RH) agonist (Buserelin) as a new therapeutic approach for endometriosis. *Fertil Steril*, 41, 862
7. Nillius, SJ, Gudmundsson, J and Bergquist, C (1984). A new superagonist of GnRH for inhibition of ovulation in women. *Ups J Med Sci*, 89, 147
8. Linde, R, Doelle, GC, Alexander, N, Kirchner, F, Vale, W, Rivier, J and Rabin, D (1981). Reversible inhibition of testicular steroidogenesis and spermatogenesis by a potent gonadotropin-releasing hormone agonist in normal men: an approach toward the development of a male contraceptive. *N Engl J Med*, 305, 663
9. Labrie, F, Dupont, A, Belanger, A, Lacourciere, Y, Raynaud, JP, Husson, JM, Gareau, J, Fazekas, ATA, Sandow, J, Monfette, G, Girard, JG, Emond, J and Houle, JG (1983). New approach in the treatment of prostate cancer: complete instead of only partial withdrawal of androgens. *Prostate*, 4, 579
10. Filicori, M, Hall, DA, Loughlin, JS, Rivier, J, Vale, W and Crowley, WF (1983). A conservative approach to the management of uterine leiomyoma: pituitary desensitization by a luteinizing hormone-releasing hormone analogue. *Am J Obstet Gynecol*, 147, 726
11. Kenigsberg, D, Littman, BA, Williams, RF and Hodgen, GD (1984). Medical hypophysectomy: II. Variability of ovarian response to gonadotropin therapy. *Fertil Steril*, 42, 116
12. Craft, IL, Porter, RN, Smith, B and Jacobs, HS (1985). Ovulation induction in the poor responder. *Fourth World Conf In Vitro Fert*, Melbourne, Australia (Abstr)

13. Wildt, L, Diedrich, K, Van der Ven, H, Al Hasani, S, Hubner, H and Klasen, R (1986). Ovarian hyperstimulation for in vitro fertilization controlled by GnRH agonist administered in combination with human menopausal gonadotropins. *Human Reprod*, 1, 15
14. Schally, AV, Arimura, A and Coy, DH (1980). Recent approaches to fertility control based on derivatives of LH-RH. *Vitam Horm*, 38, 257
15. Kenigsberg, D, Littman, BA and Hodgen, GD (1984). Medical hypophysectomy: I. Dose-response using a gonadotropin-releasing hormone antagonist. *Fertil Steril*, 42, 112
16. Norman, RL, Rivier, J, Vale, W and Spies, HG (1986). Inhibition of estradiol-induced gonadotropin-releasing hormone antagonist. *Fertil Steril*, 45, 288
17. Kenigsberg, D and Hodgen, GD (1986). Ovulation-inhibition by administration of weekly gonadotropin-releasing hormone antagonist. *J Clin Endocrinol Metab*, 62, 734
18. Bint Akhtar, F, Weinbauer, GF and Nieschlag, E (1985). Acute and chronic effects of a gonadotrophin-releasing hormone antagonist on pituitary and testicular function in monkeys. *J Endocrinol*, 104, 345
19. Balmaceda, JP, Coy, DH, Schally, AV and Asch, RH (1983). Temporal changes in FSH and LH concentrations following the administration of a potent LH-RH inhibitory analogue ($[N\text{-Ac-D-Trp}^1,^3, D\text{-p-Cl-Phe}^2, D\text{-Phe}^6, D\text{-Ala}^{10}]\text{-LH-RH}$) to oophorectomized rhesus monkeys. *Acta Eur Fertil*, 14, 249
20. Goodman, AL, Descalzi, CD, Johnson, DK and Hodgen, GD (1977). Composite pattern of circulating LH, FSH, estradiol and progesterone during the menstrual cycle in cynomolgus monkeys. *Proc Soc Exp Biol Med*, 155, 479
21. Rivier, JE, Porter, J, Rivier, CL, Perrin, M, Corrigan, A, Hook, WA, Siraganian, RP and Vale, WW. New effective gonadotropin releasing hormone antagonists with minimal potency for histamine release *In Vitro*. *J Med Chem*, in press
22. Dixon, WJ and Massey, F (eds.). "Introduction to statistical analysis". 1983. (New York: McGraw-Hill, Inc.)
23. Schriock, ED, Monroe, SE, Martin, MC, Henzl, MR and Jaffe, RB (1985). Effect on corpus luteum function of luteal phase administration of a potent gonadotropin-releasing hormone analog (nafarelin). *Fertil Steril*, 43, 844
24. Schmidt, F, Sundaram, K, Thau, RB and Bardin, CW (1984). $[Ac\text{-D-Nal}(2)^1, 4FD\text{-Phe}^2, D\text{-Trp}^3, D\text{-Arg}^6]\text{-LHRH}$, a potent antagonist of LHRH, produces transient edema and behavioral changes in rats. *Contraception*, 29, 283
25. Hahn, DW, McGuire, JL, Vale, WE and Rivier, J (1985). Reproductive/endocrine and anaphylactoid properties of an LHRH antagonist, ORF 18260 $[Ac\text{-D-Nal}(2)^1, 4FDPhe^2, D\text{-Trp}^3, D\text{-Arg}^6]\text{-GnRH}$. *Life Sci*, 37, 505
26. Cetel, NS, Rivier, J, Vale, W and Yen SSC (1983). The dynamics of gonadotropin inhibition in women induced by an antagonist analog of gonadotropin-releasing hormone. *J Clin Endocrinol Metab*, 57, 62

THE REPRODUCTIVE, ENDOCRINE AND PHARMACOLOGICAL EVALUATION OF A NEW LHRH ANTAGONIST

**Do Won HAHN, Audrey PHILLIPS, John L. McGUIRE,
David M. RITCHIE, Robert J. CAPETOLA, C. BOWERS
and Karl FOLKERS**

Ortho Research Laboratories, Ortho Pharmaceutical Corporation, Raritan
New Jersey, USA; Tulane University, New Orleans, Louisiana, USA; and
Institute for Biomedical Research, University of Texas, Austin, Texas, USA

INTRODUCTION

LHRH agonists suppress the reproductive endocrine system through desensitization and are potentially useful in the treatment of clinical disorders such as uterine fibroids, precocious puberty, endometriosis and prostatic carcinoma. The development of these peptides has progressed in recent years and their clinical efficacy and safety are established. A drawback to the use of these agonists, however, is related to the initial stimulation of the reproductive endocrine system which may exacerbate symptoms of the disease being treated. LHRH antagonists also suppress the reproductive endocrine system but do so by competitively inhibiting binding of LHRH to its receptors and so avoid the initial stimulation which occurs with agonists. Unfortunately, the development of LHRH antagonists was impeded when facial edema was observed in toxicity studies in rats being treated with one of the early antagonists, [N-Ac-D-Nal(2)¹, D-pF-Phe², D-Trp³, D-Arg⁶]LHRH (ORF 18260) [1]. Later this peptide was also shown to release histamine [2] and induce cutaneous anaphylaxis [3] which are thought to be the causes of the facial edema which had been observed. Similar reactions were subsequently observed with other LHRH antagonists. These findings raised serious doubt concerning the safety in clinical use of these LHRH antagonists.

Research efforts were then directed towards further modification of the peptide structure of the prototype LHRH antagonists in an attempt to separate the desirable LHRH-blocking property from the undesirable anaphylactoid-like activity. Over forty peptides were screened for anti-ovulatory and anaphylactoid activity in rats, and it was shown that these two activities were not linked [4-6]. These encouraging results suggested that an LHRH antagonist devoid of undesirable anaphylactoid activity could be synthesized. This effort led to the discovery of ORF 21243 ([N-Ac-D-Nal(2)¹, D-pCl-Phe², D-Pal(3)³, D-Glu(AA)⁶, D-Ala¹⁰]LHRH) as a potent LHRH antagonist with an improved therapeutic ratio relative to that of ORF 19260 [5,7]. Subsequently, a third generation peptide with an even more improved margin of safety was identified. This peptide is ORF

23541 ([N-Ac-D-Nal(2)¹, D-pCl-Phe², D-Pal(3)³, Lys(Nic)⁵, D-Lys(Nic)⁶, Lys(iPr)⁸, D-Ala¹⁰]LHRH). The evaluation of its LHRH-inhibiting and anaphylactoid activities are described in this chapter.

SUPPRESSION OF GONADOTROPIN RELEASE

ORF 23541 was characterized as an LHRH antagonist in three test systems. First, the ability to inhibit LHRH-induced LH release was examined in dispersed male rat pituitary cells in vitro. ORF 23541 has an IC₅₀ of 2×10^{-10} M in this assay [8].

The antioviulatory activity of ORF 23541 was demonstrated in proestrous rats. The antagonist was administered in the afternoon of proestrus and on the following morning, the oviducts were examined microscopically for ova. ORF 23541 blocked ovulation, with an ED₅₀ of 1.45 μ g [8].

Finally, the LHRH antagonist activity of ORF 23541 was also evaluated by its ability to suppress gonadotropin levels in cannulated ovariectomized rats. Ovarian steroids provide negative feedback to the pituitary and removal of the ovaries results in an elevation of gonadotropins. ORF 23541, like other LHRH antagonists, blocks this elevation (Fig. 1). LH levels were




FIGURE 1 The suppression of (A) plasma LH and (B) plasma FSH levels in ovariectomized cannulated rats after a single s.c. injection of ORF 23541. (Reprinted with permission from Life Sci [8])

maximally suppressed within 3 hours after subcutaneous injection of 20, 200 or 2000 $\mu\text{g/kg}$ of ORF 23541 [8]. The recovery was dose-related with a return to castrate levels by 24 or 72 hours after injection with 20 or 200 $\mu\text{g/kg}$, respectively. LH levels remained suppressed at 96 hours after injection of 2000 $\mu\text{g/kg}$. FSH levels were similarly affected except that the onset of and recovery from the inhibiting activity of ORF 23541 were delayed as compared to changes in LH levels. The persistent suppression of LH levels 96 hours after injection of 2000 $\mu\text{g/kg}$ of ORF 23541 was different from historical data obtained for ORF 21243 [9]. This prompted us to re-evaluate the two antagonists side by side. The results of this experiment confirmed the long-acting nature of ORF 23541 (Fig. 2). A single injection of 2000 $\mu\text{g/kg}$ of either




FIGURE 2 The suppression of plasma LH levels in ovariectomized cannulated rats after a single s.c. injection of 2000 $\mu\text{g/kg}$ of either ORF 23541 or ORF 21243. (Reprinted with permission from Life Sci [8])

peptide maximally suppressed LH levels within 3 hours [8]. A return to castrate levels occurred at 72 hours after injection of ORF 21243, whereas LH remained suppressed at 96 hours after injection of ORF 23541. Similar long-lasting effects of ORF 23541 were observed in castrated monkeys [10].

CUNTANEOUS ANAPHYLACTOID-LIKE ACTIVITY

The facial edema in rats resulting from administration of ORF 18260 has been characterized by reference to vascular permeability

changes. Those studies suggested that the facial reactions were identical to a cutaneous anaphylactoid-like reaction, possibly as a result of the direct action of the LHRH antagonist on mast cell mediator release. We have used the induction of cutaneous vascular permeability changes (wheal response) in rats and guinea pigs as a predictor of the risk of causing such a reaction clinically in man [11]. In this model, Evan's blue dye is injected intravenously into the tail vein. The dye binds to serum proteins. Increases in vascular permeability can then be visualized as serum proteins, carrying the dye, diffuse into the interstitium and stain connective tissue. Immediately following dye injections, various dilutions of GnRH antagonists were injected intradermally into a shaved section on the back of the animal. Fifteen minutes after the intradermal injection, the animals were killed, the dorsal skin reflected, and the area of wheal was estimated using the mean of the longest perpendicular diameters. Regression analysis was used to estimate the dose producing a 8.75 x 8.75 mm wheal.

Relatively high doses of ORF 23541 were required to cause a wheal response: 10.9 and 13.7 µg in rats and guinea pigs, respectively (Table 1). In contrast, much lower doses of ORF 21243 and ORF 18260 caused the same response. The therapeutic

Table 1. Cutaneous Anaphylactoid-Like Activity (Wheal Response) and Therapeutic Ratio of LHRH Antagonists in Rats and Guinea Pigs

Peptide	Species	ED 8.75 x 8.75 µg/wheal (95% F.L.)	Antioviulatory ED ₅₀ , µg (95% F.L.)	Therapeutic Ratio ^a
LHRH	Rat	ND ^b ---	---	
LHRH	Guinea Pig	ND ^b ---	---	
ORF 23541	Rat	10.9 (5.8-24.7)	1.45 (1.0-1.9)	7.52
ORF 23541	Guinea Pig	13.7 (7.5-32.3)	---	---
ORF 21243	Rat	0.48 (0.20-0.94)	0.50 (0.30-0.55)	0.96
ORF 21243	Guinea Pig	1.75 (1.15-2.82)	---	---
ORF 18260	Rat	0.01 (0.0004-0.05)	0.65 (0.46-0.85)	0.02
ORF 18260	Guinea Pig	0.77 (0.55-1.08)	---	---

^aTherapeutic ratio = µg (ED anaphylactoid activity)/µg (ED₅₀ anti-ovulatory activity).

^bThe slope of the line for LHRH was not significantly different from zero. An ED wheal was not determined with doses up to 10 µg.

ratio of ORF 23541 in rats (7.52) was much better than that of ORF 21243 (0.96) or ORF 18260 (0.02). These studies suggest an improved margin of safety of ORF 23541. The intradermal administration used in these studies is more provocative than the subcutaneous and intramuscular routes proposed for clinical use. Therefore, this calculated ratio may underestimate the clinical therapeutic index.

CARDIOPULMONARY EVALUATION

Mast cell degranulation with release of mediators including histamine may also affect pulmonary function as these mediators are potent bronchospastic agents. We have evaluated the cardiopulmonary effects of LHRH antagonists in anesthetized dogs and guinea pigs [11]. Guinea pigs were placed in a whole body plethysmograph. A jugular vein and a carotid artery were cannulated for compound administration and for monitoring of blood pressure, respectively. The trachea was also cannulated to allow ventilation at a constant volume via a miniature Starling pump. One differential pressure transducer was used to measure transpleural pressure and tidal volume was monitored by a second. The guinea pigs were treated with succinyl choline to arrest spontaneous respiration. Following a five-minute stabilization period one of the LHRH antagonists was administered i.v. in a volume of 1.0 ml over 30 seconds. In dogs, a femoral artery was cannulated for monitoring blood pressure and a femoral vein was cannulated for administering compounds. A tracheal cannula was inserted and connected to a pneumotachograph for the measurement of air flow. The other side of the pneumotachograph was attached to a respirator for ventilating the animal. An intrapleural cannula was surgically implanted in the 6th intercostal space to monitor intrapleural pressure. The LHRH antagonist was administered i.v. in a volume of 5.0 ml over 30 seconds. In both species, airway resistance and dynamic lung compliance were calculated by a pulmonary computer. Percent change from baseline was monitored over 30 minutes for all parameters and recorded at 1, 3, 5, and 30 minutes after administration.

ORF 23541 did not affect pulmonary function in guinea pigs (Figure 3) or dogs (Figure 4). In contrast ORF 21243 and ORF 18260 did affect pulmonary function. Bronchospasm was observed in dogs after 10mg of either ORF 18260 or ORF 21243, as evidenced by increase in airway resistance and an associated drop in compliance. A severe bronchospasm was observed with 10mg of ORF 21243 in guinea pigs. Relatively high intravenous doses (10mg) were required to observe this effect and a safety factor is demonstrable, i.e. lack of pulmonary effects at a dose of 1mg. However, the finding itself represents a potential risk with these two antagonists. These results suggest that the risk of pulmonary effects in the clinical use of ORF 23541 is very low and that this peptide has an improved safety margin compared to the other two antagonists.




FIGURE 3 Effects of LHRH antagonists on cardiopulmonary function in guinea pigs




FIGURE 4 Effects of LHRH antagonists on cardiopulmonary function in dogs

A slight decrease in arterial pressure was caused by 10mg of ORF 23541 in guinea pigs, but no such change was seen with this dose in the dog, or with lower doses in the guinea pig.

CONCLUSION

ORF 23541 represents a new generation of potent LHRH antagonists with potentially improved margin of safety. It is hoped that these encouraging findings abate the concern regarding the clinical safety of these peptides and stimulate renewed interest in the development of LHRH antagonists as therapeutic agents.

ACKNOWLEDGEMENTS

We thank C. Bishop, T. Kirchner and MT Lai for their skillful technical assistance and K. Romash and C. Maines for the preparation of this manuscript. ORF 23541 was obtained from the Contraceptive Development Branch of NICHD Bethesda, MD, US and ORF 21243 and ORF 18260 from J. Rivier of the Salk Institute, La Jolla, CA, USA.

REFERENCES

1. Schmidt, F, Sundaram K, Thau, RB and Bardin, CW (1984). [Ac-D-Nal(2)¹, 4FD-Phe², D-Trp³, D-Arg⁶]-LHRH, a potent antagonist of LHRH, produces transient edema and behavioral changes in rats. *Contraception*, 29, 293
2. Hook WA, Karten M and Siraganian, RP (1985). Histamine release by structural analogs of LHRH. *Fed. Proc.*, 44, 1323
3. Hahn, DW, McGuire, JL, Vale, WW and Rivier, J (1985). Reproductive/endocrine and anaphylactoid properties of an LHRH antagonist, ORF 18260 [Ac-DNal¹(2), 4F-OPhe², D-Trp³, D-Arg⁶] -GnRH. *Life Sci*, 37, 505
4. Phillips, A, Hahn, DW, Capetola, RJ, Bishop, C and McGuire, JL (1987). Anaphylactoid and antioviulatory activities of LHRH antagonists in rats. *Life Sci*, 41, 2017
5. Phillips, A, Hahn, DW, Capetola, RJ and McGuire, JL (1986). The antioviulatory and anaphylactoid activities of two potent LHRH antagonists. *The Pharmacologist*, 28, Abs #689
6. Rivier, JE, Porter, J, Rivier, CL, Perrin, M, Corrigan, A, Hook, WA, Siraganian, RP and Vale WW (1986). New effective gonadotropin releasing hormone antagonists with minimal potency for histamine release *in vitro*. *J Med Chem*, 29, 1846
7. Phillips, A, Hahn, DW, Bishop, C, Capetola, RJ and McGuire, JL (1987). Anaphylactoid properties of LHRH analogs. In: Vickery BH and Nestor Jr JJ (eds.) "LHRH and Its Analogs, Part 2". p.191. (Boston: MTP Press)
8. Phillips, A, Hahn, DW, McGuire, JL and Folkers, K (1988). Pharmacologic properties of a new LHRH antagonist. *Life Sci* (in press)

9. Phillips, A, Klimek, S, Lai, MT, Hahn, DW and McGuire, JL (1986). Inhibition of ovulation and estrous cyclicity in rats with ORF 21243 - A potent LHRH antagonist. Am Fert Soc, Toronto. Abs #41
10. Chillik, CF, Itskovitz, J, Hahn, DW, McGuire, JL, Danforth, DR, and Hodgen GD (1989). Second generation GnRH antagonist: characterization of pituitary response in monkeys. This volume
11. Phillips, A, Hahn, DW, Capetola, RJ, Ritchie, D, McGuire, JL and Folders, K (1988). Evaluation of the anaphylactoid activity of a new LHRH antagonist (ORF 25341). J Med Chem (in press)

PHARMACOKINETICS OF LHRH ANALOGUES

J. SANDOW, G. JERABEK-SANDOW and B. KRAUSS

Hoechst AG, Pharmacology H 821, D-6230 Frankfurt 80, FRG

INTRODUCTION

Peptides are particularly well suited for therapy, because their metabolism differs markedly from conventional drugs. Biotransformation to inactive metabolites and amino acids is rapid, the capacity of peptidase enzymes is high, and enzyme induction has not been observed. LHRH agonists have an important therapeutic role in the control of gonadal steroid secretion in reproductive disorders, and in oncological indications (prostatic carcinoma and premenopausal mammary carcinoma). Their actions on gonadotropins and gonadal steroids are highly selective, with no side effects on the secretion of other hormones or on other organ systems [1-3]. The investigation of peptide pharmacokinetics and metabolism has made rapid progress due to the advances in analytical methodology. In this review, results on buserelin [D-Ser(tBu)⁶,Pro⁹NH₂Et]LHRH and other LHRH agonists are summarized and discussed with reference to the clinical implications for suitable dose regimens in different disorders.

ORGAN DISTRIBUTION

The organ distribution of isotope-labelled LHRH (gonadorelin) and the LHRH agonist leuprolide ([D-Leu⁶,Pro⁹NH₂Et]LHRH), nafarelin ([D-Nal(2)⁶]LHRH), and buserelin has been investigated in experimental animals. Accumulation of drug is found in the liver, kidney, anterior pituitary and intestine. In studies with ¹²⁵I-buserelin in rats, the liver and kidney are the main inactivating organs [4]. High tissue:plasma ratios are reached in these tissues with a maximum about 1 h after i.v. injection [5]. In terms of the percent dose, the liver and kidneys accumulate most of the radioactive drug and metabolites. Radiolabelled metabolites are also excreted in bile. A small fraction of the dose accumulates in the anterior pituitary gland with a time course proportional to the stimulation of LH release after a single dose. The amount of radiolabel excreted in the urine is proportional to the plasma concentration. The urinary peptides can be identified mainly as metabolites [6]. Reports on the organ

distribution of 125-I-buserelin [5], and 3-H-LHRH or 14-C-nafarelin [8] reach similar conclusions on the organs which show accumulation. When 125-I labelled buserelin is administered to the rat, there is a time-dependent increase of 125-I in the thyroid gland, due to the uptake of iodine resulting from metabolic inactivation and deiodination of tyrosine (the site of labelling). No intact peptide is found in the thyroid tissue. The fraction of intact peptide circulating in the serum or plasma decreases with time after injection, thus half-life estimates




FIGURE 1 Organ distribution study of 125-I-buserelin in male rats. Separation of intact buserelin and metabolites by HPCL. Upper panel: buserelin reference compound, lower panel: plasma sample obtained 1 h after treatment with 2 uCi/1.5ng 125-I-buserelin i.v., extracted by C18-silica (SEP-PAK) and separated on NOVA-PAK C18-column in a reversed phase system.

based on radioactive drug equivalents in the serum or plasma lead to erroneous half-life estimates. We found that in rats treated with 125-I-buserelin metabolism is rapid. The plasma values obtained 1 h after injection consist mainly of metabolites (Fig. 1), when analyzed by thin layer chromatography (TLC) or by high performance liquid chromatography (HPLC). In the HPLC separated plasma extract, 12.6% intact buserelin is identified. The circulating 125-I is not retained by C8-silica extraction (SEP-PAK cartridges). The fraction of intact buserelin in plasma can be increased by saturating the inactivating enzymes by a high dose of an LHRH antagonist administered prior to 125-I-buserelin [24].

With 14-C-nafarelin and 3-H-LHRH, it has also been observed that the fraction of intact peptide in the plasma decreases rapidly after treatment, whereas the total radioactivity circulating in the plasma remains high [7]. Significant differences exist with regard to protein binding. The hydrophobic agonist, nafarelin, is about 80% bound to plasma albumin [9], whereas leuprolide and buserelin have a protein binding of 15-20%, similar to LHRH [8].

METABOLISM

The biotransformation of buserelin to inactive metabolites has been investigated in rats with 125-I-buserelin, isolating the radiolabelled metabolites from liver, kidney, urine and plasma and comparing them with the unlabelled metabolites found in plasma, bile and urine of rats. Results of different studies using identification by TLC or HPLC are in good agreement. One hour after treatment, the livers of rats contain no intact buserelin, but a spectrum of smaller C-terminal metabolites (Fig. 2). In the kidneys, intact buserelin and several C-terminal metabolites are present, one of which is the buserelin [5-9] pentapeptide (Figure 3). This metabolite is also the major product found in the urine. It has also been found in the urine of all species investigated, including the human [2]. Intact buserelin and a variable proportion of the buserelin(2-9)octapeptide are also found in the pituitary gland [4].

Inactivation by enzymes

In contrast to LHRH, which is inactivated by several exo- and endopeptidases, many of the LHRH agonists are relatively resistant to enzyme degradation [5, 9, 10]. Incubation of buserelin with pyroglutamyl-aminopeptidase gives the (2-9)octapeptide from cleavage of the N-terminal bond between pyroglutamic acid and histidine. Both intact buserelin and C-terminal metabolites, e.g. the (2-9)octapeptide can be cleaved by chymotrypsin-like enzymes acting on the Trp³-Ser⁴ and Tyr⁵-D-Ser(tBu)⁶ bonds [11]. In the nonapeptide-ethylamide agonists, inactivation by the post-proline cleaving enzyme is blocked, whereas this enzyme can act on the decapeptide agonists like nafarelin and tryptorelin ([D-Trp⁶]LHRH). The azaGly-substitution in goserelin ([D-Ser(tBu)⁶,azGly¹⁰]LHRH) should also afford protection. Substitution by D-amino acids in position 6 does not completely inhibit the inactivation by chymotrypsin, because of limited stereoselectivity. The buserelin(5-9)pentapeptide is gradually converted to the (6-9)tetrapeptide, and the nafarelin(5-10) hexapeptide can be converted to the nafarelin (6-10)pentapeptide by chymotrypsin [9]. It is difficult to assess the relative contributions of chemical substitutions to enhanced receptor affinity or to enzyme resistance. In the hydrophobic decapeptide agonists an unprotected Pro⁹-glycinamide terminus is compatible

with high potency, suggesting that such agonists may be inaccessible to receptor-associated enzymes after binding.




FIGURE 2 Organ distribution study of ^{125}I -buserelin in male rats. Extraction of peptides from liver homogenate (supernatant fraction) by C18-silica (SEP-PAK) and separation on a NOVA-PAK C18-column in a reversed phase system (28). Intact buserelin indicated by shaded area.

Biliary excretion

From studies with labelled LHRH agonists, a significant fraction of the dose is excreted with the bile and found in the intestine, [7, 9]. After unlabelled buserelin, the only metabolite excreted in the bile of rats was identified as the buserelin (5-9)penta-peptide [12]. A bile:serum ratio of 140 exists for the (5-9) pentapeptide 120 min after s.c. injection of buserelin, indicating that an active secretory mechanism concentrates the (5-9)penta-peptide in the bile. With 14-C-nafarelin, the biliary metabolites identified in the rat [9] were the (5-10)hexapeptide, and smaller amounts of the [5-7]tripeptide, the (1-7)heptapeptide, and the (1-9)nonapeptide. Biliary excretion of buserelin is less than 1% in the dog, indicating that renal elimination may be more important in this species.

Urinary excretion

Immunoreactive LHRH is excreted in the urine of rats and human in small quantities that require concentration by extraction for detection [13]. After administration of equal doses of LHRH and




FIGURE 3 Organ distribution study of ^{125}I -buserelin in male rats. Extraction of peptides from liver homogenate (in 0.1-M acetic acid, supernatant fraction) by C18-silica (SEP-PAK) and separation on a NOVA-PAK C18-column in a reversed phase system (28). The peak in fraction 9 corresponds to the buserelin (5-9) pentapeptide, intact buserelin is found in fraction 14 (shaded area).

buserelin in rats, the fraction of the dose of LHRH in the urine was 0.002%, whereas 25-30% of the buserelin dose was recovered [6]. The daily injection or infusion of 60 μg buserelin per rat resulted in a consistent urinary excretion over 14 days. For an analytical identification of the urinary peptide material, we developed a separation by high performance liquid chromatography (HPLC), with specific buserelin radioimmunoassay (RIA) as the detection method [14].

PHARMACOKINETICS

Early studies with synthetic LHRH showed that after an initial rapid distribution phase, this decapeptide has an elimination half-life of 10-12 min [15]. In contrast, highly active LHRH agonists such as leuprolide, nafarelin, goserelin and buserelin are eliminated much more slowly [8,16-19]. The peptides can be measured in the circulation by specific radioimmunoassays [8, 11, 17, 19-21], which detect the two groups of the peptide agonist with a C-terminal Pro⁹-glycinamide, and nonapeptide agonists with a C-terminal Pro⁹-ethylamide. The results of such measurements should be controlled by chromatography of serum samples to separate intact peptide and biologically inactive metabolites [15]. We have found that extraction on C18-silica cartridges, HPLC for separation, and RIA with specific buserelin




FIGURE 4 Pharmacokinetics of unlabelled buserelin in the human. Patients with endometriosis, injection of buserelin 500 μ g i.v. Serum samples taken 10, 120 and 360 min after injection. Intact buserelin and metabolites in serum extracts separated by HPLC (28) and measured by specific buserelin RIA. Intact buserelin and the main metabolite, the (5-9) pentapeptide, marked by arrows.

antiserum is a highly selective and sensitive combination detection method. In 6 women treated with an i.v. injection of 500 μ g of buserelin, the elimination half-life was 51 min. The concentrations of total immunoreactive buserelin measured in unextracted sera were 101.4 ± 33 ng/ml 20 min after injection and 1.12 ± 0.12 ng/ml 360 min after injection. In serum extracts analyzed by reversed-phase HPLC and RIA (Figure 4), intact buserelin was the main component. The main serum metabolite was the inactive buserelin(5-9)pentapeptide. In the urine collected 0-1 h after treatment, 66% intact buserelin and 28% of the buserelin(5-9)pentapeptide were found (Figure 5). In the urine collected between 6-24 h, the ratio of intact buserelin to (5-9)pentapeptide remained similar. The urinary buserelin/creatinine ratio 0-1 h after treatment was 345 ± 156 μ g/g creatinine, and 6-24 h after treatment the concentration was 4.7 μ g/g creatinine. These results confirmed that buserelin is eliminated more slowly than LHRH, and remains available to the pituitary receptors for a long period after injection of high doses. After s.c. injection of 1000 μ g, buserelin the half-life

estimate between 1 and 8 h after injection is 80 min. It is thus clear that high dose injections can saturate pituitary receptors and maintain down-regulation for a 24 h period, as illustrated by the clinical efficacy of single daily high dose injections of leuprolide in prostatic carcinoma.




FIGURE 5 Urinary excretion of buserelin and metabolites in patients with endometriosis after treatment with 500 μ g i.v. Urine samples collected between 0 and 1 h after treatment, and between 6 and 24 h. Intact buserelin and metabolites separated by HPLC (28) and measured by specific buserelin RIA. The ratio of intact buserelin and the main metabolite does not change with time after injection.

Treatment by nasal spray

The urinary excretion of immunoreactive buserelin after treatment with nasal spray is dose-related. Nasal spray medication is a significant improvement for long-term therapy with peptides. The absorption of buserelin in humans is 2.5 to 3.3% of the nasal spray dose [23]. Two bioequivalent formulations of buserelin have been investigated for their LH-releasing activity in comparison with s.c. injections. Single doses of 300 or 400 μ g nasally give comparable LH-release to 10 μ g s.c. The serum concentrations after nasal spray of buserelin and nafarelin have been monitored [11, 20, 21, 24]. Peak concentrations of immunoreactive buserelin are reached within 30-60 min. The time course of LH-release induced by the spray is similar to that after an s.c. injection

of buserelin. In the urine, maximum concentrations of immunoreactive buserelin are reached during the first 2 h after nasal treatment, with a time-dependent decline. The effective drug concentration during long-term treatment can be monitored by the daily cumulative excretion, collecting the urine for 24 h during the administration of consecutive doses (Table 1), or by collection for an 8-h period

Table 1. Excretion of immunoreactive buserelin measured by RIA in unextracted urine samples.

Buserelin Dose µg/24 h	Urinary Buserelin µg/24 h
2 x 300 µg	3.1 ± 0.7 (12)
3 x 300 µg	3.8 ± 1.1 (19)
4 x 300 µg	7.0 ± 0.7 (23)
5 x 300 µg	10.3 ± 1.4 (16)
8 x 300 µg	26.0 ± 4.1 (36)

Mean and standard error

() No. of determinations

after one nasal spray dose. Due to the limited absorption, the urinary excretion after nasal spray treatment is lower than that reached after buserelin injections [23, 25]. In contraception, a single daily dose of buserelin nasal spray is administered to reduce pituitary responsiveness and prevent the preovulatory LH-surge. In other gynecological indications, 3-4 nasal spray doses of 300µg per day are required to suppress oestrogen secretion and induce a reversible amenorrhoea [6].

Treatment by sustained release

The initial studies on buserelin pharmacokinetics were performed with s.c. infusion via an external minipump. The serum concentrations and urinary excretion of buserelin were closely related to the rate of infusion (Table 2). Since LHRH and buserelin may be inactivated by enzymes in serum at room temperature, handling of the samples under careful refrigeration or the addition of bacitracin 10^{-3} M as an enzyme inhibitor is required to obtain reproducible and consistent results [26]. Buserelin in urine samples is stable when handled repeatedly during assay procedures; freezing and thawing did not affect the analytical result [23].

During the infusion studies, steady state conditions are reached after about 6 h, and the urinary excretion of immunoreactive buserelin (µg/g creatinine) remains constant. The monitoring of clinical studies is therefore considerably facilitated. Urine samples may be obtained at predetermined intervals without having to perform a quantitative collection of urine for 24 h. After treatment with buserelin implants, the

Table 2. Buserelin measured by RIA in unextracted serum and urine samples.

Buserelin Dose μg/24 h	Serum Buserelin ng/ml	Urinary Buserelin μg/g Creatinine
25 μg		1.11 ± 0.55 (3)
50 μg	0.33 ± 0.14 (6)	16.43 ± 2.06 (6)
100 μg	0.38 ± 0.08 (14)	25.51 ± 6.51 (11)
200 μg	0.53 ± 0.05 (45)	34.42 ± 5.95 (20)
300 μg	0.95 ± 0.29 (27)	43.24 ± 12.34 (9)

Mean and standard error
() No. of determinations

urinary buserelin excretion (μg/g creatinine) reaches a plateau on day 2-3 after treatment, with a gradual subsequent decline over a period of 4-12 weeks, depending on the dose and formulation [3, 14]. In patients with prostatic carcinoma, buserelin implants containing 6.6 mg peptide in a matrix of polylactic/glycolic acid (75:25) produced urinary excretion rates of 10-15μg buserelin/g creatinine, indicating a daily release rate of 50-75μg buserelin when corrected for an average creatinine excretion of 1.5 g/ 24 h, and a fractional excretion of 30% of the dose in the urine. In several studies, the serum concentrations during implant treatment with a dose of 3.3 mg or 6.6 mg were closely correlated with the urinary excretion rates (Fig 6). It is convenient to monitor the release rate by the buserelin/creatinine ratio in daily morning urine samples.

The spectrum of metabolites identified in the urine of the mouse, rat, dog, monkey and human was similar [2], but different amounts of intact buserelin were present. The fraction of intact buserelin in the urine was well correlated with the dose requirement for suppression for gonadal steroid secretion in each species (Table 3). The mouse, being very insensitive to suppression requires an extremely high daily dose of 25 mg/kg s.c., and almost no intact buserelin is recovered from the urine, the peptide material consisting mainly of the (5-9)pentapeptide (Figure 7). In the rat, which also has a high dose requirement, 23% of intact buserelin was excreted. In the dog, which is readily suppressed by buserelin, the highest fraction of intact buserelin (70%) was found in the urine. In the monkey and human, the fraction of intact buserelin was 45-55%, and the buserelin (59)pentapeptide was the main urinary metabolite as in the other species.

In the human during treatment by buserelin injections infusions, or by biodegradable implants the fraction of the dose of immunoreactive buserelin was 16-30%, depending on the experimental conditions (Figure 8). Of this peptide material, about 50-55% was identified as intact buserelin. The excretion of




FIGURE 6 Release of busserelin in patients with prostate carcinoma treated with implants containing a dose of 3.3 or 6.6mg peptide in a matrix of polylactide/glycolide 75:25. The release is determined by specific busserelin RIA in unextracted serum samples, and in morning urine samples with a correction for creatinine excretion (28). The busserelin concentrations in serum and urine are well-correlated and dose-dependent.

Table 3. Metabolism of busserelin in different species: dose-requirement for gonadal steroid suppression* and rate of metabolism.

Species	Steroid Suppression	Dose Per Day	% Intact Busserelin in Urine
mouse	testosterone	25 mg/kg	N.D.
rat	testosterone	0.1 mg/kg	22.5
dog	testosterone	2.5 µg/kg	70.2
monkey	oestradiol	10 µg/kg	54.2
human	oestradiol	4 µg/kg	55.1

N.D. nondetectable

*Mouse and rat exhibit partial suppression at the dose indicated, dog, monkey and human are fully suppressed.

busserelin in the milk of nursing mothers was investigated in studies on post-partum contraception [27]. The highest

concentration was 8.8 ng/ml, an a maximum amount of 1-2/ μ g was calculated as the dose likely to be ingested by infants. There was no charge in urinary LH excretion in these nursing infants, and oral administration of 600 μ g busserelin had no effect on LH secretion in adult volunteers. It is concluded that busserelin does not enter breast milk in biologically effective concentrations.




FIGURE 7 Busserelin in urine of mice and rats treated with busserelin injections of 25mg/kg s.c. in mice 0.12mg/kg s.c. in rats. The release is determined by specific busserelin RIA in unextracted urine samples after separation of intact busserelin and C-terminal metabolites by HPCL (28).




FIGURE 8 Separation of the intact peptide and all C-terminal metabolites excreted in the urine of a patient with endometriosis treated with busserelin injections (35 μ g/kg s.c. per day). A 25 h urine sample was analyzed by HPLC/RIA [28]. The metabolites identified were as indicated.

REFERENCES

1. Sandow, J (1987). Pharmacology of LHRH agonists. In: Furr, BJA and Wakeling, AE (eds.) "Pharmacology and Clinical Uses of Inhibitors of Hormone Secretion and Action". p.365 (London: Baillier Tindal)
2. Sandow, J and von Rechenberg, W (1987). Pharmacokinetics, Metabolism and clinical studies with buserelin. In: Vickery, BH and Nestor, JJ Jr. (eds.) "LHRH and Its Analogs, Contraceptive and Therapeutic Applications, Part 2". p.363. (Lancaster: MTP Press Ltd.)
3. Sandow, J, Fraser, HM and Geithövel, F (1986). Pharmacology and experimental basis of therapy with LHRH agonists in women. In: Rolland R, Chadha, DR and Willemsen WNP (eds.) "Gonadotropin-Down-Regulation in Gynecological Practice", Progress in Clinical and Biological Research. vol. 255, p.1. (New York: Alan Liss, Inc.)
4. Sandow, J, Jerabek-Sandow, G, Krauss, B and Stoll, W (1982). Metabolic and dispositional studies with LHRH analogs. In: Zatuchni, GI, Shelton, JD and Sciarra, JJ (eds.) "LHRH Peptides as Female and Male Contraceptives", p.338. (Philadelphia: Harper & Row)
5. Sandow, J, and Clayton, RN (1983). The disposition, metabolism, kinetics and receptor binding properties of LHRH and its analogues. *Prog Horm Biochem Pharmacol*, 2, 63
6. Sandow, J, Seidel, HR, Krauss, B and Jerabek-Sandow, G (1987). Pharmacokinetics of LHRH agonists in different delivery systems and the relation to endocrine function. In: Klijn, JGM, Paridaens, R and Foekens, JA (eds.) "Hormonal Manipulation of Cancer: Peptides, Growth Factors, and New(Anti) Steroidal Agents" p.203. (New York: Raven Press)
7. Chu, NI, Chan, RL, Hama, KM, and Chaplin, MD (1985). Disposition of nafarelin acetate, a potent agonist of luteinizing hormone-releasing hormone, in rats and rhesus monkeys. *Drug Metab Dispos*, 13, 560
8. Clayton, RN, Bailey, LC, Cottam, J, Arkell, D, Perren, TJ, and Blackledge, GRP (1985). A radioimmunoassay for GnRH agonist analogue in serum of patients with prostate cancer treated with D-Ser(tBu)⁶Aza Gly¹⁰GnRH. *Clin Endocrinol*, 22, 453
9. Chan, RL, and Nerenberg, C (1987). Pharmacokinetics and metabolism of LHRH analogs. In: Vickery BH, Nestor, JJ Jr (eds.) LHRH and its Analogs: Contraceptive and Therapeutic Applications, Part 2". p.577. (Lancaster: MTP Press Ltd.)
10. Horsthemke, B, Knisatschek, H, Rivier, J, Sandow, J and Bauer, K (1981). Degradation of luteinizing hormone-releasing hormone and analogs by adenohipophyseal peptidases. *Biochem Biophys Res Commun*, 100, 753
11. Holland, FJ, Fishman, L, Costigan, DC, Luna, L and Leeder, S (1986). Pharmacokinetic characteristics of the gonadotropin-releasing hormone analog D-Ser(TBU)⁶-EA-10 luteinizing hormone-releasing hormone (buserelin) after subcutaneous and intranasal administration in children with central precocious puberty. *J Clin Endocrinol Metab*, 63, 1065

12. Sandow, J, von Rechenberg, W, Krauss, B and Jerabek-Sandow, G (1987). Excretion of a gonadorelin-agonist (buserelin) in the urine and bile of rats. *Naunyn-Schmiedeberg's Arch Pharmacol*, 335 (Suppl.), 23
13. Bourguignon, JP, Hoyoux, C, Reuter, A, Franchimont, P, Leinartz-Dourcy, C, Vrindts-Gevaert, Y (1979). Urinary excretion of immunoreactive luteinizing hormone-like material and gonadotropins at different stages of life. *J Clin Endocrinol Metab*, 48, 78
14. Sandow, J, Jerabek-Sandow, G, and Schmidt-Gollwitzer, M (1985). Metabolism and pharmacokinetics of LHRH agonists. In: Schmidt-Gollwitzer, M and Schley, R (eds.) "LHRH and its Analogues, Fertility and Antifertility Aspects". p.105. (Berlin: Walter de Gruyter)
15. Handelsman, DJ, and Swerdloff, RS (1986). Pharmacokinetics of gonadotropin-releasing hormone and its analogs. *Endocr Rev* 7, 95
16. Barron, JL, Millar, RP and Di Searle (1982). Metabolic clearance and plasma half-disappearance time of D-Trp⁶ and exogenous luteinizing hormone-releasing hormone. *J Clin Endocrinol Metab*, 54, 1169
17. Nerenberg, C, Foreman, J, Chu, N, Chaplin, MD, and Kushinsky, S (1984). Radioimmunoassay of nafarelin [(6-(3-(2-naphthyl)-D-alanine)-luteinizing hormone-releasing hormone in plasma or serum. *Anal Biochem*, 141, 10
18. Sennello LT, Finley RA, Chu SY, Jagst, C, Max, D, Rollins, DE, and Tolmanm KG (1986). Single-dose pharmacokinetics of leuprolide in humans following intravenous and subcutaneous administration. *J Pharm Sci*, 75, 158
19. Yamazaki, I, and Okada, H (1980). A radioimmunoassay for a highly active luteinizing hormone-releasing hormone analogue and relation between the serum level of the analogue and that of gonadotropin. *Endocrinol Japon*, 27, 593
20. Reznik, Y, Winiger, BP, Aubert, ML, and Sizonenko, PC (1987). Pharmacodynamics of (D-Ser(t-Bu)⁶,Des-Gly¹⁰)GnRH ethylamide (Buserelin). *Acta Endocrinol (Copenh)*, 115, 235
21. Saito, S, Saito, H, Yamasaki, R, Hosoi, E, Komatsu, M, Iwahana, H, and Maeda, T (1985). Radioimmunoassay of an analog of luteinizing hormone-releasing hormone, (D-Ser(tBu)⁶des-Gly-NH¹⁰ ethylamide (Buserelin). *J Immunol Methods*, 79, 173
22. Kiesel, L, Sandow, J, Bertges, K, Jerabek-Sandow, G, Trabant, H and Runnebaum, B (1987). Serum concentration and urinary excretion of the LHRH agonist buserelin in patients with endometriosis. VI World Congress on Human Reproduction, Tokyo, p.133, Abstract
23. Sandow, J, Fraser, HM, Seidel, H, Krauss, B, Jerabek-Sandow, G and von Rechenberg, W (1987). Buserelin: pharmacokinetics, metabolism and mechanisms of action. *Brit J Clin Pract*, 41, 6
24. Anik, ST, McRae, G, Nerenberg, C, Worden, A, Foreman, J, Hwang, JY, Kushinsky, S, Jones RE, Vickery, B (1984). Nasal absorption of nafarelin acetate, the decapeptide (D-Nal(2)⁶) LHRH, in rhesus monkeys I. *J Pharm Sci*, 73, 684

25. Sandow, J, Jerabek-Sandow, G, Krauss, B and von Rechenberg, W (1985). Pharmacokinetic and antigenicity studies with LHRH analogues. In: Runnebaum, B, Rabe, T and Kiesel, L (eds.) "Future Aspects in Contraception, Part 2, Female Contraception" p.129. (Boston: MTP Press Ltd.)
26. Fraser, HM, Yorkston, CE, Sandow, J, Seidel, H and von Rechenberg, W (1987). The biological evaluation of LHRH agonist (buserelin) implants in the female stump-tailed macaque. Brit J Clin Pract, 41, 14
27. Dewart, PJ, McNeilly, AS, Smith, SK, Sandow, J, Hillier, SG and Fraser, HM (1989). LRH agonist buserelin as a post-partum contraceptive: lack of biological activity of buserelin in breast milk. Acta Endocrinol (Copenh), 114, 185

MOLECULAR MECHANISMS OF GnRH ACTION ON MAMMARY TUMORS AND UTERINE LEIOMYOMATA

**Joseph LEVY, Tzvia SEGAL, Arnon WIZNITZER, Vaclav INSLER
and Yoav SHARONI**

Endocrinology Laboratory, Soroka Medical Center and
Division of Obstetrics and Gynecology, Clinical Biochemistry Unit,
Faculty of Health Sciences, Ben-Gurion University of the Negev,
Beer-Sheva, 84101 Israel

INTRODUCTION

Carcinoma of the breast is the most common and fatal form of cancer among women in western countries. Many of these tumors are hormone-dependent and are known to be affected by various hormones, such as estrogens, progesterone, corticosteroids, prolactin and insulin [1]. In recent years an additional battery of growth factors which may be released by the tumor cells to act as autocrine growth regulators has been added to the above mentioned list of hormones. The synthesis and secretion of these factors are frequently regulated by the classical hormones, especially estradiol [2]. The autocrine mechanism for the regulation of tumor cell growth, as suggested by Sporn and Todaro [3], is an attractive explanation for the unregulated growth of cancer cells.

The uterine leiomyoma is the most common benign tumor of the uterus and female pelvis. This tumor consists mainly of smooth muscle cells, as well as varying amounts of fibrous connective tissue. The pathogenesis of the myoma is unknown, but estrogens appear to play a role in its growth. The tumor thrives during the years of greatest ovarian activity. However, with the onset of menopause, the decline of ovarian estrogen secretion usually causes the cessation/regression of leiomyoma growth.

The understanding of the basic mechanisms responsible for the growth regulation of mammary tumors and uterine leiomyomata may provide the means for developing therapeutic drugs which would interfere with these mechanisms.

GnRH IN THE TREATMENT OF BREAST CANCER

GnRH analogs have recently been used in several clinical trials for treating breast cancer patients [4]. The rationale for this is illustrated in Fig. 1. According to its known mechanism of action, GnRH, when secreted in a pulsatile fashion from the hypothalamus, stimulates gonadotropin release from the pituitary. However, continuous administration of the peptide causes an




FIGURE 1 Direct and indirect effect of GnRH on mammary tumor growth. GnRH effect on mammary tumors can be explained by the involvement of two mechanisms: A. Indirect effect: Non-pulsatile GnRH administration performs "medical castration" by inhibiting pituitary gonadotropin release, and a decline in ovarian estrogen production. B. Direct interaction of GnRH with the tumor

inhibition of gonadotropin release, as shown by A, which in turn causes inhibition of gonadal steroidogenesis. It was surmised that the low estrogen level was the main reason for the inhibition of mammary tumor growth by GnRH. In this review we will summarize evidence which indicates a direct effect of GnRH analogs on mammary tumors, as illustrated by arrow B. We will discuss primarily the molecular mechanism of the direct action of GnRH in breast cancer, as no direct actions of GnRH on uterine leiomyomata have been reported.

For performing these studies two experimental models were used: (i) the hormone-independent human cell lines in culture - MDA-MB-231, and (ii) the DMBA (7,12-dimethylbenz[a]anthracene)-induced mammary tumor in the rat. The latter serves as an excellent model for the human disease, as its growth rate is

easily manipulated by hormones. Estrogens [5] and prolactin [6] stimulate its growth, while ovariectomy [5], antiestrogens [7] and drugs which decrease plasma prolactin level cause growth arrest or regression of the tumors [8]. We therefore suggested that hormonal transmembrane transducing mechanisms such as tyrosine kinase activity [8] and the phosphatidyl inositide (PI) cycle [9] may be involved in the growth of these tumors.

SPECIFIC BINDING OF GnRH ANALOGS IN EXTRAPITUITARY TISSUES

Our first aim was to search for possible specific binding sites of GnRH on the DMBA-induced mammary tumor and on uterine leiomyomata membranes. As described in Table 1, specific binding sites for GnRH in different extrapituitary tissues were reported by us and other groups. (For review, see Sandow [18]). The binding sites are characterized by their high specificity but low affinity for various GnRH analogs.

Table 1. Presence of specific binding sites for GnRH in various tissues

Tissue	Reference
Pituitary	10
Breast tumor (biopsy)	11
Breast cancer Human cell lines	12
DMBA-induced mammary tumor in rat	13
Leiomyoma	14
Placenta	15
Testis	16
Ovarian carcinoma	17

GnRH TRANSDUCING MECHANISM

By analogy to the known mechanism of GnRH action in the pituitary [19], we deduce that GnRH may act on mammary tumor membranes via the Phospholipase-C (PL-C) transducing system. As shown in Fig. 2, this transmembranal system includes a receptor which activates the PI specific PL-C effector enzyme, which in turn produces two second messengers - diacyl glycerol and inositol triphosphate.

We developed [20] an in vitro system for measuring phospholipase-C activity in isolated mammary tumor membranes, as can be seen in Fig. 3.

The effect of GnRH on mammary tumor membranes was examined using the GnRH analog [D-Trp⁶]GnRH. As shown in Fig. 4, this analog stimulates PL-C activity in a dose-dependent manner. We examined the hormonal effect with two more GnRH analogs (Fig. 5).




FIGURE 2 Molecular mechanisms for the direct action of GnRH on mammary tumor membranes. GnRH acts on mammary tumor membranes via the Phospholipase-C (PL-C) transducing system. This transmembranal system includes a receptor (R) which activates the phosphoinositide-specific PL-C effector enzyme, which in turn produces two second messengers - diacyl glycerol (DG) and inositol tris phosphate (IP₃) from phosphatidyl inositol diphosphate (PIP₂). IP₃ releases Ca²⁺ from the endoplasmic reticulum (ER), the cation activates protein kinase-C (PK-C) together with DG. G-binding proteins (G) are modulators of this transducing mechanism as shown in experiments with GTP_γS, an unhydrolyzable analog of GTP.

The stimulatory effect of buserelin [D-Ser(Bu^t)⁶, Pro⁹ NHet]GnRH on PL-C activity is greater than that of [D-Trp⁶]GnRH. The effect of 10⁻⁷ M [D-Trp⁶]GnRH was completely suppressed in the presence of 10⁻⁴ M of the GnRH antagonist ORG-30276 ([N-Ac-D-pCl-Phe¹, D-pCl-Phe², D-Trp³, D-Arg⁶, D-Ala¹⁰]GnRH).

The binding of GnRH and its stimulation of PL-C activity in mammary tumor membranes may trigger the mechanism involved in the direct effect of the decapeptide upon mammary tumor growth.




FIGURE 3 Phospholipase-C assay protocol. Isolated membranes from different tissues are incubated in the presence of labelled PIP₂. The reaction is stopped with an acidified chloroform methanol mixture. Separation to two phases is achieved by the addition of water. The upper aqueous phase is applied to Dowex 1+8 columns, and the inositol phosphates are separated by three sequential washes with increasing concentrations of ammonium formate.




FIGURE 4 Stimulation of membranal PL-C activity by [D-Trp⁶] GnRH. IP₃ produced in control incubations (without hormone) was subtracted from all experiments.

THE DIRECT EFFECT OF GnRH ON MAMMARY TUMOR CELL PROLIFERATION

Would these basic findings indicate that GnRH may be used in the therapy of breast cancer? The answer is not simple, as is evident from the low rate of response in several clinical studies reported so far [4,21]. One of the explanations for the partial success might be that insufficient emphasis was placed on the direct effect of GnRH analogs on the mammary tumor. This direct effect may be involved in the responses noted in clinical studies performed on postmenopausal breast cancer patients [21]. We have therefore studied the direct effect of buserelin and ORG 30276 on human mammary tumor cell line proliferation.




FIGURE 5 Effect of GnRH agonist and antagonist on membranal PL-C activity. The effect of ORG 30276 (10^{-4} M) shown in the insert was assayed in the presence of 10^{-7} M [D-Trp⁶]GnRH.

As depicted in Fig. 6, buserelin did not affect thymidine incorporation into the cells. The GnRH antagonist, however, significantly inhibited this incorporation. These results suggest that GnRH antagonists may directly affect mammary tumor cell proliferation. The fact that the GnRH antagonist also inhibits PL-C activity as described above may be relevant to the effect on cell proliferation. However, more studies are needed to clarify this point.




FIGURE 6 Effect of buserelin and ORG 30276 on the proliferation of MDA-MB-231 breast cancer cells in culture. Cells were plated on 96 plating flasks. After 24 hours the cells were incubated either in the presence or the absence of the agonist or antagonist for 24 additional hours before being incubated with labelled thymidine for 3 hours.

FUTURE PROSPECTS

One probably cannot expect to find the same inhibitory effect on mammary tumor growth by GnRH agonist and antagonist as found in the pituitary system in gonadotropin release studies. The receptor(s) for GnRH analogs in mammary tumors may recognize different peptides from those developed for the pituitary system. Consequently, for more effective mammary tumor treatment, the analogs should be screened in assays that will emphasize their direct effect on mammary tumors.

The first screening stage can be performed in the tube (in vitro) by measuring GnRH analog binding and modulation of PL-C activity. Membranes from experimental tumor models or from patients' biopsies will be used. The selected analogs will be tested for their effect on the proliferation of mammary tumor cell lines in culture (second stage). In the third stage, tests on animal models, such as the DMBA-induced mammary tumor in the rat, will be performed before the final clinical trials.

This procedure, which is obviously more economical, can be employed in the selection of the optimal GnRH analog, from among the many available, for the treatment of breast cancer patients.

ACKNOWLEDGEMENTS

This work was supported in part by the Chief Scientist's Office, Israel Ministry of Health, and by the Beatty Malkin Grant for cancer research. The authors wish to thank Mrs. M. Duman for her excellent editorial assistance.

We are grateful to Profs. Sandow and Rechenberg (Hoechst AG, Frankfurt, FRG) for supplying buserelin; to Profs. Ghraf and Schroder (Ferring GmbH, Kiel, FRG) for supplying [D-Trp⁶]GnRH; and to Dr. L. Tax (Organon, Oss, The Netherlands) for supplying ORG-30376.

REFERENCES

1. Sharoni, Y and Levy, J (1985). The involvement of phosphoinositides phosphorylation and tyrosine kinase activity in the growth of mammary tumors. *Rev Basic Clin Pharmacol*, 5, 57s
2. Dickson, RB and Lippman, ME (1987). Estrogenic regulation of growth and polypeptide growth factor secretion in human breast carcinoma. *Endocr Rev*, 8, 29
3. Sporn, MB and Todaro, GJ (1980). Autocrine secretion and malignant transformation of cells. *N Engl J Med*, 303, 878
4. Manni, A, Santen, R, Harvey, H, Lipton, A and Max, D (1986). Treatment of breast cancer with gonadotropin-releasing hormone. *Endocr Rev*, 7, 89
5. Levy, J, Liel, Y and Glick, SM (1981). Peroxidase activity as a marker for estrogenicity studies in uterus and mammary tumors. *Isr J Med Sci*, 17, 970
6. Johnson, ML, Levy, J and Rosen, JM (1983). Isolation and characterization of casein producing and non-producing cell populations from 7, 12 dimethylbenz(a)anthracene-induced rat mammary carcinomas. *Cancer Res* 43, 2199
7. Levy, J, Liel, Y, Feldman, B, Aflallo, L and Glick, SM (1981). Peroxidase activity in mammary tumors: effect of tamoxifen. *Eur J Cancer Clin Oncol*, 17, 1023
8. Sharoni, Y, Radian, S and Levy, J (1985). Membranal tyrosine protein kinase activity (but not cAMP dependent protein kinase activity) is associated with growth of rat mammary tumors. *FEBS Lett*, 189, 133
9. Sharoni, Y, Teuerstein, I and Levy, J (1986). Phosphoinositide phosphorylation precedes growth in rat mammary tumors. *Biochem Biophys Res Commun*, 134, 876
10. Hazum, E, Schwartz, I, Waksman, Y and Keinan, D (1986). Solubilization and purification of rat pituitary gonadotropin-releasing hormone receptor. *J Biol Chem*, 261, 13043
11. Eidne, KA, Flanagan, CA and Millar, RP (1985). Gonadotropin-releasing hormone binding sites in human breast carcinoma. *Science*, 239, 989
12. Eidne, KA, Flanagan CA, Harris, NS and Millar, RP (1987). Gonadotropin-releasing hormone (GnRH)-binding sites in human breast cancer cell lines and inhibitory effects of GnRH antagonists. *J Clin Endocrinol Metab*, 64, 425

13. Segal, Z, Feldman, B, Levy, J and Sharoni, Y (1988). The role of PK-C in the negative regulation of GnRH dependent PL-C activity in mammary tumor membranes. *Gynecol Endocrinol*, 2 (Suppl. 1), 136, abstract
14. Wiznitzer, A, Marbach, M, Sharoni, Y, Insler, V and Levy, J (1988). Gonadotropin-releasing hormone specific binding site in uterine leiomyomata. *Gynecol Endocrinol*, 2 (Suppl. 1), 184, abstract
15. Iwashita, M, Evans, MI and Catt, KJ (1986). Characterization of a gonadotropin-releasing hormone receptor site in term placenta and chorionic villi. *J Clin Endocrinol Metab*, 62, 127
16. Clayton, RN, Katikineni, M, Chan, V, Dufau, ML and Catt, KJ (1980). Direct inhibition of testicular function by gonadotropin-releasing hormone: mediation by specific gonadotropin-releasing hormone receptors in interstitial cells. *Biochemistry*, 77, 4459
17. Emons, G (1988). The use of GnRH-analogues in ovarian cancer. *Gynecol Endocrinol*, 2 (Suppl. 1), 48, abstract
18. Sandow, J (1983). The regulation of LHRH action at the pituitary and gonadal receptor level: A review. *Psychoneuro-endocrinology*, 3, 277
19. Conn, PM, Huckle, WR, Andrews, WV and McArdle, CA (1987). The molecular mechanism of action of gonadotropin releasing hormone (GnRH) in the pituitary. *Rec Prog Horm Res*, 43, 29
20. Segal T, Levy, J and Sharoni, Y (1987). GnRH analogs stimulate phospholipase C activity in mammary tumor membranes: modulation by GTP. *Mol Cell Endocrinol*, 53, 239
21. Harvey, HA, Lipton, A, Santen, RJ, Escher, GC, Hardy, MN, Glode, IM, Segaloff, A, Landau, RZ, Schneir, H and Max, DT (1981). Phase II study of a gonadotropin-releasing hormone analogue (Leuprolide) in postmenopausal advanced breast cancer patients. *Proc Am Assoc Cancer Res/Am Soc Clin Oncol*, 22, 444

KINETICS OF SUSTAINED RELEASE DELIVERY SYSTEMS

R. DEGHENGI

Debiopharm S.A., 1003 Lausanne, Switzerland

INTRODUCTION

Depot formulations of contraceptive steroids, generally in the form of subcutaneous implants of injectable particles, have been known for more than twenty years, [1, 2] but the work on sustained release of hormonal peptides was started relatively recently. Polypeptides need special delivery systems to circumvent the general lack of oral activity and the short duration of action once they are released in the systemic circulation.

The discovery of the decapeptide LHRH, presented a particular therapeutic challenge for several reasons. Stimulation is obtained when LHRH is given at low doses and intermittently (or in pulsatile fashion), but the opposite (antagonistic) effect prevails when this peptide is administered continuously. Therefore, when optimal antagonistic effect is desired, to suppress tumor growth for instance, the LHRH agonist has to be administered with a specially designed sustained release delivery system.

A group under this author's direction [3] was probably the first, in 1974, to point out the usefulness of microencapsulation for intramuscular sustained delivery of LHRH. The first description of microencapsulation of an LHRH agonist (nafarelin) using dl lactide-coglycolide (PLGA) was in 1984 [4], this was subsequently refined and characterized [5-8]. The process was also applied to tryptorelin [9]. Researchers from ICI described the use of a solid, implantable PLGA formulation with goserelin [10, 11]. IPSEN-Biotech published a pharmacokinetic study on one such formulation with tryptorelin (D-Trp⁶-LHRH) as the active principle [12]. More information on controlled release formulations of LHRH agonists and antagonists has been presented at this Symposium.

KINETICS OF RELEASE

The release pattern of the LHRH analogue is a function of the physico-chemical properties of the peptide as well as of the polymer. It has been recognized that essentially two mechanisms of release operate in sequence: 1) diffusion of the active principle through the polymeric matrix of which the delivery systems are made and 2) hydrolytic dissolution of the polymer with time, which frees additional active principle. An overlapping phase may exist between the two mechanisms. As of today, two LHRH agonists (tryptorelin and goserelin) are commercially available in a depot, slow release formulation and at least three more (nafarelin, buserelin and leuprolide) are under active development.

From my analysis of the published data, bell-shaped patterns of release are obtained as measured by plasma levels of peptide. Two general patterns are thus obtained: a symmetrical curve as in Fig. 1b, with plasma levels increasing to a peak at several days after the injection and a non-symmetrical one (Fig. 1a), characterized by a "burst effect" resulting in a peak within day one of the injection with a secondary lower peak several days later. None of these patterns follows zero-order kinetics, a fact which, at least for LHRH agonists as opposed to antagonists, may have therapeutic significance.




FIGURE 1 Patterns of release of LHRH agonists from sustained release formulations

Sustained release patterns of nafarelin acetate [7] and goserelin [8] are similar to the one depicted in pattern "b" (Fig. 1). On the contrary, tryptorelin [12] leuprolide [14] and

buserelin [15] follow pattern "a" with the early peak due to the burst effect.

Whether it is by pattern a or b, all these formulations have achieved a sustained release of the active principle and have therefore fulfilled one of the requirements for optimal antagonistic activity. The question however should be asked, whether the different release patterns reflect any clinically appreciable difference which would allow a distinction and a preference between them.

Let us examine carefully the evidence:

- 1) It has been known for some time from kinetic studies and confirmed by recent experiments [16] that the LHRH induced desensitization of rat pituitary cells is dose and time dependent. In other words the gonadotrophin release from the pituitary can be down-regulated by continuous exposure to LHRH. Higher doses result in a more rapid down-regulation, whereas preincubation with lower doses requires a longer period of pretreatment with the releasing factor to achieve the same effect.
- 2) Animal experiments conducted in male dogs [17] have shown a positive correlation between the dose of tryptorelin released on day 1 and the rate of inhibition of testosterone.
- 3) Experiments in normal volunteers receiving controlled release injectable preparations loaded with different concentrations (2, 4, 7%) of nafarelin have shown [18] that serum testosterone levels were suppressed earlier with the higher load (7%) which also gave an earlier peak of the agonist.
- 4) In patients treated with varying doses of sustained release goserelin [11] it was noted that the lowest dose (0.9mg) resulted in the longest time (up to 44 days) to achieve castrate levels of testosterone, a fact which prompted cessation of randomisation and a transfer of the patients to a higher dose (3.6mg).

These facts lead us to postulate that high initial dose and continuous administration is preferable to a gradual increase in dose and continuous administration at least for those patients who are starting treatment and are therefore at some risk of flare-up episodes due to the initial hormonal stimulation.

The "burst-effect" of pattern "a" is then the equivalent of a loading dose frequently given therapeutically when a fast initial effect is desirable.

Obviously only a careful analysis of the decrease rates of testosterone, estradiol and gonadotrophins as related to the release kinetics of various preparations of LHRH agonists will confirm our postulate.

In summary, newer delivery systems of the kind discussed here not only ensure better patient compliance, but represent an optimization of therapeutic effects in terms of safety and efficacy.

ACKNOWLEDGEMENTS

I am grateful to Dr. Andrew V. Schally for helpful discussions and advice.

REFERENCES

1. Dzuik, PJ and Cook, B (1966). Passage of steroids through silicone rubber. *Endocrinology*, 78, 208
2. Beck, LR, Cowsar, DR, Lewis, DH, Gibson, JW and Flowers, CE (1979). New long-acting injectable microcapsule contraceptive system. *Am J Obstet Gynecol*, 135, 419
3. Immer, H, Nelson, VR and Goetz, M (1974). Process for preparing LHRH, salts and compositions therefore and intermediates therefore, US Pat 3,835,108, September 10, 1974
4. Kent, JS, Sanders, LM, McRae, GI, Vickery, BH, Tice, TR and Lewis, DH (1982). Feasibility studies on the controlled release of an LHRH analogue from subcutaneously injected polymeric microspheres: early in vivo studies. *Contr Del Sys*, 3, 58
5. Sanders, LM, Kent, JS, McRae, GI, Vickery, BH, Tice, TR and Lewis, DH (1984). Controlled release of an LHRH analogue from poly (D, L lactide-co-glycolide) microspheres. *J Pharm Sci*, 73, 1294
6. Kent, JS, Sanders, LM, Tice, TR and Lewis, DH (1984). Microencapsulation of the peptide nafarelin acetate for controlled release. In: Zatuchni GI, Goldsmith, A, Shelton, JD and Sciarra, JJ (eds.) 'Long-acting Contraceptive Delivery Systems'. p.169. (Philadelphia: Harper & Row)
7. Vickery, BH and Sanders, LM (1985). Studies on controlled release systems for LHRH analogues. In: Schmidt-Gollwitzer, M and Schley, R (eds.) "LHRH and Its Analogues: Fertility and Antifertility Aspects". p.123. (Berlin: Walter de Gruyter)
8. Sanders, LM, Vitale, KM, McRae, GI and Mishky, PB (1986). Controlled delivery of nafarelin, an agonistic analogue of LHRH, from microspheres of poly (D,L-lactic-co-glycolic) acid. In: Illum, L and Tomlison (eds.) "Delivery Systems for Peptide Drugs". p.125. (New York and London: Plenum Press)
9. Redding, TW, Schally, AV, Tice, TR and Meyers, WE (1984). Long-acting delivery systems for peptides: Inhibition of rat prostate tumors by controlled release of D-Trp⁶ LHRH from injectable microcapsules. *Proc Natl Acad Sci USA*, 81, 5845
10. Walker, KJ, Turkes, AO, Turkes, A, Zwink, R, Beacock, C, Buck, AC, Pelling WB and Griffith, K (1984). Treatment of patients with advanced cancer of the prostate using a slow release (depot) formulation of the LHRH agonist ICI 118630 (Zoladex). *J Endocrinol*, 103, R1
11. Hutchinson, GFG and Furr, BJA (1987). Biodegradable carriers for the sustained release of polypeptides. *TIBTECH* 5, 102
12. Ezan, E, Drieu, K, Chapelat, M, Rougeot, C and Dray, F (1986). Radioimmunoassay of (D-Trp⁶LHRH): its application to animal pharmacokinetic studies after single injection and long-acting formulation administration. *Reg Peptides*, 14, 155

13. Perren, TJ, Clayton, RN, Blackledge, G, Bailey, LC, Lynch, SS, Arkell, David, G, Cortam, J, Farrar, D, and Young, CH (1986). Pharmacokinetic and endocrinological parameters of a slow-release depot preparation of the GnRH analogue ICI 118630 (Zoladex) compared with a subcutaneous bolus and continuous infusion of the same drug in patients with prostatic cancer. *Cancer Chemother Pharmacol*, 18, 39
14. Shimamoto, T (1987). Nasal and Depot Formulations of Leuprolide. *J Androl*, 8, S-14
15. Fraser, HM, Sandow, J, Seidel, H and von Reichenberg, W (1987). An implant of a gonadotropin releasing hormone agonist (buserelin) which suppresses ovarian function in the macaque for 3-5 months. *Acta Endocrinol (Copenh)*, 115, 521
16. Helm, KJ, Kiesel, L, Rabe, T and Runnebaum, B (1987). Kinetics of GnRH induced desensitization of rat pituitary cells. *Acta Endocrinol (Copenh)*, 114, Suppl 283, 184 (Abstract)
17. Tonelli, PA, Lawter, JR, Nicolau, G, Silber, BM, Lanc, RA, Lanzilotti, M, Dente, AR, McWilliams, WE and Yacobi, A (1986). Decapeptyl: Release from microcapsules and testosterone suppression in the dog. *Pharm Res*, 3, suppl
18. Pavlou, SN, Hoffman, PG, Wakefield, GB, Le Page, ME, Kushinsky, S and Kovacs, WJ (1987). Sustained suppression of pituitary and testicular function by controlled release injectable nafarelin. Presented at the Satellite Symposium on medical management of Prostatic Cancer, Nice, France, Sept 1987

FORMULATIONS OF GnRH ANALOGUES FOR THERAPEUTIC USE

B.J.A. FURR and F.G. HUTCHINSON

Bioscience I and Pharmaceutical Departments, ICI Pharmaceuticals plc,
Alderley Park, Macclesfield, Cheshire, SK10 4TG, UK

INTRODUCTION

Formulation of drugs rarely elicits much interest amongst bioscientists and clinicians although it is well recognised that the properties of any formulation can have a considerable impact on the successful exploitation of a pharmaceutical agent. Polypeptides usually show low oral potency because their molecular weights are too high to allow effective absorption across the intestinal mucosa and they are rapidly broken down by proteolytic enzymes in gastrointestinal fluids. This is certainly true for luteinising hormone releasing hormone (LHRH) analogues: for example, 'Zoladex' ([D-Ser(tBu)⁶,azGly¹⁰])LHRH; goserelin; ICI 118,630), which is around 2000-fold less potent by the oral route than parenterally [1]. For this reason, the parenteral route has been preferred. This paper aims to review the formulations of LHRH analogues which have been used experimentally and clinically and to discuss their advantages and disadvantages.

AQUEOUS SOLUTIONS

Simple aqueous formulations are available for LHRH, buserelin ([D-Ser(tBu)⁶,Pro⁹NH₂Et]LHRH), leuprorelin (leuprolide; [D-Leu⁶,Pro⁹NH₂Et]LHRH), tryptorelin (decapeptyl; [D-Trp⁶]LHRH) and nafarelin ([D-Nal(2)⁶]LHRH). The drugs are usually administered intravenously, subcutaneously or intramuscularly. Aqueous formulations of LHRH and analogues are appropriate for acute administration, where the objective is to elicit gonadotrophin secretion; for example, in the assessment of pituitary function. Aqueous formulations are also currently preferred to use in the treatment of male and female infertility by pulsatile injection from programmed minipumps. However, data in farm animals [2] suggest that continuous infusion of very small doses of LHRH will stimulate ovulation during anoestrus. If these data are confirmed in patients with infertility then perhaps a constant infusion or depot preparation which releases tiny amounts of LHRH or analogues will also be of value.

To achieve pituitary desensitisation and the consequent castration-like effect needed for the treatment of malignant and benign sex hormone-responsive diseases, aqueous injections of LHRH analogues need to be administered one or more times daily.

Although there are reports of the successful treatment of prostate cancer using daily injections of leuprolide [3-4], such formulations are inconvenient to administer and are likely to give compliance problems if self-administered by the patient. For this reason alternative delivery systems and routes of administration have been widely sought.

BUCCAL, RECTAL, INTRAVAGINAL AND PERCUTANEOUS ADMINISTRATION

Apart from im, sc and iv injection, other routes of administration are possible for LHRH analogues; these include intravaginal [5-8], buccal [9] and rectal [10]. Generally these alternative routes of administration are associated with a very low and variable bioavailability and none offers wholly satisfactory solutions to the problems of peptide administration. In the authors' view oral, buccal and rectal administration, even with the use of absorption enhancers or surfactants, will always pose severe problems either in terms of bioavailability of drug or frequency of administration. None of these routes is compatible with genuine long term sustained release of peptide over weeks or months following a single administration.

The transdermal route is suited to the use of long-acting formulations and indeed it has been claimed that polypeptides can be administered percutaneously [11]. It seems unlikely, however, that large molecules such as polypeptides could ever be transported across unbroken and undamaged skin.

INTRANASAL DELIVERY

Apart from simple parenteral administration, the intranasal route has been the most widely studied. Buserelin is marketed as a nasal spray and a similar formulation of nafarelin is being developed [12]. Delivery of an aqueous formulation of an LHRH analogue into the nose is usually by means of a metered dose spray pump [13]. The absorption of LHRH analogues varies between 1 and 10% when delivered in this way [13] which makes accurate dosing difficult.

It has been claimed that the absorption of drug following nasal delivery may also be influenced by rhinitis, local pathological changes in the nasal cavity, and colds [13]. However recent data on intranasal dosing of buserelin shows that experimentally-induced rhinitis has no effect on drug absorption [14]. Although effective and widely used, nasal spray formulations give a number of problems in addition to poor and variable absorption of the drug. The need for multiple daily drug administration inevitably impairs compliance with the treatment

and may, indeed, be the reason why nasal spray therapy with buserelin was less effective than daily parenteral injection [15].

The presence of a preservative in most nasal formulations is one of the most frequent causes of nasal irritations and allergic rhinitis in patients on long term nasal spray therapy according to Harris [13].

For these reasons, we have preferred the parenteral route of administration which ensures essentially complete bioavailability of compound and have taken the view that the most practical and effective formulation for the administration LHRH analogues is a depot, either as a solid sub-dermal formulation or an injectable suspension, that provides for continuous, sustained release over many weeks or even months.

DEPOT FORMULATIONS

Depot formulations of 'Zoladex', tryptorelin, buserelin and nafarelin have been described [16-19], all of which appear to be based on dispersion of the drugs in co-polymers of lactic and glycolic acids.

These polymers have been selected because long experience shows that they are biocompatible and inert in the physiological environment and degrade to toxicologically acceptable fragments [20]. 'Zoladex' and buserelin are in the form of solid depots for sc injection. Tryptorelin and nafarelin have been studied as powdered solids which have to be dispersed in a solution before administration. The administration at no more than monthly intervals means that medical or paramedical staff can give the injections which should ensure improved compliance.

BIOASSAYS OF RELEASE

Since we are most familiar with the formulation of 'Zoladex' the remainder of this paper will describe the development and properties of 'Zoladex' depots. Continuous release of 'Zoladex' in vivo was measured qualitatively by the biological effect elicited in adult female rats showing regular oestrous cycles. Normally, these rats have an oestrous cycle of 4 days and the occurrence of oestrus is indicated by the presence of cornified cells in vaginal smears. In rats given sub-dermal depots of 'Zoladex', release of drug at an effective rate causes a fall in circulating oestrogens, which in turn leads to a suppression of oestrus and absence of cornified vaginal smears. Rats, therefore, show an extended period of dioestrus.

For amorphous homo- and co-polymers of approximately the same high molecular weight, increasing lactide concentration results in slower degradation. This is reflected in the biological effect elicited in rats using sub-dermal depots containing small amounts of drug in high molecular weight carriers (Fig. 1).

When administered to female rats some drug is released initially as judged by biological response. There then follows a period during which drug is either not released at all or is released at an ineffective rate and so the rats return to oestrus. At some later time point release recommences, the biological response returns and continues until the depot is fully depleted of drug. For polymers of similar molecular weight and




FIGURE 1 Effects of treatment with 100 μ g 'Zoladex' using subcutaneous depots containing 3% w/w drug on oestrous cycles in regularly cycling female rats

distribution it can be seen that the interval between the two phases of release is shortest for the most rapidly degradable polymer. The mechanism of release has been extensively reviewed [20].

When these two phases of release do not overlap discontinuous release is observed (Fig. 2A). However, by controlling the properties of the polymer, the initial phase of release can be

made to overlap with the second phase and depots can be defined which give continuous release over not less than 28 days (Fig. 2B).

Depot formulations, which are administered by subcutaneous injection and which release over at least 28 days, have been developed for clinical use. The depots are based on a poly (d,l-lactide-co-glycolide) matrix in which 'Zoladex' (3.6 mg) is uniformly dispersed. The depot is in the form of a cylindrical rod 11 x 1.1 mm.




FIGURE 2 Effect of subdermal depots containing 300µg Zoladex on estrous cycling in rats. Depots containing (A) 3% w/w drug in high molecular weight polymer, (B) 20% w/w drug in low molecular weight polymer. I, initial release due to surface leaching; II, degradation induced release

Although the primary objective was to produce a formulation of the drug which was more convenient to administer and which would secure improved compliance, these depot formulations also appear to have improved efficacy. This was demonstrated in rats which were given a single subcutaneous bolus dose of 50 µg 'Zoladex' after either pretreatment with daily subcutaneous injections of saline or 50 µg 'Zoladex' for 6 weeks or, alternatively, at the start of the experiment and at 4 weeks, a single subcutaneous depot, calculated to release a dose of 'Zoladex' equivalent to 50 µg daily. Rats were killed at various times after the final treatment and serum LH was determined by radioimmunoassay in blood collected from the dorsal aorta. The results are shown in Fig. 3.

As expected, the bolus dose of 'Zoladex' elicited a massive secretion of LH in the rats pre-treated with saline. In spite of pre-treatment for 6 weeks with 50 μ g 'Zoladex' daily the bolus dose of the drug still caused a substantial release of LH, although a clear degree of desensitisation of the pituitary gland occurred. In contrast, there was negligible LH secretion in response to the bolus injection in rats pre-treated with the 'Zoladex' depot, indicating an improved level of pituitary desensitisation with this formulation.

Studies in the pigtailed monkey have also shown that the sc administration of a depot containing 'Zoladex' is more effective in suppressing serum testosterone than daily sc injections of the drug [21]. Recently, two groups have shown that a depot




FIGURE 3 Serum LH release in response to a bolus dose of 50 μ g 'Zoladex' in rats pretreated with saline (C) or 50 μ g 'Zoladex' (I) daily for 6 weeks. The response after treatment at the start of the experiment and at 4 weeks with single subcutaneous depots, calculated to release 50 μ g 'Zoladex' daily, is also shown. Serum LH was measured by radioimmunoassay and the values given are the means of 5 samples at each time point

containing 3.6 mg is more effective at suppressing serum testosterone in patients with prostate cancer than daily sc

injections of 250 or 500 μg 'Zoladex' in an aqueous formulation [22,23].

BREAST CANCER

The efficacy of this depot formulation of 'Zoladex' was tested in two sex hormone-responsive tumor models. The first was the dimethyl-benzanthracene (DMBA)-induced rat mammary carcinoma [24] which is known to be dependent on both oestrogen and prolactin [25]. A single sc depot containing 500 μg 'Zoladex' caused an inhibition of oestrogen secretion, the disappearance of cornified cells from vaginal smears and regression of DMBA-induced mammary tumours.

Half of the tumours present at the start of the experiment were not palpable at 28 days but all save one of them reappeared between 40 and 60 days as the single 'Zoladex' depot became exhausted. In contrast, the DMBA-induced mammary tumours increased in size by more than 50% in control animals given a placebo depot.

If single depots of the drug were given sc at weeks 0, 4 and 8 of the study, the regression was more impressive. No tumour was palpable at week 11.

Again, by week 16 regrowth of the tumours occurred because treatment stopped at week 8. By week 20, the tumours had re-attained pre-treatment size.

When given at days 30, 58 and 86 after administration of the carcinogen, single depots containing 500 μg 'Zoladex' delayed the appearance of tumours for a period of around 100 days. That is, the expected duration of action of such a treatment regimen (Fig. 4).

When given at 28 day intervals, starting on day 30 after administration of the carcinogen, single sc depots of 'Zoladex' caused a more profound inhibition of tumour appearance and only 9 out of 21 rats had mammary tumours at the end of the study on day 450. Those tumours which were found did not regress following ovariectomy and so were classified as non-hormone responsive. It is concluded from this study, which approximates the clinical adjuvant therapy setting, that 'Zoladex' may be of benefit as a treatment for primary breast cancer post-mastectomy in pre-menopausal women.

The second tumour we have used is the Dunning R3327H rat transplantable prostate adenocarcinoma, which is androgen responsive and has been used extensively as a model for the human disease [26].

Single subcutaneous depots containing 1 mg 'Zoladex' given every 28 days to rats bearing Dunning R3327H prostate tumours implanted on each flank caused a marked inhibition of tumour growth indistinguishable from that in surgically castrated rats (Fig. 5). Twenty-one days after the eighth depot was given, the rats were killed and the weights of the sex organs and serum hormone concentrations measured by radioimmunoassay (Table 1).




FIGURE 4 Effect of depots containing 500 μ g 'Zoladex' on the appearance of mammary tumours induced in rats by DMBA. The number of rats with tumours in groups given placebo depots (Control), single depots at days 30, 58 and 86 (3 x Depot), or single depots every 28 days starting at day 30 ('N' x Depot) are shown. Each group comprised 21 rats




FIGURE 5 Effect of single subcutaneous depots containing 1mg 'Zoladex' on the growth of Dunning R3327H transplantable rat prostate tumours. The depots were given every 4 weeks as shown by the arrows. The control group (---) had 13 animals and 'Zoladex'-treated (—) and surgically castrated (···) groups 12 animals each as shown by the figures in parentheses

Table 1: Sex organ weights and serum hormone concentrations in "Zoladex"-treated, and surgically castrated rats bearing Dunning R3327H prostate tumours. Control values for rats of a similar age and weight are shown for comparison.

Group	Testes	Tissue Weight (mg)		Sera Hormone Levels (ng/ml)			
		Ventral Prostrate	Seminal Vesicle	LH	FSH	Prolactin	Testosterone
Zolad	367	21	54	<0.2	175	63	<0.3
Castrate	-	20	54	13	1413	61	<0.3
Intact	3500	250	350	2	400	30	3

Testes weights were about 10% those of control rats of a similar age and weight and showed atrophic histological changes. Ventral prostate gland and seminal vesicle weights were identical to those in the surgically castrated group and, histologically, were also completely atrophic.

Serum LH and testosterone were undetectable in the group given 'Zoladex' depot and serum FSH was decreased by 60-70%. Serum prolactin doubled in rats given 'Zoladex', as it did in surgically castrated animals, probably as a consequence of androgen withdrawal. This contrasts with the effect of 'Zoladex' in female rats where there was a significant reduction in serum prolactin following oestrogen withdrawal [27].

Because of interest in the concept of 'total androgen withdrawal' persuasively supported by Labrie and co-workers [28], a study of the effects of 'Zoladex' depot and a new antiandrogen, ICI 176,334 [29], given alone and in combination, were evaluated. The results (Fig. 6) clearly show that both 'Zoladex' depot and ICI 176,334 are effective at limiting tumour growth but that the combination offers no advantage over monotherapy with 'Zoladex'.

This finding is in agreement with studies [30,31] which showed that antiandrogen failed to improve the effect of surgical castration in rats with Dunning tumours. It is possible that the rat secretes less androgen from the adrenal gland than man and that the rat may, therefore, not be the most suitable species in which to test the hypothesis. The results of the extensive, controlled, comparative clinical studies now being undertaken will, hopefully, soon answer this question.

At present, asymptomatic patients with prostate cancer usually remain untreated probably because of the unacceptability of the most widely used therapy, castration. This policy should be reconsidered now that LHRH agonists and some newer antiandrogens have been introduced. It is important to know whether treatment early in the course of the disease reduces tumour growth rate and




FIGURE 6 Effect of single s.c. 1mg 'Zoladex' depots given every 28 days, daily oral administration of 25mg ICI 176,334/kg and the combination of both treatments on the growth of Dunning prostate tumours. The values shown are means \pm SEM; n=8

improves survival. Consequently, a study was made of the growth of Dunning tumours and the survival of rats given single 1 mg 'Zoladex' depots every 28 days starting either 28 days after tumour implantation, when most tumours are not palpable, or when tumours had reached a size of 2 cm². Tumour growth was measured at weekly intervals and survival was assessed. In this experiment rats were killed, for ethical reasons, on the week the tumour burden exceeded 20 cm². The results shown in Figs. 7 and 8 demonstrate that both treatment schedules reduce tumour growth rate and increase survival but that an early start of treatment produces the best result.

This finding is in accord with the work of Schally and Redding [32] who showed that combination of an LHRH agonist with cytotoxic chemotherapy produced more impressive responses and increases in survival if started early post-transplantation of Dunning tumours. These results argue strongly that consideration should now be given to the treatment of patients with early stages of prostate cancer. It is hoped that clinical trials will soon confirm the animal data.

The promising results achieved in animal studies with 'Zoladex' therapy have now been substantiated in clinical trials in men suffering from prostatic carcinoma and in premenopausal women with advanced breast cancer [33].




FIGURE 7 Effect of single depots containing 1mg 'Zoladex' given every 28 days on growth of Dunning prostate tumours in rats. Control rats were untreated; early 'Zoladex' treatment started 28 days after tumour implantation; late 'Zoladex' treatment started when tumours measured 2cm². There were 14 rats in the control group and 13 rats in each 'Zoladex' treated group




FIGURE 8 Effect of single depots containing 1mg 'Zoladex' given every 28 days on survival of rats with Dunning prostate tumours. Control rats were untreated; early 'Zoladex' treatment started 28 days after tumour implantation; late 'Zoladex' treatment started when tumours measured 2cm². There were 14 rats in the control group and 13 rats in each 'Zoladex' treated group

CONCLUSIONS

It is concluded that LHRH analogues can be effectively administered in a variety of formulations. Biocompatible, biodegradable depots of LHRH agonists are effective, well tolerated and convenient to administer and will find utility in the chronic treatment of sex hormone-responsive benign and malignant diseases.

ACKNOWLEDGEMENTS

NIAMDD, Professor L.E. Reichert and GD Niswender for reagents for FSH, LH and prolactin radioimmunoassays. The US National Prostate Cancer Agency for supplies of rats bearing Dunning R3327H tumours. The many colleagues in ICI Pharmaceuticals Research Department who have made this development possible, particularly A.S. Dutta, J.R. Churchill, J.R. Woodburn, B. Valcaccia and B. Curry.

REFERENCES

1. Furr, BJA (1987). Pharmacological studies with 'Zoladex'. A novel luteinising hormone-releasing hormone analogue. In: Chisholm GD (ed.) "'Zoladex' a new treatment for prostatic cancer". p. 1. (London: Royal Society of Medicine International Congress Series)
2. McLeod, BJ, Haresign, W and Lamming, GE (1983). Induction of ovulation in seasonally anoestrous ewes by continuous infusion of low doses of Gn-RH. *J Reprod Fert*, 68, 489
3. Garnick MB, Glode, LM and the Leuprolide Study Group (1984). Leuprolide versus diethylstilbestrol for metastatic prostate cancer. *N Engl J Med*, 311, 1281
4. Smith, JA for the Leuprolide Study Group (1985). Clinical effects of gonadotrophin-releasing hormone analogue in metastatic carcinoma of prostate. *Urology*, 25, 106
5. Okada, H, Yamazaki, I, Ogawa, Y, Hirai, S, Yashiki, T and Mima, H (1982). Vaginal absorption of a potent luteinizing hormone-releasing hormone analog (leuprolide) in rats I: absorption by various routes and absorption enhancement. *J Pharm Sci*, 71, 1367
6. Okada, H, Yamazaki, I, Yashiki, T and Mima, H (1983). Vaginal absorption of a potent luteinizing hormone-releasing hormone analog (leuprolide) in rats II: mechanism of absorption enhancement with organic acids. *J Pharm Sci*, 72, 75
7. Okada, H, Yashiki, T and Mimo, H (1983). Vaginal absorption of a potent luteinizing hormone analog (leuprolide) in rats III: effect of estrous cycle on vaginal absorption of hydrophilic model compounds. *J Pharm Sci*, 72, 173
8. Okada, H, Yamazaki, I, Yashiki, T, Shamamoto, T and Mimi, H (1984). Vaginal absorption of a potent luteinising hormone-releasing hormone analog (leuprolide) in rats IV:

evaluation of the vaginal absorption and gonadotrophin responses by radioimmunoassay. *J Pharm Sci*, 73, 298

9. Merkle, HP, Anders, R, Sandow, J and Schurr, W (1986). Drug delivery of peptides: the buccal route. In: Davis, SS, Illum, L and Tomlinson E (eds.) "Delivery Systems for Peptide Drugs, NATO ASI Series". p. 159. (New York: Plenum Press)
10. Muranishi, S, Takada, K, Hoshikawa, H and Murakami, M (1986). Enhanced absorption and lymphatic transport of macromolecules via the rectal route. In: Davis, SS, Illum, L and Tomlinson, E (eds.) "Delivery Systems for Peptide Drugs, NATO ASI Series". p. 177. (New York: Plenum Press)
11. Uda, K and Yamada, M (1984). Percutaneous pharmaceutical compositions for external use. European Patent Application 127426
12. Anik, ST, Sanders, LM, Chaplin, MD, Kushinski, S and Nerenberg, C (1984). Delivery systems of LHRH and analogues. In: Vickery, BH, Nestor JJ Jr. and Hafez, ESE (eds.) "LHRH and its Analogs: Contraceptive and Therapeutic Applications". p. 421. (Lancaster: MTP Press)
13. Harris, AS (1986). Biopharmaceutical aspects on the intranasal administration of peptides. In: Davis, SS, Illum, L and Tomlinson, (eds.) "Delivery Systems for Peptide Drugs. NATO ASI Series A: Life Science" vol.125, p. 191. (New York: Plenum Press)
14. Larsen, C, Jorgensen, MN, Tommerup, B, Mygind, N, Dagrose, EE, Grigoleit, HG and Malerczyk (1987). Influence of experimental rhinitis on the gonadotrophin response to intranasal administration of buserelin. *Eur J Clin Pharmacol*, 33, 155
15. Rajfer, J, Handelsman, DJ, Crum, A, Steiner, B, Peterson, M and Swerdloff, RS (1986). Comparison of the efficacy of subcutaneous and nasal spray buserelin treatment in suppression of testicular steroidogenesis in men with prostate cancer. *Fertil Steril*, 47, 104
16. Hutchinson, FG and Furr, BJA (1985). Biodegradable polymers for the sustained release of peptides. *Biochem Soc Trans*, 13, 520
17. Parmar, H, Lightman, SL, Allen, L, Phillips, RH, Edwards, L and Schally, AV (1985). Randomised controlled study of orchidectomy vs long-acting D-Trp-6-LHRH microcapsules in advanced prostatic carcinoma. *Lancet*, 11, 1201
18. Waxman, JH, Sandow, J, Man, A, Barnett, MJ, Hendry, WF, Besser, GM and Oliver, RTD (1986). The first clinical use of depot buserelin for advanced prostatic carcinoma. *Cancer Chemother Pharmacol*, 18, 174
19. Sanders, LM, Kent, JS, McRae, GI, Vickery, BH, Tice, TR and Lewis, DH (1984). Controlled release of a luteinizing hormone-releasing hormone analogue from poly (d,l-lactide-co-glycolide) microspheres. *J Pharm Sci*, 73, 1294
20. Hutchinson, FG and Furr, BJA (1987). Biodegradable carriers for the sustained release of polypeptides. *Trends in Biotechnology*, 5, 102
21. Furr, BJA and Hutchinson, FG (1985). Biodegradable sustained release formulation of the LH-RH analogue 'Zoladex' for the treatment of hormone-responsive tumours. " EORTC Genitourinary

- Group Monograph 2, Part A: Therapeutic Principles in Metastatic Prostatic Cancer", p. 143. (New York: Alan Liss)
22. Williams, G, Kerle, DJ, Roe, SM, Yeo, T and Bloom SR (1985). Results obtained in the treatment of prostate cancer patients with 'Zoladex'. "EORTC Genitourinary Group Monograph 2, Part A: Therapeutic Principles in Metastatic Prostatic Cancer". p. 287. (New York: Alan Liss)
23. Grant, JBF, Ahmed, SR, Shalet, SM, Costello, CB, Howell, A and Blacklock, NJ (1986). Testosterone and gonadotrophin profiles in patients on daily or monthly LHRH analogue ICI 118,630 ('Zoladex') compared with orchidectomy. Brit J Urol, 58, 539
24. Huggins, C, Brizziarelli, G and Sutton, H (1959). Induction of mammary carcinoma in the rat and influence of hormones on the tumours. J Exper Med, 104, 25
25. Jordan VC (1982). Laboratory models of hormone dependent cancer. Clin Oncol, 1, 21
26. Smolev, JK, Heston, WDW, Scott, WW and Coffey, DS (1977). Characterisation of the Dunning R3327H prostatic adenocarcinoma: an appropriate model for prostatic cancer. Cancer Treat Rev, 61, 273
27. Furr, BJA and Nicholson, RI (1982). Use of analogues of LHRH for treatment of cancer. J Reprod Fert, 64, 529
28. Labrie, F, Dupont, A, Belanger, A, Poyet, P, Giguere, M, Lacoursiere, Y, Edmond, J, Monfette, G and Borsanyi, JP (1986). Combined treatment with flutamide and surgical or medical (LHRH agonist) castration in metastatic prostate cancer. Lancet, 1, 48
29. Furr, BJA, Valcaccia, B, Curry, B, Woodburn, JR, Chesterson, G and Tucker, H (1987). ICI 176,334: a novel, non-steroidal, peripherally selective antiandrogen. J Endocrinol, 113, R7
30. Redding, TW and Schally, AV (1985). Investigation of the combination of the agonist D-Trp-6-LH-RH and the antiandrogen flutamide in the treatment of Dunning R-3327H Prostate Cancer Model. Prostate, 6, 219
31. Ellis, WJ and Isaacs, JT (1985). Effectiveness of complete versus partial androgen withdrawal therapy for the treatment of prostatic cancer as studied in the Dunning R-3327 system of rat prostatic adenocarcinomas. Cancer Res, 45, 6041
32. Schally, AV and Redding TW (1985). Combination of long-acting microcapsules of the D-tryptophan-6-analog of luteinizing hormone-releasing hormone with chemotherapy: Investigation in the rat prostate cancer model. Proc Natl Acad Sci USA, 82, 2498
33. Furr, BJA and Milsted, RAV (1987). Role of LHRH analogues in cancer treatment. Rev Endocrine-Related Cancer, 26, 5

INDEX

- adrenal glands, GnRH receptors 72, 75
- agonistic GnRH analogues 1, 6–14, 113–23
 - breast cancer 1, 12–13, 77–8, 127–33, 149, 150
 - clinical applications 2–3, 5–6, 8–14
 - formulations 143–54
 - gonadal protection against
 - chemotherapy and radiation 17
 - metabolism and excretion 115–17
 - organ distribution 113–15
 - ovarian cancer 10, 78
 - paradoxical antifertility effects 5, 7, 91
 - pharmacokinetics 117–23
 - prostate cancer 11–12, 78, 144, 149–53
 - reduced size (3–9) 47–9, 50–1
 - side-effects 3
 - structure–activity relationships 33–6
 - sustained release formulations, *see* sustained release formulations
 - see also specific agents*
- 17-alpha-hydroxylase 72
- 20-alpha-hydroxysteroid dehydrogenase 74
- anaphylactoid-like reactions 3, 100, 105
- newer GnRH antagonists 16, 100, 107–9
- androgen blockade, total 2, 151
- antagonistic GnRH analogues 1, 14–17
 - breast cancer 13, 77, 132–3
 - clinical trials 15–16
 - effects on breast cancer cell proliferation 132–3
 - first generation: ovulation inhibition 85–92
 - gonadal protection against
 - chemotherapy and radiation 17
 - potential clinical applications 6, 17, 92
 - reduced size (3–9) 47–9, 50–1
 - second generation: characterization of pituitary response 95–101
 - side-effects 3, 16, 39, 40, 100, 105
 - structure–activity relationships 15, 36–40
 - third generation: evaluation 105–11
- antiandrogens 2, 151
- aqueous solutions 143–4
- arachidonic acid 54, 55, 59, 61–3, 65, 66
- arachidonic acid metabolites 54, 61–5, 66
- aromatase 74
- arterial blood pressure 111
- alpha-aza-amino-acid substitution of C-terminal glycine residue 35
- azaglycinamide substitution at C-terminal glycine residue 38
- beta-glucuronidase 76
- biliary excretion, GnRH agonists 116
- blood pressure, arterial 111
- breast cancer 2, 6, 12–13, 127–33
 - direct effects of GnRH analogues 76–8, 132–3
 - GnRH agonists 1, 12–13, 77–8, 127–33, 149, 150
 - GnRH antagonists 13, 77, 132–3
 - GnRH receptors 72, 77, 129, 133
 - GnRH transducing mechanism 129–30
 - LHRH-like peptides 71, 76
- breast milk
 - buserelin in 122–3
 - LHRH in 76
- bronchospasm, GnRH antagonists inducing 109–11
- buccal administration 144
- burst effect release pattern 138, 139
- buserelin 11, 35, 36
 - active reduced size fragments 45
 - aqueous solution 143
 - breast cancer 13, 77, 132–3
 - in breast milk 122–3
 - intranasal 119–20, 144–5
 - metabolism 115, 116–17
 - organ distribution 113–14
 - pharmacokinetics 117–23
 - serum protein binding 115
 - stability 120
 - stimulation of phospholipase C activity 130
 - structure 6, 34
 - sustained release formulations 120–3, 138, 139, 145

- calcium, mediation of GnRH actions 53–4, 61, 65, 74, 130
- calmodulin 54
- central nervous system, LHRH-like peptides 71
- chemotherapy
 - combined with hormonal therapy 11, 13
 - GnRH analogues for gonadal protection 17
- chloroquine 55, 59, 60
- cholecystokinin 14
- choriocarcinoma 71
- chorionic gonadotropin, human (hCG) 76
- chymotrypsin 115
- contraception
 - GnRH agonists 5, 7, 8–9
 - GnRH antagonists 6, 85–92
- controlled release formulations, *see* sustained release formulations
- corpus luteum, GnRH receptors 72, 74–5
- cryptorchidism 8
- cyclic AMP (cAMP) 53, 74
- cyclic GMP (cGMP) 53, 59
- cyclo-oxygenase inhibitors 63
- cyclophosphamide 11
- [D-Trp⁶]-LHRH, *see* tryptorelin
- decapeptyl, *see* tryptorelin
- depot formulations, *see* sustained release formulations
- desensitization, pituitary, *see* pituitary desensitization
- 17,20-desmolase 72
- Detirelix (RS 68439) 15–16, 38
- diacylglycerol (DG) 59–60, 65, 66, 129, 130
- down-regulation of pituitary GnRH receptors 7, 65, 139
- dysfunctional bleeding 2
- endometriosis 2, 5, 9
- enzyme degradation, GnRH agonists 115–16
- epidermal growth factor (EGF) 12, 14
- epoxygenase arachidonic acid metabolites 63, 65
- 5,6-epoxycosatrienoic acid (5,6-EET) 63
- estrogen challenge test, effects of GnRH antagonists 86, 87, 89, 92
- estrogens
 - effects of GnRH antagonists 86–7, 88, 92
 - mammary tumour growth and 129
 - suppression by GnRH agonists 7
- ethylamide substitution of C-terminal glycine residue 35, 36–8
- fertirelin 34
- fibroids, uterine, *see* leiomyomas, uterine
- FSH receptors, gonadal 7
- FSH release, GnRH-stimulated, *see* GnRH challenge test
- FSH suppression
 - chronic Zoladex treatment 151
 - GnRH antagonists 15, 97–8, 100–1, 106–7
- G-binding proteins 66, 130
- gamete intra-fallopian transfer (GIFT) 6, 9–10
- gastrin 14
- gastrointestinal hormones 14
- glucosamine-6-phosphate synthase 75
- GnRH (LHRH)
 - aqueous solution 143
 - elimination half-life 117
 - extrapituitary actions 1, 71–92
 - inter-species variations 33, 34
 - intracellular mechanisms of action
 - mammary tumours 129–30
 - pituitary cells 53–66
 - organ distribution 113, 115
 - stability 120
 - structure 6, 34
- GnRH (LHRH) analogues
 - chemistry 33–41
 - clinical applications 2–3, 5–18
 - extrapituitary actions 71–92
 - formulations and delivery 1–2, 143–54
 - gonadal protection against
 - chemotherapy and radiation 17
 - reduced size 1, 45–51
 - safety 3
 - see also* agonistic GnRH analogues; antagonistic GnRH analogues
- GnRH (LHRH) challenge test
 - during GnRH antagonist treatment 15, 88, 90–1, 92, 99–100, 101
 - gonadotrope-enriched pituitary cell populations 56–7
 - superfused pituitary cells 55, 56
- GnRH (LHRH) receptors
 - affinity of reduced size GnRH analogues 51
 - binding assay 45–6
 - breast cancer 72, 77, 129, 133

- GnRH (LHRH) receptors (*continued*)
 - down-regulation 7, 65, 139
 - extrapituitary tissues 72, 73–4, 75, 76, 79, 129
 - tumours 72, 75, 76, 77, 78, 79, 129
- GnRH (LHRH)-like peptides 71, 79
 - breast cancer 76
 - gonads 73, 75
 - placenta 76
- gonadotropes
 - enrichment of cell fractions 56–7
 - intracellular mechanisms of GnRH action 53–4, 58–66
- goserelin (Zoladex) 11, 33–5, 36
 - metabolism 115
 - pharmacokinetics 117
 - structure 6, 34
 - sustained release formulations 137, 138, 139, 145–54
 - bioassays of release 145–9
 - in breast cancer 149, 150
 - in prostate cancer 149–53
- granulosa cells 74
- growth factors 12, 14
- growth hormone (GH) 12, 13
- GTP-binding proteins 66, 130
- half-lives, elimination 117–19
- hexapeptide GnRH analogues 1, 45–51
- histamine release 16, 39, 40, 100, 105
 - pulmonary function and 109–11
- histrelin 6, 34, 35
- HOE766 8
- 5-hydroperoxy-6,8,11,14-eicosatetraenoic acid (5-HPETE) 55
- 5-hydroxy-6,8,11,14-eicosatetraenoic acid (5-HETE) 55
- 15-hydroxy-5,8,10,14-eicosatetraenoic acid (15-HETE) 55
- hypogonadism, therapy of 8
- ICI 176,334 151, 152
- implantation, prevention or interception 8
- implants, *see* sustained release formulations
- in vitro fertilization/embryo transfer (IVF-ET) 2, 6, 9–10
- indole-(4-9) GnRH analogues 46, 47–8
- infertility treatment 2, 7, 9
- inositol phosphates
 - measurement of GnRH-induced release 54–5
 - mediation of GnRH actions 58–61, 65
- inositol 1,4,5-trisphosphate (IP₃) 59, 61, 65, 66, 129, 130, 131
- insulin-like growth factor 1 (IGF-1) 12
- insulinomas 71
- interception of implantation 8
- intestines, accumulation of GnRH agonists 113, 116
- intramuscular administration 143
- intranasal administration 1–2, 119–20, 144–5
- intravaginal administration 144
- intravenous administration 143
- kidneys
 - accumulation of GnRH agonists 113, 115
 - direct effects of GnRH 76
- leiomyomas, uterine 2, 5, 127
 - GnRH receptors 72, 75
 - GnRH therapy 10
- leukotrienes 55, 63–5, 66
- leuprolide (leuporelin) 13, 35, 36
 - active reduced size fragments 47
 - aqueous solution 143
 - organ distribution 113
 - pharmacokinetics 117
 - prostate cancer 11, 144
 - serum protein binding 115
 - structure 6, 34
 - sustained release formulations 138–9
- leuporelin, *see* leuprolide
- Leydig cells 72, 73
- LH receptors, gonadal 7
- LH release
 - assay for test GnRH analogues 46
 - GnRH-stimulated, *see* GnRH challenge test
 - intracellular mechanisms 53–66
 - superfused pituitary cells 55–6
 - Zoladex-induced 148
- LH suppression
 - GnRH agonists 7, 151
 - GnRH antagonists 15, 97–8, 100–1, 106–7
 - mid-cycle surge, by GnRH antagonists 15, 86–9, 92
- LHRH, *see* GnRH
- lipoygenase arachidonic acid metabolites 55, 63–4
- liver, accumulation of GnRH agonists 113, 115
- lutrelin 6, 34

- mammary tumours, see breast cancer
- melittin 55, 58–9, 60
- metabolism of GnRH agonists 115–17
- microcapsules 11, 137
- milk, see breast milk
- nafarelin 35
 - contraceptive effects 8
 - formulations 137, 143, 144, 145
 - metabolism 115, 116
 - organ distribution 113, 114, 115
 - pharmacokinetics 117, 119, 138–9
 - structure 6, 34
 - naphthyl (4–9) GnRH analogues 48–9
 - nasal administration 1–2, 119–20, 144–5
 - Novantrone 11
 - O-benzyl substitution of Ser⁴ side chain 49–50, 51
 - oocyte harvest 2, 10
 - oral administration 144
 - ORF 18260 38
 - anaphylactoid-like reactions 105, 107–9
 - pulmonary effects 109–10
 - ORF 21243 105, 107
 - anaphylactoid-like reactions 108–9
 - pulmonary effects 109–10
 - ORF 23541 105–11
 - cardiopulmonary effects 109–11
 - cutaneous anaphylactoid-like activity 107–9
 - suppression of gonadotropin release 106–7
 - ORG 30093 38
 - ORG 30276 14–15, 16, 38
 - breast cancer cell growth and 132–3
 - inhibition of phospholipase C activity 130, 132
 - ornithine decarboxylase 75, 76
 - osteoporosis 3, 8
 - osteosarcomas 2
 - ovarian cancer 2, 6, 10, 72, 78
 - ovarian disease, polycystic 2, 6, 9
 - ovarian hyperstimulation syndrome (OHSS) 9
 - ovary, direct actions of GnRH 71, 72, 74–5
 - ovulation
 - induction 8, 9, 10, 74
 - inhibition 8, 15, 85–92, 106
 - pancreas 71
 - pancreatic cancer 2, 13–14
 - percutaneous administration 144
 - pharmacokinetics 117–23, 138–9
 - phosphatidyl inositol diphosphate (PIP₂) 130, 131
 - phosphatidylinositol (PI) cycle
 - direct actions of GnRH in mammary tumours 129–30, 131
 - pituitary actions of GnRH 58–61, 65, 66
 - phosphoinositol-1,4,5-triphosphate (PIP₃) 59
 - phospholipase A₂ (PLA₂) 55, 58–9, 65, 66
 - phospholipase C (PL-C)
 - assay 131
 - effects of GnRH analogues 129–30, 131, 132, 133
 - stimulation of LH release 58
 - phospholipid turnover 54, 58–61, 74
 - pituitary cells
 - gonadotrope enrichment 56–7
 - LH secretion 46, 55–6
 - measurement of inositol phosphate release 54–5
 - post-receptor actions of GnRH 53–66
 - preparation 54
 - pituitary desensitization 7, 144
 - role of protein kinase C 65
 - sustained release formulations 139, 148
 - pituitary microsomal fractions, inositol phosphate binding 61
 - pituitary plasma membranes, GnRH receptor binding assay 45–6
 - placenta, direct actions of GnRH 71–2, 76
 - plasma protein binding 115
 - plasminogen activator 74
 - platelet-derived growth factor (PDGF) 12
 - poly (*d,l*-lactide-co-glycolide) (PLGA) 137, 147
 - poly(A)-polymerase 76
 - polycystic ovarian disease 2, 6, 9
 - precocious puberty 3, 5
 - progesterone
 - direct effects of GnRH agonists 74
 - GnRH antagonists suppressing 15, 87, 88
 - progestins 66, 92
 - prolactin
 - mammary tumours 13, 129
 - prostate cancer 11–12

- prolactin release-inhibiting factor (PIF) 13
- prostaglandins 62–3, 66
- prostate cancer 2, 6, 11–12
 - GnRH agonist therapy 11–12, 78, 144, 149–53
 - GnRH receptors 72, 75, 78
- prostate gland 7, 75
- prostatic hyperplasia, benign (BPH) 2, 5, 12
- protein binding, plasma 115
- protein kinase C (PKC) 54, 60, 65, 66, 74, 130
- puberty, precocious 3, 5
- pulmonary function, effects of GnRH antagonists 109–11
- radiation, GnRH analogues for gonadal protection 17
- rectal administration 144
- reduced size GnRH analogues 1, 45–51
 - measurement of biological activity 45–6
 - structure–activity relationships 46–51
 - synthesis 45
- rhinitis, intranasal administration and 144–5
- RS 68439 (Detirelix) 15–16, 38
- secretin 14
- seminal vesicles 7, 75
- Sertoli cells 71, 73
- slow release formulations, see sustained release formulations
- somatomedins 12
- somatostatin analogues 2
 - breast cancer 13
 - pancreatic cancer 13–14
 - prostate cancer 11–12
- spermatogenesis, suppression of 8, 9, 15
- steroidogenesis, inhibition of gonadal 7, 72, 73, 74
- structure–activity relationships
 - agonistic GnRH analogues 33–6
 - antagonistic GnRH analogues 15, 36–40
 - reduced size (3–9) GnRH analogues 46–51
- subcutaneous administration 143
- sustained release formulations (depot formulations) 2, 40, 137–9, 145–54
 - assays of biological effects 145–9
 - in breast cancer 13, 149, 150
 - GnRH antagonists 17
 - pharmacokinetics 120–3, 138–9
 - in prostate cancer 11, 149–53
- testes
 - direct effects of GnRH 71, 72–4
 - protection against chemotherapy and radiation damage 17
- testosterone
 - direct suppression by LHRH 73
 - effects of GnRH antagonists 15
 - GnRH agonists suppressing 7, 139, 148–9, 151
- 12-O-tetradecanoyl-phorbol-13-acetate (TPA) 60
- theca cells 74
- total androgen blockade 2, 151
- TPA (12-O-tetradecanoyl-phorbol-13-acetate) 60
- transdermal administration 144
- transforming growth factor (TGF) 12
- tryptorelin ([D-Trp⁶]-LHRH, decapeptyl) 35
 - aqueous solution 143
 - breast cancer 12–13
 - gonadal protection against chemotherapy and radiation 17
 - gynecological disorders 9, 10
 - metabolism 115
 - ovarian cancer 10
 - pancreatic cancer 13–14
 - prostate cancer 11, 78
 - stimulating phospholipase C in mammary tumour membranes 129–30, 131, 132
 - structure 6, 34
 - sustained release formulations 137, 138–9, 145
- tyrosine kinase, see protein kinase C
- urinary excretion, GnRH agonists 113, 116–17
- uterine leiomyomas, see leiomyomas, uterine
- uterus, direct effects of GnRH 75
- wheel response 108–9
- Zoladex, see goserelin

WK 515 G572 1989-1990
GnRH analogues in cancer and human
Countway Library
AJH1543



3 2044 016 369 910

DUE DATE

[illegible]

X 515 G572 v.1 1989

S 2044 016 369 910

HD